



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:48 PM UTC

PDB ID : 5K6C / pdb_00005k6c
Title : Crystal structure of prefusion-stabilized RSV F single-chain 9-10 DS-Cav1 variant.
Authors : Joyce, M.G.; Zhang, B.; Lai, Y.T.; Mascola, J.R.; Kwong, P.D.
Deposited on : 2016-05-24
Resolution : 3.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1521 (3.66-3.50)
Clashscore	190562	1595 (3.66-3.50)
Ramachandran outliers	187476	1551 (3.66-3.50)
Sidechain outliers	187428	1551 (3.66-3.50)
RSRZ outliers	180081	1520 (3.66-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	F	445	 81% 16% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fusion glycoprotein F0,Fusion glycoprotein F0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	F	445	Total	C	N	O	S	0	0	0
			3442	2165	573	681	23			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	102	ALA	PRO	engineered mutation	UNP P03420
F	143	GLY	-	linker	UNP P03420
F	144	SER	-	linker	UNP P03420
F	155	CYS	SER	engineered mutation	UNP P03420
F	190	PHE	SER	engineered mutation	UNP P03420
F	207	LEU	VAL	engineered mutation	UNP P03420
F	290	CYS	SER	engineered mutation	UNP P03420
F	373	ARG	LEU	engineered mutation	UNP P03420
F	379	VAL	ILE	engineered mutation	UNP P03420
F	447	VAL	MET	engineered mutation	UNP P03420

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	F	1	Total	O	S	0	0
			5	4	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fusion glycoprotein F0,Fusion glycoprotein F0



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	171.22Å 171.22Å 171.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.43 – 3.58 49.43 – 3.58	Depositor EDS
% Data completeness (in resolution range)	91.7 (49.43-3.58) 85.0 (49.43-3.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.79 (at 3.57Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.227 , 0.262 0.230 , 0.275	Depositor DCC
R_{free} test set	465 reflections (4.36%)	wwPDB-VP
Wilson B-factor (Å ²)	114.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 100.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3447	wwPDB-VP
Average B, all atoms (Å ²)	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	F	0.24	1/3492 (0.0%)	0.43	5/4732 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	265	PRO	N-CD	5.21	1.55	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	264	MET	CA-C-N	-6.11	112.84	120.79
1	F	264	MET	C-N-CA	-6.11	112.84	120.79
1	F	290	CYS	N-CA-C	5.50	120.00	112.68
1	F	361	GLN	N-CA-C	-5.18	102.01	110.20
1	F	162	GLY	N-CA-C	-5.14	108.53	115.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3442	0	3478	76	0
2	F	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3447	0	3478	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:379:VAL:HG23	1:F:391:TYR:HE2	1.14	1.11
1:F:161:GLU:OE1	1:F:162:GLY:N	1.94	1.00
1:F:379:VAL:HG23	1:F:391:TYR:CE2	2.00	0.97
1:F:391:TYR:CD1	1:F:392:ASP:N	2.35	0.95
1:F:310:ASP:OD1	1:F:364:ARG:NH2	2.00	0.94
1:F:143:GLY:HA2	1:F:144:SER:HB3	1.49	0.94
1:F:391:TYR:HD1	1:F:392:ASP:H	1.14	0.91
1:F:237:PHE:CE2	1:F:289:MET:HB2	2.11	0.85
1:F:265:PRO:O	1:F:266:ILE:HG23	1.76	0.82
1:F:259:SER:O	1:F:263:ASP:OD1	2.02	0.77
1:F:290:CYS:O	1:F:291:ILE:HG23	1.85	0.76
1:F:361:GLN:OE1	1:F:362:SER:N	2.19	0.75
1:F:379:VAL:HA	1:F:391:TYR:HD2	1.50	0.74
1:F:291:ILE:O	1:F:291:ILE:HD12	1.89	0.73
1:F:388:ASN:HD22	1:F:388:ASN:N	1.93	0.67
1:F:362:SER:O	1:F:363:ASN:HB3	1.95	0.66
1:F:379:VAL:CG2	1:F:391:TYR:HE2	2.02	0.65
1:F:143:GLY:CA	1:F:144:SER:HB3	2.27	0.62
1:F:379:VAL:HA	1:F:391:TYR:CD2	2.33	0.62
1:F:144:SER:OG	1:F:145:GLY:N	2.26	0.62
1:F:289:MET:HA	1:F:299:TYR:CB	2.30	0.61
1:F:364:ARG:CG	1:F:364:ARG:HH11	2.14	0.60
1:F:143:GLY:HA2	1:F:144:SER:CB	2.29	0.59
1:F:291:ILE:HD12	1:F:291:ILE:C	2.27	0.59
1:F:237:PHE:CZ	1:F:289:MET:HB2	2.37	0.58
1:F:391:TYR:HD1	1:F:392:ASP:N	1.87	0.58
1:F:61:LEU:HD11	1:F:297:LEU:HD23	1.86	0.56
1:F:93:LEU:HD23	1:F:292:ILE:HD11	1.87	0.56
1:F:290:CYS:O	1:F:291:ILE:CG2	2.54	0.55
1:F:167:ILE:HD12	1:F:189:THR:HG21	1.89	0.55
1:F:206:ILE:HG23	1:F:213:SER:HB3	1.89	0.55
1:F:407:ILE:HD11	1:F:457:TYR:HB3	1.88	0.54
1:F:151:GLY:O	1:F:155:CYS:SG	2.65	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:359:LYS:C	1:F:360:VAL:HG23	2.31	0.54
1:F:338:ASP:O	1:F:342:TYR:OH	2.22	0.52
1:F:75:LYS:HE3	1:F:217:ILE:HB	1.92	0.51
1:F:265:PRO:O	1:F:266:ILE:CG2	2.54	0.51
1:F:263:ASP:OD1	1:F:263:ASP:N	2.43	0.51
1:F:364:ARG:NH1	1:F:364:ARG:HG3	2.25	0.51
1:F:60:GLU:HB2	1:F:191:LYS:HD2	1.92	0.50
1:F:76:VAL:HG11	1:F:212:CYS:HB3	1.94	0.50
1:F:59:ILE:HB	1:F:297:LEU:HG	1.92	0.49
1:F:86:TYR:HB2	1:F:227:ASN:HD21	1.76	0.49
1:F:291:ILE:C	1:F:291:ILE:CD1	2.85	0.49
1:F:266:ILE:HG13	1:F:271:LYS:HG3	1.95	0.49
1:F:359:LYS:C	1:F:360:VAL:CG2	2.85	0.48
1:F:426:ASN:HB3	1:F:429:ARG:HG2	1.95	0.48
1:F:265:PRO:O	1:F:265:PRO:HG2	2.14	0.47
1:F:354:GLN:O	1:F:357:THR:OG1	2.32	0.47
1:F:289:MET:HA	1:F:299:TYR:HB2	1.95	0.47
1:F:293:LYS:HD3	1:F:293:LYS:HA	1.69	0.47
1:F:288:ILE:O	1:F:300:VAL:N	2.48	0.47
1:F:318:THR:HG23	1:F:339:ARG:HB3	1.96	0.46
1:F:273:LEU:HD11	1:F:364:ARG:HD3	1.96	0.46
1:F:388:ASN:HD22	1:F:388:ASN:H	1.60	0.46
1:F:290:CYS:C	1:F:291:ILE:HG23	2.39	0.46
1:F:143:GLY:CA	1:F:144:SER:CB	2.90	0.45
1:F:157:VAL:HG21	1:F:183:ASN:HB2	1.99	0.45
1:F:77:LYS:O	1:F:81:GLN:HG2	2.16	0.45
1:F:143:GLY:HA3	1:F:144:SER:C	2.43	0.43
1:F:53:TYR:CD2	1:F:264:MET:HE2	2.52	0.43
1:F:60:GLU:O	1:F:196:LYS:N	2.51	0.43
1:F:388:ASN:HA	1:F:389:PRO:HD3	1.86	0.43
1:F:143:GLY:HA3	1:F:145:GLY:N	2.33	0.43
1:F:146:SER:OG	1:F:370:MET:SD	2.77	0.42
1:F:394:LYS:HA	1:F:491:SER:HA	2.02	0.42
1:F:175:ASN:OD1	1:F:175:ASN:N	2.49	0.42
1:F:252:LEU:HD12	1:F:252:LEU:HA	1.89	0.42
1:F:58:THR:HG23	1:F:296:VAL:HG13	2.01	0.42
1:F:30:GLU:CD	1:F:410:LEU:HB2	2.45	0.41
1:F:289:MET:HE1	1:F:297:LEU:HB2	2.02	0.41
1:F:278:VAL:O	1:F:282:ARG:HG3	2.21	0.41
1:F:235:ARG:HG3	1:F:236:GLU:N	2.36	0.40
1:F:56:VAL:HB	1:F:189:THR:HG22	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:388:ASN:H	1:F:388:ASN:ND2	2.18	0.40
1:F:93:LEU:HG	1:F:289:MET:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	F	443/445 (100%)	421 (95%)	21 (5%)	1 (0%)	43 72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	360	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	F	406/406 (100%)	394 (97%)	12 (3%)	36 60

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	160	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	161	GLU
1	F	263	ASP
1	F	265	PRO
1	F	289	MET
1	F	359	LYS
1	F	361	GLN
1	F	364	ARG
1	F	388	ASN
1	F	390	LYS
1	F	429	ARG
1	F	460	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	27	ASN
1	F	388	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	F	601	-	4,4,4	0.35	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	F	445/445 (100%)	-0.08	2 (0%) 88 70	65, 114, 193, 268	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	102	ALA	2.1
1	F	392	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	F	601	5/5	0.67	0.16	248,260,272,278	0

6.5 Other polymers [i](#)

There are no such residues in this entry.