



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:30 PM UTC

PDB ID : 5K6E / pdb_00005k6e
Title : Crystal structure of Retinoic acid receptor-related orphan receptor (ROR) gamma ligand binding domain complex with SBI0654919
Authors : Lu, J.; Rastinejad, F.
Deposited on : 2016-05-24
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	4% 77% 13% 9%
1	B	247	7% 79% 13% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3841 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nuclear receptor ROR-gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1850	1171	336	329	14			
1	B	234	Total	C	N	O	S	0	0	0
			1909	1212	345	338	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	261	GLY	-	expression tag	UNP P51449
A	262	SER	-	expression tag	UNP P51449
A	263	HIS	-	expression tag	UNP P51449
A	264	MET	-	expression tag	UNP P51449
B	261	GLY	-	expression tag	UNP P51449
B	262	SER	-	expression tag	UNP P51449
B	263	HIS	-	expression tag	UNP P51449
B	264	MET	-	expression tag	UNP P51449

- Molecule 2 is 2-chloranyl-6-fluoranyl- {N}-[(2 {R})-4-(4-fluorophenyl)sulfonyl-2-(hydroxymethyl)-2,3-dihydro-1,4-benzoxazin-6-yl]benzamide (CCD ID: 6QP) (formula: C₂₂H₁₇ClF₂N₂O₅S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	S	0	0
			33	22	1	2	2	5	1		
2	B	1	Total	C	Cl	F	N	O	S	0	0
			33	22	1	2	2	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total	O	0	0
			6	6		
3	B	10	Total	O	0	0
			10	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nuclear receptor ROR-gamma



• Molecule 1: Nuclear receptor ROR-gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	117.67Å 139.93Å 120.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.16 – 2.80 42.16 – 2.80	Depositor EDS
% Data completeness (in resolution range)	80.6 (42.16-2.80) 81.5 (42.16-2.80)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.42 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.209 , 0.259 0.215 , 0.261	Depositor DCC
R_{free} test set	1043 reflections (4.12%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3841	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6QP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	1/1887 (0.1%)	0.90	5/2537 (0.2%)
1	B	0.56	2/1948 (0.1%)	0.90	6/2622 (0.2%)
All	All	0.56	3/3835 (0.1%)	0.90	11/5159 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	354	LYS	C-O	5.85	1.31	1.24
1	B	351	VAL	C-O	5.65	1.30	1.24
1	B	354	LYS	C-O	5.41	1.30	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	470	GLY	N-CA-C	-7.77	104.58	114.37
1	A	357	ALA	CA-C-N	-7.16	109.42	120.31
1	A	357	ALA	C-N-CA	-7.16	109.42	120.31
1	B	357	ALA	CA-C-N	-6.68	110.16	120.31
1	B	357	ALA	C-N-CA	-6.68	110.16	120.31
1	A	471	LYS	N-CA-C	-5.81	104.54	111.69
1	B	426	ILE	N-CA-C	5.51	112.10	106.21
1	B	490	HIS	CA-C-N	-5.48	112.99	119.84
1	B	490	HIS	C-N-CA	-5.48	112.99	119.84
1	A	269	GLU	N-CA-C	-5.42	106.51	113.23
1	B	428	ALA	N-CA-C	5.34	119.67	113.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1850	0	1849	22	0
1	B	1909	0	1912	25	0
2	A	33	0	0	0	0
2	B	33	0	0	1	0
3	A	6	0	0	1	0
3	B	10	0	0	0	0
All	All	3841	0	3761	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:427:ASN:HD22	1:B:427:ASN:C	1.84	0.85
1:B:427:ASN:ND2	1:B:429:HIS:H	1.78	0.82
1:A:452:HIS:NE2	3:A:701:HOH:O	2.13	0.79
1:A:363:VAL:O	1:A:366:CYS:HB3	1.91	0.71
1:A:392:GLY:HA2	1:B:283:GLU:HB3	1.76	0.66
1:B:407:LEU:HD22	1:B:412:PHE:HE2	1.65	0.62
1:B:388:PHE:HB3	1:B:391:LEU:HD12	1.85	0.59
1:B:363:VAL:O	1:B:366:CYS:HB3	2.04	0.58
1:A:470:GLY:HA2	1:A:473:ARG:HG2	1.84	0.58
1:B:427:ASN:C	1:B:427:ASN:ND2	2.56	0.58
1:B:427:ASN:HD22	1:B:428:ALA:N	2.03	0.56
1:A:303:GLU:OE1	1:A:303:GLU:N	2.31	0.55
1:A:470:GLY:HA2	1:A:473:ARG:HB2	1.89	0.54
1:A:392:GLY:CA	1:B:283:GLU:HB3	2.38	0.54
1:A:362:LEU:O	1:A:365:MET:HG2	2.09	0.53
1:A:396:LEU:HD13	1:A:478:GLN:HB3	1.91	0.53
1:B:427:ASN:HD22	1:B:429:HIS:H	1.53	0.52
1:B:356:GLY:HA3	1:B:425:LEU:HD23	1.90	0.52
1:A:412:PHE:HB3	1:A:417:ILE:HG13	1.92	0.51
1:B:490:HIS:N	1:B:491:PRO:HD3	2.26	0.51
1:A:288:ARG:NH2	1:A:291:ASP:OD2	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:427:ASN:HD21	1:B:429:HIS:H	1.59	0.49
1:A:346:GLN:O	1:A:350:ILE:HG12	2.13	0.49
1:A:267:LEU:O	1:A:268:THR:HB	2.13	0.49
1:B:486:PHE:O	1:B:490:HIS:HB2	2.13	0.48
1:A:469:LYS:HE2	1:A:473:ARG:NH1	2.28	0.48
1:B:309:GLN:NE2	1:B:386:GLU:HB2	2.29	0.47
1:B:492:ILE:O	1:B:492:ILE:HG13	2.16	0.45
1:B:360:VAL:HG21	1:B:425:LEU:HG	1.98	0.44
1:A:488:HIS:CD2	1:B:342:MET:HE3	2.53	0.44
1:B:306:THR:HA	1:B:309:GLN:HB2	2.00	0.44
1:A:281:TYR:OH	1:A:367:ARG:NE	2.51	0.44
1:B:459:ARG:O	1:B:462:ILE:HG12	2.18	0.43
1:B:396:LEU:HD13	1:B:478:GLN:HG2	2.00	0.43
1:A:266:SER:OG	1:A:267:LEU:N	2.52	0.42
1:A:359:GLU:HB3	1:A:472:LEU:HD21	2.02	0.42
1:B:353:LEU:HD23	1:B:353:LEU:HA	1.88	0.42
1:A:364:ARG:O	1:A:367:ARG:HB2	2.20	0.41
1:A:469:LYS:HE2	1:A:473:ARG:CZ	2.50	0.41
1:B:393:CYS:O	1:B:397:ILE:HG12	2.20	0.41
1:A:316:MET:HG2	1:A:390:ALA:HB3	2.03	0.41
1:B:365:MET:HE2	2:B:601:6QP:C17	2.51	0.41
1:A:289:LEU:O	1:A:293:LEU:HG	2.20	0.40
1:B:437:ARG:HG3	1:B:437:ARG:HH11	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/247 (91%)	215 (96%)	7 (3%)	2 (1%)	14	41
1	B	232/247 (94%)	219 (94%)	10 (4%)	3 (1%)	9	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	456/494 (92%)	434 (95%)	17 (4%)	5 (1%)	11	36

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	494	VAL
1	B	497	ALA
1	A	354	LYS
1	A	488	HIS
1	B	354	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/220 (92%)	197 (98%)	5 (2%)	42	76
1	B	208/220 (94%)	203 (98%)	5 (2%)	43	77
All	All	410/440 (93%)	400 (98%)	10 (2%)	43	77

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	THR
1	A	275	GLN
1	A	366	CYS
1	A	473	ARG
1	A	490	HIS
1	B	366	CYS
1	B	395	GLU
1	B	427	ASN
1	B	429	HIS
1	B	469	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	370	ASN
1	A	441	GLN
1	A	453	HIS
1	B	309	GLN
1	B	427	ASN
1	B	429	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	6QP	B	601	-	36,36,36	2.85	10 (27%)	50,53,53	3.42	19 (38%)
2	6QP	A	601	-	36,36,36	2.74	10 (27%)	50,53,53	3.53	21 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6QP	B	601	-	-	2/22/34/34	0/4/4/4
2	6QP	A	601	-	-	5/22/34/34	0/4/4/4

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	6QP	C05-C04	-7.69	1.26	1.39
2	A	601	6QP	C05-C04	-7.10	1.27	1.39
2	B	601	6QP	C24-C07	6.62	1.53	1.39
2	B	601	6QP	C06-N13	-6.51	1.36	1.43
2	A	601	6QP	C24-C07	6.40	1.52	1.39
2	B	601	6QP	C12-N13	-6.38	1.40	1.48
2	A	601	6QP	C12-N13	-6.25	1.40	1.48
2	B	601	6QP	C25-C04	-5.53	1.29	1.39
2	A	601	6QP	C06-N13	-5.52	1.37	1.43
2	A	601	6QP	C25-C04	-5.36	1.30	1.39
2	A	601	6QP	C25-C24	4.07	1.45	1.38
2	B	601	6QP	C25-C24	4.07	1.45	1.38
2	B	601	6QP	O08-C07	3.96	1.43	1.38
2	A	601	6QP	O08-C07	3.83	1.43	1.38
2	B	601	6QP	C05-C06	-3.46	1.34	1.39
2	A	601	6QP	C05-C06	-3.22	1.34	1.39
2	A	601	6QP	S14-N13	3.15	1.69	1.65
2	A	601	6QP	C02-N03	3.06	1.44	1.35
2	B	601	6QP	C02-N03	2.89	1.44	1.35
2	B	601	6QP	S14-N13	2.02	1.68	1.65

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	6QP	O23-S14-O22	-11.47	101.69	119.59
2	A	601	6QP	O23-S14-O22	-11.37	101.84	119.59
2	A	601	6QP	C24-C25-C04	-9.09	109.83	120.30
2	B	601	6QP	C24-C07-C06	-8.95	105.04	120.32
2	B	601	6QP	C24-C25-C04	-8.78	110.19	120.30
2	A	601	6QP	C24-C07-C06	-8.47	105.86	120.32
2	A	601	6QP	O08-C07-C06	7.80	134.57	122.03
2	B	601	6QP	O08-C07-C06	7.63	134.29	122.03
2	A	601	6QP	O22-S14-N13	6.44	114.19	106.69
2	A	601	6QP	C25-C04-C05	6.17	127.12	119.66
2	B	601	6QP	C25-C04-C05	6.07	126.99	119.66
2	A	601	6QP	C07-C06-N13	-5.92	112.42	118.99
2	B	601	6QP	C07-C06-N13	-5.53	112.85	118.99

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	6QP	O23-S14-N13	5.51	113.11	106.69
2	B	601	6QP	O23-S14-N13	5.04	112.56	106.69
2	B	601	6QP	O22-S14-N13	4.93	112.43	106.69
2	A	601	6QP	C25-C24-C07	4.66	128.88	120.06
2	B	601	6QP	C25-C24-C07	4.55	128.67	120.06
2	A	601	6QP	C04-C05-C06	3.25	125.01	119.48
2	B	601	6QP	C15-S14-N13	3.24	111.20	106.97
2	B	601	6QP	C04-C05-C06	3.14	124.83	119.48
2	B	601	6QP	C05-C06-C07	3.06	124.27	119.78
2	A	601	6QP	O08-C09-C10	2.91	111.28	106.79
2	A	601	6QP	C15-S14-N13	2.90	110.75	106.97
2	A	601	6QP	C25-C04-N03	-2.57	111.80	120.41
2	B	601	6QP	C32-C26-C02	-2.56	118.33	122.21
2	A	601	6QP	C26-C02-N03	2.55	119.49	114.81
2	B	601	6QP	O08-C07-C24	2.45	120.67	116.20
2	A	601	6QP	C05-C06-C07	2.39	123.30	119.78
2	A	601	6QP	C21-C20-C18	2.38	120.82	118.38
2	B	601	6QP	C25-C04-N03	-2.37	112.47	120.41
2	B	601	6QP	O08-C09-C10	2.35	110.42	106.79
2	B	601	6QP	C07-O08-C09	-2.34	109.52	114.61
2	A	601	6QP	C12-N13-S14	-2.30	112.07	117.37
2	B	601	6QP	C26-C02-N03	2.29	119.01	114.81
2	A	601	6QP	C07-O08-C09	-2.23	109.75	114.61
2	A	601	6QP	O01-C02-C26	-2.11	117.95	121.06
2	B	601	6QP	C16-C17-C18	2.08	120.52	118.38
2	A	601	6QP	C31-C32-C26	-2.07	119.69	123.48
2	A	601	6QP	F33-C32-C26	2.01	121.67	118.15

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	6QP	O08-C09-C10-O11
2	A	601	6QP	C12-C09-C10-O11
2	A	601	6QP	C06-N13-S14-O22
2	B	601	6QP	C06-N13-S14-O22
2	A	601	6QP	C06-N13-S14-C15
2	B	601	6QP	C06-N13-S14-C15
2	A	601	6QP	C12-N13-S14-O22


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	6QP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/247 (91%)	0.24	10 (4%) 39 31	15, 32, 59, 92	0
1	B	234/247 (94%)	0.49	17 (7%) 21 15	16, 38, 64, 78	0
All	All	460/494 (93%)	0.37	27 (5%) 28 21	15, 35, 63, 92	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	498	PHE	6.9
1	B	494	VAL	6.7
1	B	265	ALA	4.9
1	B	266	SER	4.3
1	A	266	SER	4.1
1	A	265	ALA	3.8
1	B	489	LEU	3.7
1	B	355	ALA	3.6
1	B	288	ARG	3.5
1	B	493	VAL	3.4
1	B	354	LYS	3.4
1	A	267	LEU	3.2
1	B	339	SER	3.1
1	A	469	LYS	3.1
1	B	497	ALA	3.1
1	B	356	GLY	3.1
1	B	492	ILE	2.9
1	A	355	ALA	2.8
1	B	496	ALA	2.7
1	A	471	LYS	2.6
1	A	354	LYS	2.5
1	A	470	GLY	2.4
1	A	275	GLN	2.4
1	B	469	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	488	HIS	2.3
1	B	294	ARG	2.2
1	A	452	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	6QP	B	601	33/33	0.85	0.17	29,47,75,125	0
2	6QP	A	601	33/33	0.88	0.14	22,33,61,111	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 6QP B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 6QP A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.