



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:01 PM UTC

PDB ID : 7K75 / pdb_00007k75
Title : Crystal structure of MAD2-6 IgA Fab in complex with PfCSP N-terminal peptide.
Authors : Pholcharee, T.; Wilson, I.A.
Deposited on : 2020-09-22
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.



■ Percentile relative to all X-ray structures

Percentile relative to X-ray structures of similar resolution

MetricWhole archive
(#Entries)

Similar resolution
(#Entries, resolution range(Å))

 R_{free}

Clashscore

Ramachandran outliers

Sidechain outliers

RSRZ outliers

180053

190562

187476

187428

180081

5829 (2.50-2.50)

6492 (2.50-2.50)

6378 (2.50-2.50)

6380 (2.50-2.50)

5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Quality of chain

2%

80%

20%

2%

80%

20%

16%

6%

0%

3%

78%

0%

85%

15%

Continued on next page...

Mol	Chain	Length	Quality of chain
2	D	213	89%11%
2	J	213	85%15%
2	N	213	2%86%14%
3	E	8	12%88%12%
3	F	8	100%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13557 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of MAD2-6 IgA Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	225	Total	C	N	O	S	0	2	0
			1701	1078	282	333	8			
1	C	225	Total	C	N	O	S	0	0	0
			1683	1069	277	329	8			
1	I	222	Total	C	N	O	S	0	0	0
			1578	1007	260	303	8			
1	M	222	Total	C	N	O	S	0	1	0
			1647	1048	274	317	8			

- Molecule 2 is a protein called Light chain of MAD2-6 IgA Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	213	Total	C	N	O	S	0	0	0
			1619	1009	274	329	7			
2	D	213	Total	C	N	O	S	0	1	0
			1621	1011	274	329	7			
2	J	213	Total	C	N	O	S	0	1	0
			1615	1006	272	330	7			
2	N	213	Total	C	N	O	S	0	1	0
			1612	1003	271	331	7			

- Molecule 3 is a protein called PfCSP N-terminal peptide P17.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	8	Total	C	N	O	0	0	0
			66	44	14	8			
3	F	8	Total	C	N	O	0	0	0
			66	44	14	8			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	73	Total 73	O 73	0	0
4	B	50	Total 50	O 50	0	0
4	D	46	Total 46	O 46	0	0
4	C	76	Total 76	O 76	0	0
4	I	6	Total 6	O 6	0	0
4	J	43	Total 43	O 43	0	0
4	M	26	Total 26	O 26	0	0
4	N	25	Total 25	O 25	0	0
4	E	3	Total 3	O 3	0	0
4	F	1	Total 1	O 1	0	0

- Molecule 1: Heavy chain of MAD2-6 IgA Fab







- Molecule 2: Light chain of MAD2-6 IgA Fab




- Molecule 2: Light chain of MAD2-6 IgA Fab



- Molecule 2: Light chain of MAD2-6 IgA Fab



- Molecule 2: Light chain of MAD2-6 IgA Fab



- Molecule 3: PfCSP N-terminal peptide P17

Chain E:  12% 88% 12%



- Molecule 3: PfCSP N-terminal peptide P17

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.14Å 153.68Å 193.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.96 – 2.50 41.96 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.8 (41.96-2.50) 98.9 (41.96-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.19_4092: ???)	Depositor
R, R_{free}	0.192 , 0.236 0.192 , 0.235	Depositor DCC
R_{free} test set	3739 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13557	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	J	1615	0	1512	29	0
2	N	1612	0	1505	17	0
3	E	66	0	81	1	0
3	F	66	0	81	0	0
4	A	73	0	0	2	0
4	B	50	0	0	1	0
4	C	76	0	0	0	0
4	D	46	0	0	0	0
4	E	3	0	0	0	0
4	F	1	0	0	0	0
4	I	6	0	0	0	0
4	J	43	0	0	0	0
4	M	26	0	0	0	0
4	N	25	0	0	0	0
All	All	13557	0	12505	176	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 176 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2:ILE:HD12	2:B:93:GLU:HB2	1.54	0.90
2:J:189:LYS:HE3	2:J:209:ASN:HB3	1.56	0.88
2:J:186:GLU:HA	2:J:210:ARG:HH11	1.42	0.83
2:B:15:VAL:HG11	1:C:9:PRO:HB3	1.60	0.82
1:M:92:THR:HG23	1:M:120:THR:HA	1.66	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/225 (100%)	215 (96%)	10 (4%)	0	100	100
1	C	223/225 (99%)	214 (96%)	9 (4%)	0	100	100
1	I	218/225 (97%)	201 (92%)	16 (7%)	1 (0%)	24	43
1	M	219/225 (97%)	209 (95%)	10 (5%)	0	100	100
2	B	211/213 (99%)	202 (96%)	9 (4%)	0	100	100
2	D	212/213 (100%)	205 (97%)	7 (3%)	0	100	100
2	J	212/213 (100%)	205 (97%)	7 (3%)	0	100	100
2	N	212/213 (100%)	201 (95%)	11 (5%)	0	100	100
3	E	6/8 (75%)	6 (100%)	0	0	100	100
3	F	6/8 (75%)	6 (100%)	0	0	100	100
All	All	1744/1768 (99%)	1664 (95%)	79 (4%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	9	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/194 (98%)	187 (98%)	4 (2%)	47	74
1	C	189/194 (97%)	183 (97%)	6 (3%)	34	62
1	I	160/194 (82%)	154 (96%)	6 (4%)	29	56
1	M	181/194 (93%)	172 (95%)	9 (5%)	22	44
2	B	181/187 (97%)	179 (99%)	2 (1%)	65	84
2	D	180/187 (96%)	177 (98%)	3 (2%)	53	78
2	J	178/187 (95%)	176 (99%)	2 (1%)	65	84
2	N	178/187 (95%)	175 (98%)	3 (2%)	53	78
3	E	7/8 (88%)	7 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	F	7/8 (88%)	7 (100%)	0	100	100
All	All	1452/1540 (94%)	1417 (98%)	35 (2%)	43	70

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	81	SER
1	M	165	SER
2	N	72	THR
1	C	201	LEU
1	C	200	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	152	GLN
1	M	218	GLN
1	I	212	HIS
3	E	4	HIS
1	M	144	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/225 (100%)	-0.12	4 (1%) 67 64	18, 40, 61, 92	2 (0%)
1	C	225/225 (100%)	-0.20	4 (1%) 67 64	26, 38, 67, 88	0
1	I	222/225 (98%)	1.11	37 (16%) 4 3	40, 78, 102, 123	0
1	M	222/225 (98%)	0.53	7 (3%) 50 46	28, 63, 84, 106	1 (0%)
2	B	213/213 (100%)	-0.30	0 100 100	27, 38, 56, 68	0
2	D	213/213 (100%)	-0.18	0 100 100	23, 38, 62, 74	1 (0%)
2	J	213/213 (100%)	-0.00	1 (0%) 87 85	22, 44, 82, 90	1 (0%)
2	N	213/213 (100%)	0.26	4 (1%) 66 62	30, 51, 79, 101	1 (0%)
3	E	8/8 (100%)	0.07	1 (12%) 8 6	38, 44, 53, 67	0
3	F	8/8 (100%)	-0.18	0 100 100	36, 39, 51, 60	0
All	All	1762/1768 (99%)	0.14	58 (3%) 49 45	18, 46, 86, 123	6 (0%)

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	N	213	CYS	4.6
1	I	18	LEU	4.6
1	I	121	VAL	4.2
1	I	90	ALA	4.0
1	I	139	THR	3.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.