



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:16 PM UTC

PDB ID : 4K9E / pdb_00004k9e
Title : Crystal structure of KIT D4D5 fragment in complex with anti-Kit antibodies Fab79D
Authors : Resheynyak, A.V.; Boggon, T.J.; Lax, I.; Schlessinger, J.
Deposited on : 2013-04-19
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	228	
2	H	221	
3	C	214	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	221	Total	C	N	O	S	0	0	0
			1686	1054	291	336	5			

- Molecule 2 is a protein called heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	0	0
			1649	1047	272	323	7			


- Molecule 3 is a protein called Mast/stem cell growth factor receptor Kit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	193	Total	C	N	O	S	0	0	0
			1521	974	244	296	7			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	305	GLY	-	expression tag	UNP P10721
C	306	ALA	-	expression tag	UNP P10721
C	307	MET	-	expression tag	UNP P10721

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	24	Total	O	0	0
			24	24		
5	H	26	Total	O	0	0
			26	26		
5	C	29	Total	O	0	0
			29	29		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: light chain



- Molecule 2: heavy chain



- Molecule 3: Mast/stem cell growth factor receptor Kit





4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	94.58Å 94.58Å 320.88Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.78 – 2.70 44.78 – 2.70	Depositor EDS
% Data completeness (in resolution range)	90.8 (44.78-2.70) 90.8 (44.78-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	21.47 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.247 , 0.282 0.247 , 0.283	Depositor DCC
R_{free} test set	1138 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4949	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.31	0/1720	0.75	0/2334
2	H	0.27	0/1692	0.72	0/2309
3	C	0.30	0/1558	0.75	0/2122
All	All	0.29	0/4970	0.74	0/6765

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1686	0	1637	33	0
2	H	1649	0	1610	21	0
3	C	1521	0	1428	34	0
4	C	14	0	13	0	0
5	C	29	0	0	9	0
5	H	26	0	0	2	0
5	L	24	0	0	4	0
All	All	4949	0	4688	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:127:GLN:NE2	5:L:301:HOH:O	1.99	0.94
3:C:352:ASN:ND2	5:C:728:HOH:O	2.08	0.84
1:L:211:SER:OG	5:L:308:HOH:O	1.96	0.82
3:C:389:THR:O	5:C:711:HOH:O	1.97	0.81
2:H:0:SER:N	5:H:302:HOH:O	2.20	0.74
1:L:111:ARG:NH1	1:L:112:THR:O	2.22	0.73
1:L:192:HIS:O	1:L:214:ARG:NH1	2.21	0.72
3:C:439:ASP:OD1	5:C:718:HOH:O	2.08	0.71
3:C:408:TYR:OH	5:C:717:HOH:O	1.90	0.69
3:C:379:LEU:HB3	3:C:382:LEU:HD11	1.77	0.67
2:H:134:SER:OG	2:H:141:ALA:O	2.13	0.66
3:C:458:ASP:OD2	5:C:715:HOH:O	2.14	0.65
3:C:360:GLU:OE1	3:C:378:HIS:NE2	2.24	0.65
2:H:131:SER:OG	2:H:132:SER:N	2.30	0.64
1:L:108:GLU:OE2	1:L:176:TYR:OH	2.16	0.64
1:L:17:ASP:OD1	1:L:18:ARG:N	2.29	0.61
3:C:360:GLU:CD	3:C:378:HIS:HE2	2.09	0.61
3:C:455:LEU:HD12	3:C:456:PRO:HD2	1.83	0.60
1:L:102:GLY:O	5:L:323:HOH:O	2.17	0.59
1:L:38:TYR:HE1	1:L:91:GLN:HG2	1.67	0.59
2:H:26:GLY:O	3:C:326:ASN:ND2	2.35	0.58
2:H:67:ARG:NH2	2:H:90:ASP:OD2	2.36	0.58
3:C:385:THR:OG1	3:C:386:GLU:N	2.37	0.57
3:C:432:GLY:HA2	3:C:471:LYS:HD2	1.86	0.57
1:L:39:GLN:HB2	1:L:49:LEU:HD11	1.86	0.56
1:L:148:LYS:HB3	1:L:200:THR:HG23	1.87	0.56
2:H:83:MET:HB3	2:H:86:LEU:HD21	1.87	0.55
3:C:406:ASN:ND2	5:C:704:HOH:O	2.31	0.55
2:H:63:SER:O	2:H:67:ARG:NH1	2.39	0.55
1:L:6:GLN:HE21	1:L:21:ILE:HD11	1.72	0.55
1:L:109:ILE:HG13	1:L:169:GLN:OE1	2.08	0.53
1:L:142:PHE:O	1:L:175:THR:OG1	2.26	0.53
1:L:78:SER:OG	5:L:311:HOH:O	2.18	0.53
3:C:413:PRO:HB3	3:C:436:PRO:HB3	1.90	0.53
2:H:91:THR:HG23	2:H:114:THR:HA	1.91	0.53
1:L:201:HIS:CD2	1:L:203:GLY:H	2.26	0.52
2:H:102:HIS:HE2	3:C:360:GLU:CD	2.17	0.52
3:C:456:PRO:HA	3:C:478:ILE:HG22	1.92	0.52
2:H:98:ARG:NH2	2:H:105:ASP:OD2	2.41	0.51
3:C:327:ASP:OD1	5:C:705:HOH:O	2.19	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:187:ALA:O	1:L:191:LYS:HD3	2.11	0.51
2:H:130:PRO:HB3	2:H:142:LEU:HB3	1.94	0.49
3:C:355:PHE:CE2	3:C:377:LEU:HD11	2.47	0.49
1:L:123:PRO:HD3	1:L:135:VAL:HG22	1.95	0.49
1:L:173:ASP:OD1	1:L:175:THR:HG22	2.13	0.48
1:L:8:PRO:HG2	1:L:11:LEU:HD11	1.95	0.48
1:L:19:VAL:HG12	1:L:80:LEU:HD21	1.95	0.48
3:C:324:PHE:HE2	3:C:434:PRO:HG3	1.79	0.47
3:C:342:LYS:NZ	3:C:344:GLU:OE2	2.34	0.47
3:C:345:HIS:ND1	3:C:395:SER:OG	2.46	0.47
3:C:469:PHE:N	5:C:724:HOH:O	2.47	0.47
1:L:31:ARG:NH1	3:C:363:PRO:O	2.47	0.47
2:H:123:PRO:HD2	2:H:209:THR:HG21	1.95	0.47
3:C:427:GLN:NE2	3:C:475:GLN:OE1	2.48	0.47
3:C:323:VAL:HB	3:C:407:VAL:HG22	1.96	0.47
3:C:355:PHE:CE2	3:C:357:ASP:HB2	2.51	0.46
3:C:442:PHE:HB3	3:C:452:ALA:HB3	1.97	0.46
2:H:58:THR:HG21	5:H:317:HOH:O	2.16	0.45
3:C:333:LEU:HD12	3:C:377:LEU:HD23	1.99	0.45
1:L:37:TRP:CZ3	1:L:90:CYS:HB3	2.52	0.45
2:H:28:ASN:N	2:H:28:ASN:OD1	2.50	0.45
1:L:116:PRO:HB3	1:L:142:PHE:HB3	2.00	0.44
1:L:123:PRO:HG2	1:L:133:ALA:HB1	1.99	0.44
1:L:170:ASP:OD1	1:L:171:SER:N	2.50	0.44
2:H:32:TYR:OH	3:C:330:ASN:O	2.34	0.44
1:L:40:GLN:O	1:L:86:ALA:HB1	2.18	0.44
1:L:175:THR:OG1	1:L:176:TYR:N	2.40	0.44
2:H:33:MET:HB3	2:H:99:TYR:HB3	1.99	0.43
1:L:140:ASN:OD1	1:L:141:ASN:ND2	2.52	0.43
2:H:131:SER:H	2:H:134:SER:HB3	1.83	0.43
3:C:423:ASN:N	5:C:729:HOH:O	2.51	0.42
3:C:471:LYS:HB2	3:C:472:LEU:H	1.70	0.42
2:H:109:GLN:H	2:H:109:GLN:CD	2.28	0.42
1:L:38:TYR:CE1	1:L:91:GLN:HG2	2.50	0.42
3:C:413:PRO:O	3:C:500:THR:OG1	2.31	0.41
1:L:63:ARG:NE	1:L:84:ASP:OD2	2.40	0.41
1:L:2:ILE:HA	1:L:3:GLN:HA	1.65	0.41
2:H:135:THR:HA	2:H:140:ALA:HA	2.03	0.41
1:L:127:GLN:HG2	1:L:132:THR:O	2.21	0.41
1:L:110:LYS:HA	1:L:143:TYR:OH	2.21	0.41
2:H:83:MET:HE2	2:H:86:LEU:HD21	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:156:VAL:HG22	2:H:202:VAL:HG22	2.02	0.40
3:C:359:TRP:HB3	3:C:377:LEU:HD13	2.02	0.40
3:C:346:GLN:HA	3:C:393:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	217/228 (95%)	202 (93%)	14 (6%)	1 (0%)	24	48
2	H	218/221 (99%)	213 (98%)	5 (2%)	0	100	100
3	C	187/214 (87%)	176 (94%)	11 (6%)	0	100	100
All	All	622/663 (94%)	591 (95%)	30 (5%)	1 (0%)	43	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	79	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	189/199 (95%)	177 (94%)	12 (6%)	16	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	H	183/184 (100%)	176 (96%)	7 (4%)	29	58
3	C	167/188 (89%)	163 (98%)	4 (2%)	43	72
All	All	539/571 (94%)	516 (96%)	23 (4%)	26	54

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	19	VAL
1	L	20	THR
1	L	21	ILE
1	L	60	VAL
1	L	80	LEU
1	L	91	GLN
1	L	98	LEU
1	L	109	ILE
1	L	126	SER
1	L	152	LYS
1	L	200	THR
1	L	205	SER
2	H	28	ASN
2	H	65	LYS
2	H	131	SER
2	H	142	LEU
2	H	188	VAL
2	H	195	THR
2	H	211	VAL
3	C	356	THR
3	C	399	VAL
3	C	409	VAL
3	C	443	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	6	GLN
1	L	91	GLN
1	L	92	GLN
1	L	141	ASN
3	C	370	ASN
3	C	400	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	427	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	601	3	14,14,15	0.26	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	601	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	221/228 (96%)	1.50	56 (25%) 1 1	34, 61, 99, 114	0
2	H	220/221 (99%)	0.68	17 (7%) 19 17	25, 48, 74, 95	0
3	C	193/214 (90%)	1.50	52 (26%) 1 1	27, 55, 96, 107	0
All	All	634/663 (95%)	1.22	125 (19%) 3 2	25, 53, 91, 114	0

All (125) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	470	GLY	7.1
1	L	227	ASP	7.0
1	L	13	ALA	6.7
3	C	469	PHE	6.2
1	L	104	GLY	6.2
1	L	2	ILE	6.2
1	L	97	SER	6.0
1	L	96	HIS	5.7
3	C	482	ALA	5.7
3	C	382	LEU	5.4
3	C	486	ASN	4.9
3	C	366	GLU	4.7
1	L	216	GLU	4.7
1	L	15	VAL	4.3
1	L	98	LEU	4.1
1	L	14	SER	4.0
2	H	219	SER	4.0
3	C	485	HIS	4.0
3	C	409	VAL	3.9
1	L	78	SER	3.9
1	L	226	ASP	3.9
3	C	361	ASP	3.8
1	L	222	TYR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	452	ALA	3.7
3	C	449	ARG	3.6
2	H	137	GLY	3.6
1	L	10	SER	3.6
1	L	111	ARG	3.5
1	L	215	GLY	3.5
1	L	107	VAL	3.5
3	C	444	PRO	3.5
1	L	77	ILE	3.4
3	C	344	GLU	3.4
3	C	368	GLU	3.4
3	C	448	GLN	3.3
3	C	487	GLY	3.3
1	L	109	ILE	3.3
1	L	174	SER	3.3
3	C	360	GLU	3.3
1	L	29	GLY	3.3
1	L	82	PRO	3.2
3	C	422	VAL	3.2
3	C	384	GLY	3.1
3	C	481	SER	3.1
3	C	508	PHE	3.0
3	C	509	LYS	2.9
1	L	24	ARG	2.9
1	L	175	THR	2.9
3	C	488	THR	2.9
3	C	483	PHE	2.9
1	L	11	LEU	2.9
1	L	80	LEU	2.9
1	L	79	SER	2.9
2	H	0	SER	2.9
2	H	42	GLY	2.9
3	C	465	SER	2.8
3	C	463	ASN	2.8
3	C	471	LYS	2.8
1	L	108	GLU	2.8
1	L	21	ILE	2.8
3	C	410	ASN	2.8
2	H	218	LYS	2.7
3	C	351	MET	2.7
3	C	385	THR	2.7
1	L	62	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	138	GLY	2.7
3	C	311	GLY	2.7
1	L	76	THR	2.7
3	C	328	GLY	2.6
3	C	380	THR	2.6
1	L	3	GLN	2.6
3	C	497	VAL	2.6
3	C	327	ASP	2.6
3	C	358	LYS	2.6
3	C	407	VAL	2.6
1	L	12	SER	2.5
1	L	83	GLU	2.5
3	C	443	CYS	2.5
1	L	89	TYR	2.5
3	C	324	PHE	2.5
1	L	173	ASP	2.5
2	H	65	LYS	2.5
1	L	172	LYS	2.4
1	L	85	PHE	2.4
1	L	223	LYS	2.4
1	L	28	ARG	2.4
3	C	472	LEU	2.4
2	H	134	SER	2.4
3	C	383	LYS	2.4
1	L	171	SER	2.4
1	L	70	GLY	2.4
3	C	442	PHE	2.4
3	C	408	TYR	2.3
3	C	424	GLY	2.3
1	L	87	THR	2.3
2	H	133	LYS	2.3
3	C	387	GLY	2.3
2	H	200	CYS	2.3
1	L	86	ALA	2.3
1	L	113	VAL	2.3
2	H	197	THR	2.2
1	L	168	GLU	2.2
3	C	414	GLU	2.2
1	L	31	ARG	2.2
1	L	224	ASP	2.2
1	L	16	GLY	2.2
2	H	118	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	145	ARG	2.2
2	H	136	SER	2.2
1	L	64	PHE	2.2
1	L	144	PRO	2.2
2	H	28	ASN	2.2
3	C	450	CYS	2.1
2	H	217	PRO	2.1
1	L	81	GLN	2.1
3	C	451	SER	2.1
3	C	484	LYS	2.1
2	H	132	SER	2.0
1	L	22	THR	2.0
3	C	329	GLU	2.0
3	C	479	ASP	2.0
3	C	342	LYS	2.0
1	L	61	PRO	2.0
2	H	196	GLN	2.0
1	L	213	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	601	14/15	0.89	0.13	29,32,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.