



wwPDB EM Validation Summary Report ⓘ

Mar 7, 2026 – 02:40 AM UTC

PDB ID : 6K9Q / pdb_00006k9q
EMDB ID : EMD-9952
Title : Structure of the native supercoiled hook as a universal joint
Authors : Kato, T.; Miyata, T.; Makino, F.; Horvath, P.; Namba, K.
Deposited on : 2019-06-17
Resolution : 3.10 Å (reported)
Based on initial models : 3A69, 5JXL

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14724 (2.60 - 3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	28% 79% 20% .
1	B	402	41% 82% 17%
1	C	402	23% 84% 15%
1	D	402	21% 82% 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	402	23% 80% 19%
1	F	402	23% 79% 21%
1	G	402	22% 80% 20%
1	H	402	25% 78% 22%
1	I	402	22% 76% 24%
1	J	402	18% 79% 21%
1	K	402	21% 84% 15%
1	L	402	21% 80% 20%
1	M	402	24% 75% 25%
1	N	402	26% 78% 22%
1	O	402	19% 77% 23%
1	P	402	22% 79% 21%
1	Q	402	25% 79% 21%
1	R	402	24% 78% 22%
1	S	402	33% 76% 23%
1	T	402	24% 76% 24%
1	U	402	34% 80% 19%
1	V	402	38% 81% 19%
1	W	402	30% 82% 16%
1	X	402	48% 79% 21%
1	Y	402	37% 80% 20%
1	Z	402	46% 79% 21%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 76902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Flagellar hook protein FlgE.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	B	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	C	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	D	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	E	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	F	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	G	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	H	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	I	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	J	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	K	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	L	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	M	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	N	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	O	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	P	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	Q	401	Total	C	N	O	S	0	0
			2952	1817	510	617	8		



Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	S	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	T	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	U	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	V	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	W	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	X	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	Y	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		
1	Z	402	Total	C	N	O	S	0	0
			2958	1820	511	619	8		




• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE





• Molecule 1: Flagellar hook protein FlgE








- Molecule 1: Flagellar hook protein FlgE





- Molecule 1: Flagellar hook protein FlgE







- Molecule 1: Flagellar hook protein FlgE





- Molecule 1: Flagellar hook protein FlgE








• Molecule 1: Flagellar hook protein FlgE



• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE








• Molecule 1: Flagellar hook protein FlgE





• Molecule 1: Flagellar hook protein FlgE








• Molecule 1: Flagellar hook protein FlgE







• Molecule 1: Flagellar hook protein FlgE










• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE





• Molecule 1: Flagellar hook protein FlgE








• Molecule 1: Flagellar hook protein FlgE




• Molecule 1: Flagellar hook protein FlgE





• Molecule 1: Flagellar hook protein FlgE



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	157334	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	JEOL CRYO ARM 200	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	0.87	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	50000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	1.728	Depositor
Minimum map value	-0.954	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.165	Depositor
Recommended contour level	0.54	Depositor
Map size (Å)	351.04, 351.04, 351.04	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.097, 1.097, 1.097	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/3002	0.63	2/4090 (0.0%)
1	B	0.29	0/3002	0.66	2/4090 (0.0%)
1	C	0.31	0/3002	0.70	2/4090 (0.0%)
1	D	0.43	3/3002 (0.1%)	0.68	4/4090 (0.1%)
1	E	0.37	1/3002 (0.0%)	0.66	3/4090 (0.1%)
1	F	0.30	0/3002	0.67	0/4090
1	G	0.38	0/3002	0.67	1/4090 (0.0%)
1	H	0.31	0/3002	0.64	0/4090
1	I	0.32	0/3002	0.67	0/4090
1	J	0.32	0/3002	0.63	0/4090
1	K	0.30	0/3002	0.61	0/4090
1	L	0.31	0/3002	0.67	0/4090
1	M	0.39	0/3002	0.66	0/4090
1	N	0.32	0/3002	0.65	0/4090
1	O	0.33	0/3002	0.69	0/4090
1	P	0.31	0/3002	0.64	0/4090
1	Q	0.34	0/2996	0.71	0/4082
1	R	0.30	0/3002	0.64	2/4090 (0.0%)
1	S	0.38	1/3002 (0.0%)	0.70	4/4090 (0.1%)
1	T	0.32	0/3002	0.68	0/4090
1	U	0.41	1/3002 (0.0%)	0.69	1/4090 (0.0%)
1	V	0.28	0/3002	0.62	0/4090
1	W	0.58	7/3002 (0.2%)	0.72	3/4090 (0.1%)
1	X	0.26	0/3002	0.60	0/4090
1	Y	0.30	0/3002	0.63	2/4090 (0.0%)
1	Z	0.28	0/3002	0.64	0/4090
All	All	0.34	13/78046 (0.0%)	0.66	26/106332 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	Y	0	1
1	Z	0	1
All	All	0	3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	W	81	PHE	C-O	-6.73	1.15	1.23
1	W	19	ILE	C-O	-6.42	1.15	1.24
1	D	3	SER	C-O	-5.80	1.17	1.24
1	S	387	THR	CA-C	-5.64	1.44	1.52
1	U	93	TYR	CA-C	-5.40	1.46	1.52

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	74	VAL	CB-CA-C	-11.92	98.47	111.59
1	W	80	GLY	N-CA-C	7.61	122.45	112.25
1	W	20	GLY	N-CA-C	-6.63	104.47	112.49
1	U	332	VAL	CB-CA-C	-6.57	100.52	111.29
1	A	77	SER	CA-C-N	5.74	132.49	121.54

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	284	LYS	Peptide
1	Y	268	GLN	Peptide
1	Z	77	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2958	0	2855	60	0
1	B	2958	0	2855	60	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2958	0	2855	42	0
1	D	2958	0	2855	48	0
1	E	2958	0	2855	58	0
1	F	2958	0	2855	68	0
1	G	2958	0	2855	74	0
1	H	2958	0	2855	77	0
1	I	2958	0	2855	70	0
1	J	2958	0	2855	68	0
1	K	2958	0	2855	46	0
1	L	2958	0	2855	59	0
1	M	2958	0	2855	72	0
1	N	2958	0	2855	78	0
1	O	2958	0	2855	77	0
1	P	2958	0	2855	63	0
1	Q	2952	0	2847	73	0
1	R	2958	0	2855	70	0
1	S	2958	0	2855	76	0
1	T	2958	0	2855	71	0
1	U	2958	0	2855	60	0
1	V	2958	0	2855	56	0
1	W	2958	0	2855	63	0
1	X	2958	0	2855	58	0
1	Y	2958	0	2855	50	0
1	Z	2958	0	2855	58	0
All	All	76902	0	74222	1446	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 1446 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:388:ILE:HG13	1:S:389:LYS:H	1.31	0.91
1:A:49:LEU:HD12	1:A:49:LEU:O	1.71	0.90
1:G:74:VAL:HG22	1:G:75:ALA:N	1.87	0.90
1:W:21:ASN:OD1	1:W:25:ASN:OD1	1.95	0.85
1:U:82:PHE:HB2	1:U:94:SER:O	1.77	0.84

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/402 (100%)	372 (93%)	25 (6%)	3 (1%)	16	47
1	B	400/402 (100%)	372 (93%)	27 (7%)	1 (0%)	36	67
1	C	400/402 (100%)	374 (94%)	25 (6%)	1 (0%)	36	67
1	D	400/402 (100%)	377 (94%)	23 (6%)	0	100	100
1	E	400/402 (100%)	375 (94%)	24 (6%)	1 (0%)	36	67
1	F	400/402 (100%)	370 (92%)	30 (8%)	0	100	100
1	G	400/402 (100%)	380 (95%)	20 (5%)	0	100	100
1	H	400/402 (100%)	383 (96%)	17 (4%)	0	100	100
1	I	400/402 (100%)	374 (94%)	26 (6%)	0	100	100
1	J	400/402 (100%)	376 (94%)	24 (6%)	0	100	100
1	K	400/402 (100%)	370 (92%)	30 (8%)	0	100	100
1	L	400/402 (100%)	375 (94%)	25 (6%)	0	100	100
1	M	400/402 (100%)	372 (93%)	28 (7%)	0	100	100
1	N	400/402 (100%)	380 (95%)	20 (5%)	0	100	100
1	O	400/402 (100%)	375 (94%)	25 (6%)	0	100	100
1	P	400/402 (100%)	375 (94%)	25 (6%)	0	100	100
1	Q	399/402 (99%)	379 (95%)	20 (5%)	0	100	100
1	R	400/402 (100%)	379 (95%)	21 (5%)	0	100	100
1	S	400/402 (100%)	372 (93%)	27 (7%)	1 (0%)	36	67
1	T	400/402 (100%)	377 (94%)	23 (6%)	0	100	100
1	U	400/402 (100%)	374 (94%)	26 (6%)	0	100	100
1	V	400/402 (100%)	369 (92%)	31 (8%)	0	100	100
1	W	400/402 (100%)	367 (92%)	32 (8%)	1 (0%)	36	67
1	X	400/402 (100%)	374 (94%)	26 (6%)	0	100	100
1	Y	400/402 (100%)	374 (94%)	26 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Z	400/402 (100%)	369 (92%)	30 (8%)	1 (0%)	36	67
All	All	10399/10452 (100%)	9734 (94%)	656 (6%)	9 (0%)	49	78

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	LYS
1	S	389	LYS
1	A	78	GLN
1	A	285	PRO
1	B	78	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/322 (100%)	322 (100%)	0	100	100
1	B	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	C	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	D	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	E	322/322 (100%)	320 (99%)	2 (1%)	78	83
1	F	322/322 (100%)	322 (100%)	0	100	100
1	G	322/322 (100%)	320 (99%)	2 (1%)	78	83
1	H	322/322 (100%)	320 (99%)	2 (1%)	78	83
1	I	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	J	322/322 (100%)	322 (100%)	0	100	100
1	K	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	L	322/322 (100%)	320 (99%)	2 (1%)	78	83
1	M	322/322 (100%)	322 (100%)	0	100	100
1	N	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	O	322/322 (100%)	320 (99%)	2 (1%)	78	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	322/322 (100%)	322 (100%)	0	100	100
1	Q	321/322 (100%)	320 (100%)	1 (0%)	86	87
1	R	322/322 (100%)	320 (99%)	2 (1%)	78	83
1	S	322/322 (100%)	322 (100%)	0	100	100
1	T	322/322 (100%)	322 (100%)	0	100	100
1	U	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	V	322/322 (100%)	322 (100%)	0	100	100
1	W	322/322 (100%)	320 (99%)	2 (1%)	78	83
1	X	322/322 (100%)	321 (100%)	1 (0%)	86	87
1	Y	322/322 (100%)	322 (100%)	0	100	100
1	Z	322/322 (100%)	321 (100%)	1 (0%)	86	87
All	All	8371/8372 (100%)	8347 (100%)	24 (0%)	84	87

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	46	LYS
1	R	53	VAL
1	Q	53	VAL
1	R	74	VAL
1	G	365	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 140 such sidechains are listed below:

Mol	Chain	Res	Type
1	W	25	ASN
1	W	382	GLN
1	Y	373	ASN
1	I	328	GLN
1	I	318	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-9952. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 174



Y Index: 155




Z Index: 170


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.54. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 493 nm³; this corresponds to an approximate mass of 445 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	3.61	4.20	3.69
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from author-provided FSC intersecting FSC 0.143 CUT-OFF 3.61 differs from the reported value 3.1 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-9952 and PDB model 6K9Q. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.54 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.54).

9.4 Atom inclusion [i](#)



At the recommended contour level, 69% of all backbone atoms, 51% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.54) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5060	 0.4440
A	 0.4990	 0.4380
B	 0.4440	 0.4300
C	 0.5250	 0.4580
D	 0.5240	 0.4580
E	 0.5310	 0.4540
F	 0.5260	 0.4390
G	 0.5300	 0.4450
H	 0.5150	 0.4470
I	 0.5270	 0.4550
J	 0.5640	 0.4690
K	 0.5370	 0.4560
L	 0.5350	 0.4500
M	 0.5360	 0.4560
N	 0.5180	 0.4530
O	 0.5450	 0.4600
P	 0.5330	 0.4500
Q	 0.5180	 0.4370
R	 0.5220	 0.4470
S	 0.4800	 0.4340
T	 0.5210	 0.4460
U	 0.4810	 0.4360
V	 0.4690	 0.4260
W	 0.4900	 0.4360
X	 0.4090	 0.4090
Y	 0.4610	 0.4380
Z	 0.4220	 0.4270

