



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:50 AM UTC

PDB ID : 2KAU / pdb_00002kau
Title : THE CRYSTAL STRUCTURE OF UREASE FROM KLEBSIELLA AEROGENES AT 2.2 ANGSTROMS RESOLUTION
Authors : Jabri, E.; Carr, M.B.; Hausinger, R.P.; Karplus, P.A.
Deposited on : 1995-02-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	0% 96% •
2	B	106	0% 83% 11% • 5%
3	C	567	3% 84% 14% •

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE (GAMMA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			775	491	134	145	5			

- Molecule 2 is a protein called UREASE (BETA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	101	Total	C	N	O	S	0	0	0
			784	496	150	135	3			

- Molecule 3 is a protein called UREASE (ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	566	Total	C	N	O	S	0	0	0
			4228	2652	741	812	23			

- Molecule 4 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Ni	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	22	Total	O	0	0
			22	22		
5	B	17	Total	O	0	0
			17	17		
5	C	176	Total	O	0	0
			176	176		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




- Molecule 1: UREASE (GAMMA CHAIN)



- Molecule 2: UREASE (BETA CHAIN)



- Molecule 3: UREASE (ALPHA CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	170.80Å 170.80Å 170.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 10.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.6 (10.00-2.00) 98.8 (10.00-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.00Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.185 , 0.225 0.181 , 0.217	Depositor DCC
R_{free} test set	2824 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	10.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.041 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6004	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/786	0.83	0/1061
2	B	0.43	0/804	0.91	1/1087 (0.1%)
3	C	0.51	0/4300	1.03	28/5860 (0.5%)
All	All	0.50	0/5890	0.99	29/8008 (0.4%)

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	436	SER	N-CA-C	-8.32	99.58	110.07
3	C	173	GLY	CA-C-N	8.19	127.76	119.24
3	C	173	GLY	C-N-CA	8.19	127.76	119.24
3	C	283	ASP	N-CA-C	7.67	122.39	112.26
3	C	299	ASN	N-CA-C	7.61	126.63	109.81
3	C	298	THR	N-CA-C	-7.35	99.03	110.42
3	C	50	VAL	N-CA-C	7.26	118.92	111.00
2	B	38	SER	N-CA-C	6.88	118.78	111.28
3	C	247	SER	N-CA-C	6.39	118.11	110.19
3	C	201	GLN	CA-C-N	6.19	126.38	119.32
3	C	201	GLN	C-N-CA	6.19	126.38	119.32
3	C	290	HIS	N-CA-C	6.05	117.69	110.07
3	C	562	GLN	N-CA-C	5.89	119.93	112.87
3	C	409	ILE	N-CA-C	5.82	116.60	110.72
3	C	565	PHE	N-CA-C	5.82	118.44	109.07
3	C	199	VAL	CB-CA-C	-5.79	103.75	111.80
3	C	361	SER	N-CA-C	5.79	119.20	110.64
3	C	320	HIS	N-CA-C	5.65	119.87	113.15
3	C	355	SER	N-CA-C	5.42	120.05	113.38
3	C	96	ILE	N-CA-C	-5.37	98.75	107.28
3	C	245	LEU	N-CA-C	5.21	117.89	109.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	369	GLU	N-CA-C	5.21	119.18	112.41
3	C	354	PHE	N-CA-C	-5.19	100.78	109.24
3	C	101	ASN	CA-C-N	5.10	124.76	119.56
3	C	101	ASN	C-N-CA	5.10	124.76	119.56
3	C	173	GLY	N-CA-C	5.09	122.72	112.34
3	C	308	THR	N-CA-C	5.04	117.17	111.02
3	C	312	HIS	N-CA-C	5.03	117.66	111.82
3	C	275	GLY	N-CA-C	5.02	120.70	114.37

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	775	0	807	3	0
2	B	784	0	775	9	0
3	C	4228	0	4176	40	0
4	C	2	0	0	0	0
5	A	22	0	0	1	0
5	B	17	0	0	1	0
5	C	176	0	0	1	0
All	All	6004	0	5758	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:83:ALA:HB1	2:B:88:ARG:HH21	1.56	0.70
2:B:72:GLU:H	2:B:75:GLN:HE21	1.41	0.69
3:C:324:PRO:HA	3:C:330:VAL:CG1	2.25	0.67
3:C:218:ILE:HD13	3:C:243:VAL:HG13	1.76	0.65
3:C:201:GLN:HE21	3:C:203:ASP:HB2	1.66	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:236:ALA:O	3:C:518:ARG:HD2	2.01	0.60
3:C:316:LEU:O	3:C:317:MET:HG2	2.02	0.58
2:B:72:GLU:H	2:B:75:GLN:NE2	2.00	0.57
3:C:297:SER:OG	3:C:349:HIS:HE1	1.88	0.57
3:C:545:VAL:O	3:C:548:GLU:HG2	2.04	0.56
3:C:324:PRO:HA	3:C:330:VAL:HG11	1.88	0.54
3:C:300:PRO:HD3	3:C:365:GLY:HA2	1.91	0.52
3:C:395:ASN:HD22	3:C:397:ASN:H	1.57	0.52
3:C:199:VAL:HG23	3:C:205:LEU:HD21	1.92	0.51
2:B:90:VAL:HB	2:B:97:VAL:HG11	1.93	0.50
3:C:60:MET:HE2	3:C:111:PRO:HB2	1.93	0.50
1:A:53:VAL:O	1:A:57:MET:HG3	2.13	0.49
3:C:296:SER:HB3	3:C:356:LEU:HB2	1.94	0.48
3:C:395:ASN:HD22	3:C:395:ASN:C	2.21	0.48
2:B:50:LYS:HB2	5:B:116:HOH:O	2.12	0.48
3:C:237:ASP:OD1	3:C:518:ARG:HD3	2.14	0.47
3:C:281:ALA:HA	3:C:282:PRO:HA	1.75	0.46
3:C:20:LYS:HE3	3:C:394:ASP:O	2.16	0.46
3:C:218:ILE:HD13	3:C:243:VAL:CG1	2.46	0.46
3:C:201:GLN:HE21	3:C:203:ASP:H	1.64	0.46
2:B:4:GLY:HA2	3:C:22:ARG:O	2.15	0.45
3:C:75:ILE:O	3:C:81:ILE:HA	2.17	0.45
3:C:271:PHE:O	3:C:272:HIS:C	2.59	0.45
3:C:218:ILE:HD11	3:C:232:ALA:CB	2.46	0.45
3:C:298:THR:CG2	3:C:360:ASP:HB2	2.47	0.45
3:C:65:CYS:HA	3:C:111:PRO:HG2	1.99	0.45
3:C:419:HIS:CD2	3:C:420:GLU:HG2	2.52	0.45
2:B:100:PRO:O	2:B:101:LEU:CB	2.66	0.44
3:C:285:ILE:HD11	3:C:348:LEU:HD12	1.99	0.44
3:C:326:ILE:O	3:C:330:VAL:HG22	2.17	0.43
3:C:236:ALA:C	3:C:518:ARG:HD2	2.43	0.43
3:C:505:LEU:HB2	3:C:507:LEU:HD13	1.99	0.43
3:C:246:HIS:CE1	3:C:278:GLY:HA3	2.53	0.43
3:C:381:MET:HE2	3:C:381:MET:HA	2.01	0.43
3:C:324:PRO:HA	3:C:330:VAL:HG13	1.99	0.42
1:A:10:LYS:HD2	1:A:44:MET:SD	2.59	0.42
3:C:440:PHE:O	3:C:560:MET:HE2	2.19	0.42
3:C:217:KCX:CX	3:C:219:HIS:HD2	2.33	0.42
2:B:89:ALA:HA	2:B:100:PRO:HA	2.02	0.41
2:B:90:VAL:HB	2:B:97:VAL:CG1	2.50	0.41
3:C:460:ASP:HB3	3:C:463:ALA:HB2	2.00	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:447:VAL:HB	3:C:455:ILE:HG22	2.02	0.41
1:A:10:LYS:NZ	5:A:111:HOH:O	2.54	0.41
3:C:418:ALA:HB3	5:C:817:HOH:O	2.20	0.41
3:C:138:ILE:HD12	3:C:362:GLN:HB2	2.02	0.40
3:C:92:ARG:HD3	3:C:427:GLY:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	B	99/106 (93%)	94 (95%)	5 (5%)	0	100	100
3	C	563/567 (99%)	531 (94%)	28 (5%)	4 (1%)	18	14
All	All	760/773 (98%)	722 (95%)	34 (4%)	4 (0%)	24	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	360	ASP
3	C	317	MET
3	C	364	MET
3	C	481	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	85 (100%)	0	100	100
2	B	78/83 (94%)	76 (97%)	2 (3%)	40	44
3	C	442/443 (100%)	430 (97%)	12 (3%)	39	42
All	All	605/611 (99%)	591 (98%)	14 (2%)	44	49

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	81	LEU
2	B	97	VAL
3	C	179	ARG
3	C	199	VAL
3	C	282	PRO
3	C	314	ASP
3	C	332	PHE
3	C	349	HIS
3	C	394	ASP
3	C	395	ASN
3	C	449	LYS
3	C	465	ILE
3	C	507	LEU
3	C	508	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	75	GLN
3	C	7	GLN
3	C	142	GLN
3	C	201	GLN
3	C	349	HIS
3	C	362	GLN
3	C	395	ASN
3	C	469	GLN
3	C	486	HIS
3	C	506	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	KCX	C	217	3,4	10,11,12	0.88	1 (10%)	6,12,14	1.68	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	217	3,4	-	0/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	217	KCX	OQ1-CX	2.01	1.25	1.21

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	217	KCX	OQ1-CX-NZ	-3.66	119.37	124.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	217	KCX	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/100 (100%)	-0.72	1 (1%) 79 79	3, 8, 17, 20	0
2	B	101/106 (95%)	-0.22	1 (0%) 79 79	7, 15, 21, 25	0
3	C	565/567 (99%)	-0.59	19 (3%) 48 47	2, 7, 21, 57	0
All	All	766/773 (99%)	-0.56	21 (2%) 56 55	2, 7, 21, 57	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	331	ALA	5.8
3	C	330	VAL	4.8
3	C	316	LEU	4.7
3	C	326	ILE	4.5
3	C	327	ALA	3.8
3	C	332	PHE	3.5
3	C	321	HIS	3.5
3	C	313	LEU	3.4
3	C	322	LEU	3.4
3	C	324	PRO	3.2
3	C	333	ALA	3.0
3	C	323	ASP	2.8
3	C	320	HIS	2.8
3	C	325	ASP	2.8
1	A	100	ILE	2.5
3	C	318	VAL	2.5
3	C	314	ASP	2.4
3	C	548	GLU	2.3
3	C	317	MET	2.3
3	C	319	CYS	2.2
2	B	54	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KCX	C	217	12/13	0.96	0.05	5,5,6,6	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NI	C	774	1/1	0.99	0.05	13,13,13,13	0
4	NI	C	775	1/1	0.99	0.07	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.