



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 11:24 PM UTC

PDB ID : 5KAH / pdb_00005kah
Title : Crystal structure of a dioxygenase in the Crotonase superfamily in P21, V425T mutant
Authors : Li, K.; Fielding, E.N.; Condurso, H.L.; Bruner, S.D.
Deposited on : 2016-06-01
Resolution : 2.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	 86% 8% . .
1	B	438	 83% 11% . 5%
1	C	438	 81% 13% . 5%
1	D	438	 84% 11% .
1	E	438	 83% 11% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	438	 80% 15% . .
1	G	438	 83% 11% . .
1	H	438	 83% 12% . 5%
1	I	438	 82% 13% . .
1	J	438	 84% 11% . .
1	K	438	 84% 12% .
1	L	438	 83% 11% . 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 40568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	421	Total	C	N	O	S	0	0	0
			3272	2055	611	596	10			
1	B	418	Total	C	N	O	S	0	0	0
			3251	2042	607	592	10			
1	C	417	Total	C	N	O	S	0	0	0
			3244	2038	606	590	10			
1	D	421	Total	C	N	O	S	0	0	0
			3273	2055	611	597	10			
1	E	421	Total	C	N	O	S	0	0	0
			3272	2055	611	596	10			
1	F	421	Total	C	N	O	S	0	0	0
			3272	2055	611	596	10			
1	G	419	Total	C	N	O	S	0	0	0
			3256	2045	608	593	10			
1	H	418	Total	C	N	O	S	0	0	0
			3251	2042	607	592	10			
1	I	421	Total	C	N	O	S	0	0	0
			3272	2055	611	596	10			
1	J	419	Total	C	N	O	S	0	0	0
			3253	2043	608	592	10			
1	K	419	Total	C	N	O	S	0	0	0
			3260	2047	608	595	10			
1	L	418	Total	C	N	O	S	0	0	0
			3246	2040	607	589	10			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	425	THR	VAL	engineered mutation	UNP Q8KLK7
B	425	THR	VAL	engineered mutation	UNP Q8KLK7
C	425	THR	VAL	engineered mutation	UNP Q8KLK7
D	425	THR	VAL	engineered mutation	UNP Q8KLK7
E	425	THR	VAL	engineered mutation	UNP Q8KLK7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	425	THR	VAL	engineered mutation	UNP Q8KLLK7
G	425	THR	VAL	engineered mutation	UNP Q8KLLK7
H	425	THR	VAL	engineered mutation	UNP Q8KLLK7
I	425	THR	VAL	engineered mutation	UNP Q8KLLK7
J	425	THR	VAL	engineered mutation	UNP Q8KLLK7
K	425	THR	VAL	engineered mutation	UNP Q8KLLK7
L	425	THR	VAL	engineered mutation	UNP Q8KLLK7

- Molecule 2 is [(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-4-HYDROXY-3-(PHOSPHONOXY)TETRAHYDROFURAN-2-YL]METHYL (3R)-4-({3-[(2-{[(3,5-DIHYDROXYPHENYL)ACETYL]AMINO}ETHYL)AMINO]-3-OXOPROPYL}AMINO)-3-HYDROXY-2,2-DIMETHYL-4-OXOBUTYL DIHYDROGEN DIPHOSPHATE (CCD ID: YE1) (formula: C₂₉H₄₃N₈O₁₉P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	B	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	C	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	D	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	E	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	F	1	Total	C	N	O	P	0	0
			59	29	8	19	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	G	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	H	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	I	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	J	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	K	1	Total	C	N	O	P	0	0
			59	29	8	19	3		
2	L	1	Total	C	N	O	P	0	0
			59	29	8	19	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		
3	B	56	Total	O	0	0
			56	56		
3	C	62	Total	O	0	0
			62	62		
3	D	61	Total	O	0	0
			61	61		
3	E	74	Total	O	0	0
			74	74		
3	F	60	Total	O	0	0
			60	60		
3	G	65	Total	O	0	0
			65	65		
3	H	66	Total	O	0	0
			66	66		
3	I	53	Total	O	0	0
			53	53		
3	J	53	Total	O	0	0
			53	53		
3	K	67	Total	O	0	0
			67	67		
3	L	63	Total	O	0	0
			63	63		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase






- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain E: 83% 11% . .




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain F: 80% 15% . .




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain G: 83% 11% . .



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain H: 83% 12% 5% . .




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain I: 82% 13% . .



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain J: 84% 11% . .




- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain K: 84% 12% .



- Molecule 1: (3,5-dihydroxyphenyl)acetyl-CoA 1,2-dioxygenase

Chain L: 83% 11% . 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	139.77Å 171.96Å 156.11Å 90.00° 90.02° 90.00°	Depositor
Resolution (Å)	39.03 – 2.78 39.03 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.03-2.78) 99.7 (39.03-2.78)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.77Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.198 , 0.231 0.193 , 0.226	Depositor DCC
R_{free} test set	9405 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.885	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 11.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.447 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	40568	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.46 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.6739e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YE1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/3333	0.52	2/4515 (0.0%)
1	B	0.20	0/3311	0.49	0/4485
1	C	0.18	0/3304	0.48	0/4475
1	D	0.20	0/3333	0.49	1/4513 (0.0%)
1	E	0.28	1/3333 (0.0%)	0.51	0/4515
1	F	0.32	1/3333 (0.0%)	0.52	0/4515
1	G	0.20	0/3316	0.50	0/4492
1	H	0.20	0/3311	0.47	0/4485
1	I	0.20	0/3333	0.50	0/4515
1	J	0.20	0/3313	0.49	0/4487
1	K	0.18	0/3320	0.47	0/4497
1	L	0.20	0/3306	0.49	0/4478
All	All	0.22	2/39846 (0.0%)	0.49	3/53972 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	317	LEU	C-N	14.53	1.45	1.33
1	E	324	ILE	C-N	11.18	1.46	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	189	GLU	N-CA-C	6.54	120.53	110.20
1	A	272	ARG	CA-C-N	-5.62	113.99	119.78
1	A	272	ARG	C-N-CA	-5.62	113.99	119.78

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3272	0	3293	27	0
1	B	3251	0	3270	34	0
1	C	3244	0	3263	37	0
1	D	3273	0	3292	25	0
1	E	3272	0	3293	37	0
1	F	3272	0	3293	45	0
1	G	3256	0	3275	38	0
1	H	3251	0	3270	38	0
1	I	3272	0	3293	57	0
1	J	3253	0	3271	32	0
1	K	3260	0	3276	31	0
1	L	3246	0	3269	41	0
2	A	59	0	37	3	0
2	B	59	0	39	3	0
2	C	59	0	39	6	0
2	D	59	0	38	2	0
2	E	59	0	39	4	0
2	F	59	0	38	4	0
2	G	59	0	37	7	0
2	H	59	0	39	3	0
2	I	59	0	38	7	0
2	J	59	0	38	2	0
2	K	59	0	39	4	0
2	L	59	0	38	2	0
3	A	58	0	0	0	0
3	B	56	0	0	2	0
3	C	62	0	0	1	0
3	D	61	0	0	3	0
3	E	74	0	0	5	0
3	F	60	0	0	4	0
3	G	65	0	0	3	0
3	H	66	0	0	4	0
3	I	53	0	0	1	0
3	J	53	0	0	1	0
3	K	67	0	0	4	0
3	L	63	0	0	2	0
All	All	40568	0	39817	431	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 431 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:319:ALA:HB1	1:E:324:ILE:O	1.54	1.08
1:I:322:GLU:CG	1:I:428:LYS:HG3	1.87	1.04
1:E:322:GLU:HG3	1:E:428:LYS:HE2	1.07	1.02
1:E:355:LYS:HB2	3:E:633:HOH:O	1.65	0.96
1:E:322:GLU:HG3	1:E:428:LYS:CE	1.99	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/438 (96%)	401 (96%)	12 (3%)	6 (1%)	9	26
1	B	414/438 (94%)	404 (98%)	7 (2%)	3 (1%)	18	44
1	C	413/438 (94%)	405 (98%)	8 (2%)	0	100	100
1	D	417/438 (95%)	411 (99%)	6 (1%)	0	100	100
1	E	419/438 (96%)	399 (95%)	16 (4%)	4 (1%)	12	35
1	F	419/438 (96%)	407 (97%)	8 (2%)	4 (1%)	12	35
1	G	415/438 (95%)	405 (98%)	9 (2%)	1 (0%)	43	70
1	H	414/438 (94%)	405 (98%)	9 (2%)	0	100	100
1	I	419/438 (96%)	407 (97%)	11 (3%)	1 (0%)	43	70
1	J	415/438 (95%)	405 (98%)	9 (2%)	1 (0%)	43	70
1	K	415/438 (95%)	405 (98%)	10 (2%)	0	100	100
1	L	414/438 (94%)	402 (97%)	11 (3%)	1 (0%)	43	70
All	All	4993/5256 (95%)	4856 (97%)	116 (2%)	21 (0%)	30	57

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	HIS
1	A	322	GLU
1	B	270	ASP
1	F	40	SER
1	F	322	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/345 (96%)	317 (96%)	14 (4%)	26	58
1	B	330/345 (96%)	323 (98%)	7 (2%)	47	76
1	C	329/345 (95%)	321 (98%)	8 (2%)	43	74
1	D	332/345 (96%)	321 (97%)	11 (3%)	33	66
1	E	331/345 (96%)	320 (97%)	11 (3%)	33	66
1	F	331/345 (96%)	322 (97%)	9 (3%)	39	71
1	G	330/345 (96%)	321 (97%)	9 (3%)	39	71
1	H	330/345 (96%)	322 (98%)	8 (2%)	43	74
1	I	331/345 (96%)	321 (97%)	10 (3%)	36	69
1	J	329/345 (95%)	320 (97%)	9 (3%)	39	71
1	K	331/345 (96%)	321 (97%)	10 (3%)	36	69
1	L	328/345 (95%)	319 (97%)	9 (3%)	39	71
All	All	3963/4140 (96%)	3848 (97%)	115 (3%)	37	70

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	340	ARG
1	L	303	VAL
1	H	163	GLU
1	L	163	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	K	148	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	222	HIS
1	L	111	GLN
1	H	64	HIS
1	L	222	HIS
1	I	111	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	YE1	C	501	-	60,62,62	2.13	12 (20%)	87,92,92	1.35	9 (10%)
2	YE1	G	501	-	60,62,62	2.23	12 (20%)	87,92,92	1.25	10 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	YE1	J	501	-	60,62,62	2.04	12 (20%)	87,92,92	1.29	10 (11%)
2	YE1	A	501	-	60,62,62	2.24	12 (20%)	87,92,92	1.25	10 (11%)
2	YE1	I	501	-	60,62,62	2.06	14 (23%)	87,92,92	1.25	11 (12%)
2	YE1	E	501	-	60,62,62	1.97	11 (18%)	87,92,92	1.28	10 (11%)
2	YE1	K	501	-	60,62,62	2.06	14 (23%)	87,92,92	1.32	12 (13%)
2	YE1	B	501	-	60,62,62	1.98	10 (16%)	87,92,92	1.24	7 (8%)
2	YE1	F	501	-	60,62,62	2.05	12 (20%)	87,92,92	1.33	12 (13%)
2	YE1	H	501	-	60,62,62	1.98	13 (21%)	87,92,92	1.33	10 (11%)
2	YE1	L	501	-	60,62,62	2.00	11 (18%)	87,92,92	1.27	11 (12%)
2	YE1	D	501	-	60,62,62	2.00	10 (16%)	87,92,92	1.19	8 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	YE1	C	501	-	-	12/55/71/71	0/4/4/4
2	YE1	G	501	-	-	14/55/71/71	0/4/4/4
2	YE1	J	501	-	-	18/55/71/71	0/4/4/4
2	YE1	A	501	-	-	14/55/71/71	0/4/4/4
2	YE1	I	501	-	-	9/55/71/71	0/4/4/4
2	YE1	E	501	-	-	10/55/71/71	0/4/4/4
2	YE1	K	501	-	-	9/55/71/71	0/4/4/4
2	YE1	B	501	-	-	12/55/71/71	0/4/4/4
2	YE1	F	501	-	-	7/55/71/71	0/4/4/4
2	YE1	H	501	-	-	19/55/71/71	0/4/4/4
2	YE1	L	501	-	-	22/55/71/71	0/4/4/4
2	YE1	D	501	-	-	10/55/71/71	0/4/4/4

The worst 5 of 143 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	YE1	P3'-O3'	13.24	1.82	1.59
2	G	501	YE1	P3'-O3'	13.20	1.82	1.59
2	K	501	YE1	P3'-O3'	11.56	1.79	1.59
2	C	501	YE1	P3'-O3'	11.42	1.79	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	501	YE1	P3'-O3'	11.39	1.79	1.59

The worst 5 of 120 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	YE1	CAC-CAB-NAA	-4.16	110.28	116.06
2	B	501	YE1	O3'-P3'-O7A	-3.89	95.46	109.33
2	F	501	YE1	O3'-P3'-O7A	-3.87	95.52	109.33
2	H	501	YE1	O3'-P3'-O7A	-3.86	95.56	109.33
2	E	501	YE1	O3'-P3'-O7A	-3.84	95.64	109.33

There are no chirality outliers.

5 of 156 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	YE1	C2'-C3'-O3'-P3'
2	A	501	YE1	C5'-O5'-P1A-O1A
2	A	501	YE1	C5'-O5'-P1A-O3A
2	A	501	YE1	C12-O6A-P2A-O5A
2	A	501	YE1	C14-C11-C12-O6A


There are no ring outliers.



12 monomers are involved in 47 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	YE1	6	0
2	G	501	YE1	7	0
2	J	501	YE1	2	0
2	A	501	YE1	3	0
2	I	501	YE1	7	0
2	E	501	YE1	4	0
2	K	501	YE1	4	0
2	B	501	YE1	3	0
2	F	501	YE1	4	0
2	H	501	YE1	3	0
2	L	501	YE1	2	0
2	D	501	YE1	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand YE1 J 501**Ligand YE1 A 501**


Ligand YE1 I 501**Ligand YE1 E 501**


Ligand YE1 K 501



Ligand YE1 B 501







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	421/438 (96%)	-1.26	0 100 100	24, 40, 88, 104	0
1	B	418/438 (95%)	-1.36	0 100 100	23, 38, 59, 72	0
1	C	417/438 (95%)	-1.28	0 100 100	23, 38, 66, 78	0
1	D	421/438 (96%)	-1.40	0 100 100	20, 30, 48, 73	0
1	E	421/438 (96%)	-1.39	0 100 100	21, 36, 55, 72	0
1	F	421/438 (96%)	-1.35	0 100 100	21, 34, 54, 77	0
1	G	419/438 (95%)	-1.39	0 100 100	20, 32, 54, 72	0
1	H	418/438 (95%)	-1.32	0 100 100	23, 39, 57, 76	0
1	I	421/438 (96%)	-1.34	0 100 100	21, 36, 54, 67	0
1	J	419/438 (95%)	-1.25	0 100 100	26, 42, 75, 91	0
1	K	419/438 (95%)	-1.38	0 100 100	20, 34, 52, 64	0
1	L	418/438 (95%)	-1.26	0 100 100	22, 37, 77, 96	0
All	All	5033/5256 (95%)	-1.33	0 100 100	20, 36, 63, 104	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	YE1	A	501	59/59	0.99	0.04	32,46,70,79	0
2	YE1	B	501	59/59	0.99	0.04	30,48,66,72	0
2	YE1	C	501	59/59	0.99	0.04	31,43,58,66	0
2	YE1	D	501	59/59	0.99	0.04	24,35,44,48	0
2	YE1	E	501	59/59	0.99	0.04	30,41,51,55	0
2	YE1	F	501	59/59	0.99	0.03	27,37,50,57	0
2	YE1	G	501	59/59	0.99	0.03	25,39,63,71	0
2	YE1	H	501	59/59	0.99	0.04	32,38,45,46	0
2	YE1	I	501	59/59	0.99	0.04	28,36,42,48	0
2	YE1	J	501	59/59	0.99	0.04	37,48,62,64	0
2	YE1	K	501	59/59	0.99	0.04	26,35,46,54	0
2	YE1	L	501	59/59	0.99	0.04	35,40,50,55	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around YE1 A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YE1 B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YE1 C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YE1 D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YE1 E 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YE1 F 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YE1 G 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YE1 H 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YE1 I 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YE1 J 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YE1 K 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around YE1 L 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.