



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:10 PM UTC

PDB ID : 8KAP / pdb_00008kap
Title : Glycoside hydrolase family 1 beta-glucosidase from *Streptomyces griseus* (ligand-free)
Authors : Kumakura, H.; Motouchi, S.; Nakai, H.; Nakajima, M.
Deposited on : 2023-08-03
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	4% 84% 11% • 5%
1	B	414	% 88% 7% 5%
1	C	414	5% 85% 9% 5%
1	D	414	3% 89% 6% 5%

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 12655 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SGR_2426.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	0	0
			3072	1942	538	584	8			
1	B	393	Total	C	N	O	S	0	0	0
			3056	1932	535	581	8			
1	C	393	Total	C	N	O	S	0	0	0
			3056	1932	535	581	8			
1	D	393	Total	C	N	O	S	0	0	0
			3056	1932	535	581	8			

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula: C₈H₁₉NO₅).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			10	6	4		
4	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			16	10	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	78	Total	O	0	0
			78	78		
6	B	80	Total	O	0	0
			80	80		
6	C	65	Total	O	0	0
			65	65		
6	D	79	Total	O	0	0
			79	79		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SGR_2426




• Molecule 1: SGR_2426




• Molecule 1: SGR_2426



• Molecule 1: SGR_2426





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.86Å 97.76Å 184.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.26 – 2.20 46.26 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.26-2.20) 99.9 (46.26-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
R, R_{free}	0.232 , 0.257 (Not available) , 0.221	Depositor DCC
R_{free} test set	4438 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 23.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.035 for k,h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12655	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, PEG, BTB, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/3161	0.81	0/4326
1	B	0.47	0/3145	0.81	0/4305
1	C	0.47	0/3145	0.86	3/4305 (0.1%)
1	D	0.46	0/3145	0.82	0/4305
All	All	0.47	0/12596	0.82	3/17241 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	186	PHE	CB-CA-C	8.23	119.18	109.47
1	C	186	PHE	CA-CB-CG	6.14	119.94	113.80
1	C	235	PRO	N-CA-CB	-5.07	97.93	103.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3072	0	2907	25	0
1	B	3056	0	2887	14	0
1	C	3056	0	2887	21	0
1	D	3056	0	2887	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	7	0	10	1	0
3	A	14	0	19	0	0
3	B	14	0	19	0	0
3	C	28	0	38	4	0
3	D	14	0	19	0	0
4	B	10	0	14	0	0
4	D	10	0	14	0	0
5	B	16	0	22	0	0
6	A	78	0	0	0	0
6	B	80	0	0	0	0
6	C	65	0	0	0	0
6	D	79	0	0	0	0
All	All	12655	0	11723	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:MET:HE1	1:A:186:PHE:HE1	1.66	0.61
1:A:214:ARG:NH1	1:A:260:ASP:OD1	2.34	0.61
1:C:314:LEU:HB2	1:C:350:ILE:HG22	1.86	0.56
1:A:174:VAL:HG12	1:A:175:MET:HE2	1.89	0.55
1:D:271:ARG:HD2	1:D:278:ILE:HD11	1.89	0.55
1:B:64:TRP:N	1:B:65:PRO:CD	2.72	0.53
1:C:64:TRP:N	1:C:65:PRO:CD	2.71	0.52
1:A:64:TRP:N	1:A:65:PRO:CD	2.73	0.52
1:B:59:ASP:OD2	1:B:63:ARG:NH1	2.43	0.52
1:C:314:LEU:HB2	1:C:350:ILE:CG2	2.40	0.51
1:C:356:TRP:NE1	3:C:501:BTB:H42	2.25	0.51
1:C:170:ASN:HB3	1:C:252:PHE:CD2	2.45	0.51
1:C:169:PRO:HG3	1:C:224:TRP:CE3	2.45	0.51
1:C:267:TYR:OH	3:C:501:BTB:H41	2.10	0.51
1:A:15:PHE:HE1	1:A:401:LEU:HD12	1.75	0.51
1:B:170:ASN:HB3	1:B:252:PHE:CD2	2.47	0.50
1:B:322:ALA:HB2	1:B:371:PHE:CE1	2.45	0.50
1:C:129:TRP:O	1:C:133:ARG:HG3	2.12	0.50
1:A:170:ASN:HB3	1:A:252:PHE:CD2	2.47	0.49
1:B:289:THR:O	1:B:290:THR:OG1	2.24	0.49
1:A:53:PRO:HB2	1:A:55:LEU:HD13	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:64:TRP:N	1:D:65:PRO:CD	2.75	0.49
1:D:289:THR:O	1:D:290:THR:HB	2.14	0.48
1:A:263:GLY:HA2	1:A:315:ILE:O	2.12	0.48
1:A:16:PRO:O	1:A:19:PHE:HB2	2.15	0.47
1:D:104:ARG:HD3	1:D:154:PRO:O	2.14	0.47
1:C:356:TRP:CE2	3:C:501:BTB:H42	2.49	0.47
1:A:306:ALA:HB2	1:A:314:LEU:HD11	1.96	0.47
1:C:20:LEU:HB2	1:C:352:GLY:HA3	1.95	0.46
1:D:376:ILE:CD1	1:D:383:ARG:HD2	2.45	0.46
1:A:64:TRP:CG	1:A:65:PRO:HD3	2.49	0.46
1:D:169:PRO:HG3	1:D:224:TRP:CE3	2.50	0.46
1:A:15:PHE:CE1	1:A:401:LEU:CD1	2.99	0.46
1:B:373:LEU:HB3	1:B:389:ALA:HB2	1.98	0.46
1:B:230:VAL:HG12	1:B:230:VAL:O	2.17	0.45
1:D:129:TRP:O	1:D:133:ARG:HB2	2.17	0.45
1:B:20:LEU:HB2	1:B:352:GLY:HA3	1.98	0.44
1:B:356:TRP:HA	1:B:357:SER:HA	1.70	0.44
1:B:95:PHE:CE2	1:B:147:ARG:HD3	2.53	0.44
1:C:303:ARG:HA	1:C:348:LEU:HD21	2.00	0.44
1:A:373:LEU:HB3	1:A:389:ALA:HB2	2.00	0.44
1:C:234:LEU:O	1:C:235:PRO:C	2.61	0.43
1:C:264:VAL:HG21	1:C:302:LEU:HD23	2.00	0.43
1:A:153:ALA:N	1:A:154:PRO:HD2	2.33	0.43
1:B:265:GLN:HB3	1:B:318:GLU:HB2	2.00	0.43
1:C:373:LEU:HB3	1:C:389:ALA:HB2	2.00	0.43
1:D:20:LEU:HB2	1:D:352:GLY:HA3	2.00	0.43
1:A:39:TRP:O	1:A:43:GLU:HG3	2.19	0.43
1:A:231:TYR:CD1	1:A:272:ILE:HD11	2.54	0.43
1:D:153:ALA:N	1:D:154:PRO:HD2	2.33	0.43
3:C:501:BTB:O6	3:C:501:BTB:H81	2.19	0.42
1:A:356:TRP:HA	1:A:357:SER:HA	1.77	0.42
1:D:230:VAL:HG12	1:D:230:VAL:O	2.20	0.42
1:A:20:LEU:HB2	1:A:352:GLY:HA3	2.00	0.42
1:C:246:HIS:HA	1:C:250:ASP:HB2	2.01	0.42
1:C:356:TRP:HA	1:C:357:SER:HA	1.80	0.42
1:A:395:LEU:HA	1:A:398:THR:HB	2.02	0.42
1:D:20:LEU:HD22	1:D:78:ASP:OD2	2.20	0.42
1:A:225:THR:HG22	1:A:263:GLY:HA3	2.00	0.42
1:B:70:LEU:HD23	1:B:70:LEU:HA	1.91	0.41
1:B:166:ILE:O	1:B:224:TRP:HB2	2.20	0.41
1:C:246:HIS:N	1:C:247:PRO:HD2	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:356:TRP:HA	1:D:357:SER:HA	1.68	0.41
1:A:15:PHE:HE1	1:A:401:LEU:CD1	2.33	0.41
1:A:78:ASP:OD1	1:A:116:ARG:HB2	2.21	0.41
1:B:169:PRO:HG3	1:B:224:TRP:CE3	2.55	0.41
1:C:164:CYS:SG	1:C:225:THR:HG23	2.60	0.41
1:A:62:HIS:HA	2:A:501:PEG:H31	2.01	0.41
1:A:169:PRO:HD2	1:A:225:THR:O	2.20	0.41
1:C:374:ILE:CG2	1:C:383:ARG:HB3	2.51	0.41
1:C:11:THR:HG23	1:C:403:ARG:HG3	2.03	0.40
1:C:302:LEU:HD22	1:C:350:ILE:HG21	2.03	0.40
1:A:289:THR:O	1:A:290:THR:CB	2.69	0.40
1:D:133:ARG:NH2	1:D:139:GLU:O	2.41	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/414 (95%)	377 (96%)	15 (4%)	1 (0%)	36	42
1	B	391/414 (94%)	382 (98%)	9 (2%)	0	100	100
1	C	391/414 (94%)	375 (96%)	14 (4%)	2 (0%)	24	27
1	D	391/414 (94%)	375 (96%)	16 (4%)	0	100	100
All	All	1566/1656 (95%)	1509 (96%)	54 (3%)	3 (0%)	43	51

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	236	GLY
1	C	235	PRO
1	C	249	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/323 (94%)	291 (96%)	11 (4%)	31	42
1	B	305/323 (94%)	302 (99%)	3 (1%)	68	81
1	C	305/323 (94%)	294 (96%)	11 (4%)	31	42
1	D	305/323 (94%)	301 (99%)	4 (1%)	61	76
All	All	1217/1292 (94%)	1188 (98%)	29 (2%)	43	58

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LYS
1	A	11	THR
1	A	55	LEU
1	A	105	ARG
1	A	180	LYS
1	A	222	VAL
1	A	269	ARG
1	A	272	ILE
1	A	284	VAL
1	A	337	LEU
1	A	398	THR
1	B	155	VAL
1	B	175	MET
1	B	222	VAL
1	C	55	LEU
1	C	137	THR
1	C	139	GLU
1	C	160	VAL
1	C	186	PHE
1	C	235	PRO
1	C	278	ILE
1	C	284	VAL
1	C	350	ILE
1	C	379	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	398	THR
1	D	11	THR
1	D	83	VAL
1	D	186	PHE
1	D	238	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	265	GLN
1	A	319	ASN
1	B	94	HIS
1	B	265	GLN
1	C	265	GLN
1	C	361	ASN
1	D	361	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BTB	A	502	-	13,13,13	1.18	2 (15%)	7,16,16	0.57	0
4	PGE	B	501	-	9,9,9	0.16	0	8,8,8	0.12	0
3	BTB	C	502	-	13,13,13	1.12	3 (23%)	7,16,16	0.23	0
3	BTB	B	502	-	13,13,13	1.05	1 (7%)	7,16,16	0.21	0
5	1PE	B	503	-	15,15,15	0.19	0	14,14,14	0.10	0
2	PEG	A	501	-	6,6,6	0.14	0	5,5,5	0.11	0
3	BTB	C	501	-	13,13,13	1.35	3 (23%)	7,16,16	0.47	0
4	PGE	D	501	-	9,9,9	0.20	0	8,8,8	0.13	0
3	BTB	D	502	-	13,13,13	1.19	3 (23%)	7,16,16	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BTB	A	502	-	-	3/21/21/21	-
4	PGE	B	501	-	-	1/7/7/7	-
3	BTB	C	502	-	-	2/21/21/21	-
3	BTB	B	502	-	-	4/21/21/21	-
5	1PE	B	503	-	-	4/13/13/13	-
2	PEG	A	501	-	-	4/4/4/4	-
3	BTB	C	501	-	-	8/21/21/21	-
4	PGE	D	501	-	-	4/7/7/7	-
3	BTB	D	502	-	-	2/21/21/21	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	501	BTB	C2-N	3.33	1.55	1.48
3	D	502	BTB	C2-N	2.83	1.54	1.48
3	C	502	BTB	C2-N	2.56	1.53	1.48
3	C	501	BTB	C7-N	2.51	1.51	1.48
3	A	502	BTB	C2-N	2.49	1.53	1.48
3	A	502	BTB	C7-N	2.48	1.51	1.48
3	B	502	BTB	C2-N	2.41	1.53	1.48
3	C	501	BTB	C5-N	2.30	1.51	1.48
3	D	502	BTB	C7-N	2.15	1.51	1.48
3	D	502	BTB	C5-N	2.08	1.50	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	502	BTB	C5-N	2.08	1.50	1.48
3	C	502	BTB	C7-N	2.00	1.50	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (32) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	BTB	C1-C2-C3-O3
3	A	502	BTB	N-C2-C3-O3
3	A	502	BTB	C8-C7-N-C5
3	B	502	BTB	O1-C1-C2-C4
3	B	502	BTB	O1-C1-C2-N
3	C	501	BTB	C1-C2-N-C5
3	C	501	BTB	C1-C2-N-C7
3	C	501	BTB	C3-C2-N-C5
3	C	501	BTB	C3-C2-N-C7
3	C	501	BTB	C4-C2-N-C5
3	C	501	BTB	C4-C2-N-C7
3	C	501	BTB	C8-C7-N-C5
5	B	503	1PE	OH6-C15-C25-OH5
5	B	503	1PE	OH5-C14-C24-OH4
4	D	501	PGE	O3-C5-C6-O4
3	B	502	BTB	N-C5-C6-O6
3	D	502	BTB	N-C7-C8-O8
4	D	501	PGE	O2-C3-C4-O3
3	C	501	BTB	N-C5-C6-O6
2	A	501	PEG	O1-C1-C2-O2
2	A	501	PEG	O2-C3-C4-O4
4	D	501	PGE	O1-C1-C2-O2
5	B	503	1PE	OH2-C12-C22-OH3
3	B	502	BTB	N-C7-C8-O8
2	A	501	PEG	C4-C3-O2-C2
4	D	501	PGE	C4-C3-O2-C2
3	D	502	BTB	N-C5-C6-O6
2	A	501	PEG	C1-C2-O2-C3
3	C	502	BTB	N-C7-C8-O8
4	B	501	PGE	C6-C5-O3-C4
3	C	502	BTB	N-C5-C6-O6
5	B	503	1PE	OH4-C13-C23-OH3

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PEG	1	0
3	C	501	BTB	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/414 (95%)	0.33	17 (4%) 40 36	17, 23, 47, 54	0
1	B	393/414 (94%)	0.21	6 (1%) 72 69	17, 23, 36, 48	0
1	C	393/414 (94%)	0.54	19 (4%) 35 32	17, 27, 49, 63	0
1	D	393/414 (94%)	0.38	12 (3%) 51 48	16, 25, 45, 57	0
All	All	1574/1656 (95%)	0.37	54 (3%) 48 45	16, 24, 45, 63	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	11	THR	5.4
1	C	186	PHE	5.1
1	A	281	ALA	4.9
1	C	11	THR	4.7
1	D	12	PRO	3.9
1	D	235	PRO	3.7
1	C	280	ALA	3.7
1	D	13	VAL	3.6
1	C	284	VAL	3.4
1	D	403	ARG	3.3
1	B	11	THR	3.3
1	A	185	SER	3.3
1	A	236	GLY	3.2
1	C	282	ASP	3.0
1	B	399	ARG	3.0
1	A	183	ASP	3.0
1	A	237	ALA	2.9
1	C	236	GLY	2.8
1	C	278	ILE	2.8
1	C	281	ALA	2.8
1	A	179	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	311	ASP	2.7
1	D	278	ILE	2.7
1	A	10	LYS	2.6
1	C	283	ASP	2.6
1	D	281	ALA	2.6
1	D	186	PHE	2.5
1	B	185	SER	2.5
1	A	274	PRO	2.5
1	A	186	PHE	2.5
1	D	311	ASP	2.4
1	D	282	ASP	2.4
1	A	235	PRO	2.3
1	A	238	GLU	2.3
1	A	234	LEU	2.3
1	A	280	ALA	2.3
1	A	11	THR	2.3
1	C	158	GLU	2.3
1	C	12	PRO	2.2
1	B	139	GLU	2.2
1	C	401	LEU	2.2
1	B	183	ASP	2.2
1	D	275	GLU	2.2
1	C	402	PRO	2.2
1	C	398	THR	2.2
1	A	275	GLU	2.1
1	B	182	GLY	2.1
1	A	282	ASP	2.1
1	C	188	PRO	2.1
1	C	335	GLY	2.1
1	C	399	ARG	2.0
1	D	271	ARG	2.0
1	A	276	GLY	2.0
1	C	277	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BTB	A	502	14/14	0.77	0.16	29,31,33,33	0
4	PGE	D	501	10/10	0.77	0.19	38,39,39,39	0
5	1PE	B	503	16/16	0.79	0.19	49,51,52,52	0
3	BTB	B	502	14/14	0.81	0.16	31,32,34,35	0
3	BTB	C	501	14/14	0.84	0.13	29,29,31,31	0
2	PEG	A	501	7/7	0.89	0.12	28,28,28,28	0
4	PGE	B	501	10/10	0.89	0.10	27,28,28,28	0
3	BTB	D	502	14/14	0.91	0.10	28,28,29,29	0
3	BTB	C	502	14/14	0.91	0.10	26,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.