



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 10:33 AM UTC

PDB ID : 9KBH / pdb_00009kbh
EMDB ID : EMD-62225
Title : The structure of B19V NS1_2-570/ssDNA/AMPPNP
Authors : Gan, J.; Zhang, Y.
Deposited on : 2024-10-30
Resolution : 3.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	15087 (2.80 - 3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	569	5% 42% 8% 49%
1	B	569	45% 7% 48%
1	C	569	44% 7% 49%
1	D	569	47% 49%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	569	5%43%7%49%
1	F	569	23%39%10%51%
2	H	67	9%91%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Non-structural protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	290	Total	C	N	O	S	0	0
			2213	1433	365	403	12		
1	B	294	Total	C	N	O	S	0	0
			2285	1474	383	416	12		
1	C	292	Total	C	N	O	S	0	0
			2285	1473	383	416	13		
1	D	292	Total	C	N	O	S	0	0
			2285	1473	383	416	13		
1	E	292	Total	C	N	O	S	0	0
			2282	1471	383	416	12		
1	F	277	Total	C	N	O	S	0	0
			2105	1357	355	382	11		

- Molecule 2 is a DNA chain called DNA (67-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	6	Total	C	N	O	P	0	0
			122	60	18	38	6		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms					AltConf
3	A	1	Total 27	C 10	N 6	O 9	P 2	0
3	B	1	Total 31	C 10	N 6	O 12	P 3	0
3	C	1	Total 31	C 10	N 6	O 12	P 3	0
3	D	1	Total 31	C 10	N 6	O 12	P 3	0
3	E	1	Total 31	C 10	N 6	O 12	P 3	0
3	F	1	Total 31	C 10	N 6	O 12	P 3	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
4	B	1	Total Mg 1 1	0
4	C	1	Total Mg 1 1	0
4	D	1	Total Mg 1 1	0
4	E	1	Total Mg 1 1	0


- Molecule 1: Non-structural protein 1






• Molecule 1: Non-structural protein 1

Chain C: 44% 7% 49%




• Molecule 1: Non-structural protein 1


Chain D: 47% 49%



• Molecule 1: Non-structural protein 1



• Molecule 1: Non-structural protein 1



THR
THR
PRO
ILE
VAL
THR
ASP
THR
SER
ILE
SER
SER
SER
GLY
GLY
GLY
SER
SER
SER
GLU
GLU
LEU
SER
GLU
SER
SER
SER
PHE
PHE
ASN
LEU
ILE
THR
PRO
GLY
ALA
CYS
ASN
THR
GLU
THR
PRO
ARG
SER
SER
THR
PRO
ILE
PRO
GLY
THR
SER
SER
GLY
GLU
SER
LEU
VAL
GLY
SER
PRO
VAL

SER
SER
GLU
VAL
VAL
ALA
ALA
SER
TRP
GLU
GLU

- Molecule 2: DNA (67-MER)



DT
DG
DT
DA
DC
DG
DG
DA
DA
DG
DT
DC
DC
DC
DG
DC
DC
DA
DA
DC
DC
DG
DG
DC
DG
DG
DC
DC
DG
DA
DC
DG
DG
DC
DC
DG
DA
DT
DC
DT
DG
DA
DT
DT
DG
DG
DT
DG
DT
DC
DT
DT
DT
DT
DT



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	25268	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	2.342	Depositor
Minimum map value	-1.017	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.070	Depositor
Recommended contour level	0.45	Depositor
Map size ($\text{\AA}$)	345.24, 345.24, 345.24	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.959, 0.959, 0.959	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.15	0/2268	0.36	0/3093
1	B	0.14	0/2341	0.28	0/3189
1	C	0.16	0/2342	0.30	0/3188
1	D	0.14	0/2342	0.29	0/3188
1	E	0.20	0/2339	0.39	0/3185
1	F	0.18	0/2154	0.41	0/2936
2	H	0.55	0/135	0.54	0/206
All	All	0.17	0/13921	0.34	0/18985

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2213	0	2163	30	0
1	B	2285	0	2271	23	0
1	C	2285	0	2273	22	0
1	D	2285	0	2274	19	0
1	E	2282	0	2270	29	0
1	F	2105	0	2055	41	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	122	0	71	0	0
3	A	27	0	12	1	0
3	B	31	0	13	0	0
3	C	31	0	13	4	0
3	D	31	0	13	1	0
3	E	31	0	13	2	0
3	F	31	0	13	2	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
All	All	13763	0	13454	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:395:VAL:O	1:C:402:SER:HA	1.78	0.83
1:A:433:LYS:NZ	1:A:437:GLU:OE2	2.17	0.77
1:B:371:TRP:HB3	1:B:412:ILE:HG13	1.71	0.72
1:D:209:GLY:HA3	1:E:244:LEU:HD11	1.72	0.72
3:D:601:ANP:O2A	3:D:601:ANP:N3B	2.26	0.69

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	286/569 (50%)	263 (92%)	22 (8%)	1 (0%)	36 65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	290/569 (51%)	270 (93%)	20 (7%)	0	100	100
1	C	288/569 (51%)	272 (94%)	14 (5%)	2 (1%)	18	49
1	D	288/569 (51%)	271 (94%)	17 (6%)	0	100	100
1	E	288/569 (51%)	263 (91%)	20 (7%)	5 (2%)	7	30
1	F	273/569 (48%)	250 (92%)	22 (8%)	1 (0%)	30	60
All	All	1713/3414 (50%)	1589 (93%)	115 (7%)	9 (0%)	26	55

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	399	MET
1	C	402	SER
1	E	371	TRP
1	E	372	ASP
1	E	373	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/494 (47%)	231 (99%)	3 (1%)	61	74
1	B	249/494 (50%)	249 (100%)	0	100	100
1	C	251/494 (51%)	249 (99%)	2 (1%)	73	79
1	D	251/494 (51%)	251 (100%)	0	100	100
1	E	250/494 (51%)	244 (98%)	6 (2%)	43	65
1	F	221/494 (45%)	221 (100%)	0	100	100
All	All	1456/2964 (49%)	1445 (99%)	11 (1%)	70	79

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	369	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	370	VAL
1	E	396	ASP
1	E	375	ILE
1	C	438	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	397	GLN
1	E	477	ASN
1	F	352	ASN
1	E	443	ASN
1	C	290	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	B	601	4	33,33,33	0.93	4 (12%)	45,52,52	0.58	0
3	ANP	E	601	4	33,33,33	0.92	4 (12%)	45,52,52	0.61	1 (2%)
3	ANP	A	601	-	28,29,33	0.69	2 (7%)	40,45,52	0.74	1 (2%)
3	ANP	C	601	4	33,33,33	0.94	4 (12%)	45,52,52	0.59	0
3	ANP	F	601	-	33,33,33	0.93	4 (12%)	45,52,52	0.59	0
3	ANP	D	601	4	33,33,33	1.09	5 (15%)	45,52,52	0.94	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	B	601	4	-	6/18/38/38	0/3/3/3
3	ANP	E	601	4	-	7/18/38/38	0/3/3/3
3	ANP	A	601	-	-	4/13/32/38	0/3/3/3
3	ANP	C	601	4	-	5/18/38/38	0/3/3/3
3	ANP	F	601	-	-	7/18/38/38	0/3/3/3
3	ANP	D	601	4	-	9/18/38/38	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	601	ANP	PG-O3G	-2.81	1.49	1.56
3	D	601	ANP	PG-O1G	2.74	1.50	1.46
3	F	601	ANP	PG-O1G	2.57	1.50	1.46
3	C	601	ANP	PB-O3A	-2.57	1.55	1.59
3	D	601	ANP	PB-O1B	2.54	1.50	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	ANP	PB-O3A-PA	-4.21	118.63	132.10
3	D	601	ANP	O2B-PB-O1B	3.85	118.13	109.87
3	D	601	ANP	O3G-PG-O1G	-2.51	107.16	113.45
3	E	601	ANP	O1B-PB-N3B	-2.02	108.80	111.77

There are no chirality outliers.

5 of 38 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	ANP	C5'-O5'-PA-O1A
3	A	601	ANP	O4'-C4'-C5'-O5'
3	A	601	ANP	C3'-C4'-C5'-O5'
3	B	601	ANP	PB-N3B-PG-O1G
3	B	601	ANP	O4'-C4'-C5'-O5'

There are no ring outliers.


5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	601	ANP	2	0
3	A	601	ANP	1	0
3	C	601	ANP	4	0
3	F	601	ANP	2	0
3	D	601	ANP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand ANP E 601




Ligand ANP A 601









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-62225. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 187




Y Index: 195




Z Index: 170


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 359


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.45. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 83 nm³; this corresponds to an approximate mass of 75 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.303 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.303 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.30	3.97	3.39
Unmasked-calculated*	7.06	9.72	7.32

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.06 differs from the reported value 3.3 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-62225 and PDB model 9KBH. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.45 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.45).

9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.45) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7370	0.4850
A	0.7250	0.4450
B	0.8220	0.5130
C	0.8620	0.5310
D	0.8440	0.5260
E	0.6990	0.4970
F	0.4580	0.3960
H	0.5160	0.3510

1.0

0.0

<0.0