



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 06:05 PM UTC

PDB ID : 1KDX / pdb_00001kdx
Title : KIX DOMAIN OF MOUSE CBP (CREB BINDING PROTEIN) IN COMPLEX WITH PHOSPHORYLATED KINASE INDUCIBLE DOMAIN (PKID) OF RAT CREB (CYCLIC AMP RESPONSE ELEMENT BINDING PROTEIN), NMR 17 STRUCTURES
Authors : Radhakrishnan, I.; Perez-Alvarado, G.C.; Dyson, H.J.; Wright, P.E.
Deposited on : 1997-09-16

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	81	
2	B	28	

2 Ensemble composition and analysis

This entry contains 17 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:589-A:665, B:124-B:132, B:134-B:146 (99)	0.87	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 8, 10, 11, 15
2	12, 13, 17
3	7, 14
4	5, 9
Single-model clusters	4; 6; 16

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1809 atoms, of which 902 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CBP.

Mol	Chain	Residues	Atoms						Trace
1	A	81	Total	C	H	N	O	S	0
			1337	426	665	120	124	2	

- Molecule 2 is a protein called CREB.

Mol	Chain	Residues	Atoms						Trace
2	B	28	Total	C	H	N	O	P	0
			472	139	237	47	48	1	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	133	SEP	SER	engineered mutation	UNP P15337

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: CBP



- Molecule 2: CREB




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: CBP



- Molecule 2: CREB



4.2.2 Score per residue for model 2

- Molecule 1: CBP

Chain A:  85% 10% 5%



- Molecule 2: CREB

Chain B:  64% 14% 21%



4.2.3 Score per residue for model 3


- Molecule 1: CBP

Chain A:  83% 11% 5%



- Molecule 2: CREB

Chain B:  71% 7% 21%



4.2.4 Score per residue for model 4

- Molecule 1: CBP

Chain A:  86% 7% 5%



- Molecule 2: CREB


Chain B:  68% 11% 21%



4.2.5 Score per residue for model 5

- Molecule 1: CBP

Chain A:  85% 10% 5%



- Molecule 2: CREB

Chain B:  61% 14% 21%



4.2.6 Score per residue for model 6


- Molecule 1: CBP

Chain A:  81% 12% 5%



- Molecule 2: CREB

Chain B:  79% 21%



4.2.7 Score per residue for model 7


- Molecule 1: CBP

Chain A:  83% 11% 5%



- Molecule 2: CREB

Chain B:  75% 21%



4.2.8 Score per residue for model 8


- Molecule 1: CBP

Chain A:  86% 9% 5%



- Molecule 2: CREB

Chain B:  68% 11% 21%



4.2.9 Score per residue for model 9

- Molecule 1: CBP

Chain A:  88% 7% 5%



- Molecule 2: CREB

Chain B:  68% 11% 21%



4.2.10 Score per residue for model 10

- Molecule 1: CBP

Chain A:  85% 7% 5%




- Molecule 2: CREB

Chain B:  79% 21%



4.2.11 Score per residue for model 11

- Molecule 1: CBP




- Molecule 2: CREB




4.2.12 Score per residue for model 12

- Molecule 1: CBP




- Molecule 2: CREB




4.2.13 Score per residue for model 13

- Molecule 1: CBP



- Molecule 2: CREB



4.2.14 Score per residue for model 14

- Molecule 1: CBP

Chain A:  88% 7% 5%



- Molecule 2: CREB

Chain B:  68% 11% 21%



4.2.15 Score per residue for model 15


- Molecule 1: CBP

Chain A:  83% 11% 5%



- Molecule 2: CREB


Chain B:  75% 0% 21%



4.2.16 Score per residue for model 16


- Molecule 1: CBP

Chain A:  88% 6% 5%



- Molecule 2: CREB

Chain B:  68% 11% 21%



4.2.17 Score per residue for model 17

- Molecule 1: CBP

Chain A:  85% 10% 5%



- Molecule 2: CREB

Chain B:  79% 21%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DG*, *SA*.

Of the 40 calculated structures, 17 were deposited, based on the following criterion: *LOWEST CONSTRAINT ENERGIES*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DIANA	refinement	
MSI FELIX95	structure solution	FELIX95

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.64±0.01	0±0/657 (0.0± 0.0%)	1.25±0.03	1±1/887 (0.1± 0.1%)
2	B	0.59±0.01	0±0/187 (0.0± 0.0%)	1.09±0.04	0±0/248 (0.0± 0.0%)
All	All	0.63	0/14348 (0.0%)	1.22	11/19295 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.0±0.8
2	B	0.0±0.0	0.9±0.8
All	All	0	67

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	661	GLN	CA-C-N	5.84	128.59	120.29	7	1
1	A	661	GLN	C-N-CA	5.84	128.59	120.29	7	1
1	A	651	HIS	CB-CG-CD2	-5.45	124.11	131.20	7	7
1	A	612	PHE	CA-CB-CG	5.03	118.83	113.80	10	1
1	A	605	HIS	CB-CG-CD2	-5.02	124.67	131.20	11	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	649	TYR	Sidechain	17
1	A	612	PHE	Sidechain	12
2	B	134	TYR	Sidechain	10
1	A	640	TYR	Sidechain	9
1	A	650	TYR	Sidechain	6
1	A	658	TYR	Sidechain	4
2	B	130	ARG	Sidechain	2
2	B	125	ARG	Sidechain	2
2	B	135	ARG	Sidechain	1
2	B	131	ARG	Sidechain	1
1	A	600	ARG	Sidechain	1
1	A	646	ARG	Sidechain	1
1	A	624	ARG	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	641	634	634	0±1
2	B	186	196	196	0±0
All	All	14059	14110	14110	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:599:LEU:C	1:A:599:LEU:HD13	0.48	2.33	12	1
2:B:134:TYR:CE2	2:B:138:LEU:HD11	0.46	2.45	5	1
2:B:141:LEU:HD23	2:B:141:LEU:C	0.44	2.37	16	1
1:A:595:VAL:HG13	1:A:599:LEU:HD12	0.42	1.91	12	1
1:A:602:HIS:CD2	1:A:602:HIS:C	0.41	2.98	15	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	77/81 (95%)	69±2 (89±3%)	7±2 (9±2%)	1±1 (2±1%)	9	53
2	B	21/28 (75%)	19±1 (91±6%)	2±1 (8±6%)	0±0 (1±2%)	18	68
All	All	1666/1853 (90%)	1493 (90%)	148 (9%)	25 (2%)	11	57

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	642	SER	7
1	A	590	GLY	5
1	A	613	PRO	2
1	A	589	LYS	2
1	A	591	TRP	2
1	A	615	PRO	1
2	B	131	ARG	1
2	B	126	GLU	1
2	B	130	ARG	1
1	A	614	THR	1
1	A	624	ARG	1
1	A	645	SER	1

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	68/71 (96%)	64±1 (94±2%)	4±1 (6±2%)	19	68
2	B	21/26 (81%)	20±1 (95±5%)	1±1 (5±5%)	24	75
All	All	1513/1649 (92%)	1426 (94%)	87 (6%)	20	69

All 31 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	606	LYS	10
1	A	644	ASN	9
1	A	589	LYS	8
1	A	623	ARG	6
1	A	650	TYR	5
1	A	651	HIS	5
2	B	142	SER	4
1	A	627	ASN	4
2	B	126	GLU	3
1	A	592	HIS	2
1	A	614	THR	2
2	B	125	ARG	2
1	A	597	GLN	2
1	A	602	HIS	2
1	A	660	ILE	2
2	B	135	ARG	2
2	B	134	TYR	2
1	A	616	ASP	2
1	A	620	LEU	2
1	A	593	GLU	2
1	A	605	HIS	1
1	A	663	GLU	1
2	B	124	ARG	1
1	A	656	LYS	1
1	A	624	ARG	1
1	A	621	LYS	1
2	B	136	LYS	1
2	B	130	ARG	1
2	B	128	LEU	1
1	A	646	ARG	1
1	A	662	LYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	SEP	B	133	2	8,9,10	1.04±0.02	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	SEP	B	133	2	7,12,14	1.95±0.33	2±1 (22±9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	B	133	2	-	0±0,6,8,10	-

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	133	SEP	OG-CB-CA	5.71	113.70	108.14	1	16
2	B	133	SEP	O3P-P-OG	2.15	101.06	106.67	9	1
2	B	133	SEP	O3P-P-O2P	2.08	115.59	107.80	15	10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided