



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 12:05 PM UTC

PDB ID : 5KDS / pdb_00005kds
Title : ZmpB metallopeptidase in complex with an O-glycopeptide (a2,6-sialylated core-3 pentapeptide).
Authors : Noach, I.; Ficko-Blean, E.; Stuart, C.; Boraston, A.B.
Deposited on : 2016-06-08
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	5	60% 40% 60%
2	A	530	3% 90% 5% • •
3	B	3	33% 67%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha2,6-sialylated core-3 pentapeptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	G	5	Total	C	N	O	0	0	0
			27	16	5	6			

- Molecule 2 is a protein called F5/8 type C domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	511	Total	C	N	O	S	0	7	0
			4127	2618	691	807	11			

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	474	MET	-	initiating methionine	UNP A0A0H2YN38
A	475	GLY	-	expression tag	UNP A0A0H2YN38
A	476	SER	-	expression tag	UNP A0A0H2YN38
A	477	SER	-	expression tag	UNP A0A0H2YN38
A	478	HIS	-	expression tag	UNP A0A0H2YN38
A	479	HIS	-	expression tag	UNP A0A0H2YN38
A	480	HIS	-	expression tag	UNP A0A0H2YN38
A	481	HIS	-	expression tag	UNP A0A0H2YN38
A	482	HIS	-	expression tag	UNP A0A0H2YN38
A	483	HIS	-	expression tag	UNP A0A0H2YN38
A	484	SER	-	expression tag	UNP A0A0H2YN38
A	485	SER	-	expression tag	UNP A0A0H2YN38
A	486	GLY	-	expression tag	UNP A0A0H2YN38
A	487	LEU	-	expression tag	UNP A0A0H2YN38
A	488	VAL	-	expression tag	UNP A0A0H2YN38
A	489	PRO	-	expression tag	UNP A0A0H2YN38
A	490	ARG	-	expression tag	UNP A0A0H2YN38
A	491	GLY	-	expression tag	UNP A0A0H2YN38
A	492	SER	-	expression tag	UNP A0A0H2YN38
A	493	HIS	-	expression tag	UNP A0A0H2YN38

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	494	MET	-	expression tag	UNP A0A0H2YN38
A	495	ALA	-	expression tag	UNP A0A0H2YN38
A	496	SER	-	expression tag	UNP A0A0H2YN38

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	3	Total	C	N	O	0	0	0
			48	27	3	18			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: C₄H₆O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	8	Total	O	0	0
			8	8		
8	A	786	Total	O	0	2
			788	788		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha2,6-sialylated core-3 pentapeptide



- Molecule 2: F5/8 type C domain protein



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-[N-acetyl-alpha-neuraminic acid-(2-6)]2-acetamido-2-deoxy-alpha-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.18Å 68.95Å 171.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.60 40.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	95.9 (40.00-1.60) 95.9 (40.00-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.83 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.158 , 0.185 0.159 , 0.187	Depositor DCC
R_{free} test set	4992 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5044	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, TLA, EDO, NAG, A2G, EPE, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	1.05	0/27	0.99	0/36
2	A	0.90	2/4241 (0.0%)	0.94	1/5736 (0.0%)
All	All	0.90	2/4268 (0.0%)	0.94	1/5772 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	935	ASN	CA-C	5.65	1.60	1.52
2	A	680	THR	N-CA	5.01	1.49	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	696	MET	CA-CB-CG	-5.37	103.36	114.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	934	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	27	0	23	6	0
2	A	4127	0	4006	23	0
3	B	48	0	40	5	0
4	A	1	0	0	0	0
5	A	10	0	3	2	0
6	A	15	0	18	1	0
7	A	20	0	30	0	0
8	A	788	0	0	3	0
8	G	8	0	0	4	0
All	All	5044	0	4120	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:G:202:HOH:O	3:B:2:NAG:H61	1.83	0.78
2:A:784:LYS:HE3	8:A:1799:HOH:O	1.85	0.76
2:A:493:HIS:CE1	2:A:618:LYS:HD2	2.24	0.72
2:A:859:GLU:H	2:A:859:GLU:CD	2.05	0.64
1:G:4:THR:HA	5:A:1102:TLA:O11	2.05	0.56
1:G:4:THR:N	5:A:1102:TLA:HB	2.04	0.55
1:G:6:PRO:CD	2:A:725:GLY:O	2.58	0.52
1:G:6:PRO:HD3	2:A:725:GLY:O	2.11	0.51
2:A:610:LYS:NZ	8:A:1206:HOH:O	2.44	0.50
2:A:680:THR:HB	2:A:681:PRO:HD2	1.94	0.50
2:A:539:ASP:OD1	2:A:607:ARG:NH2	2.32	0.49
2:A:723:SER:HA	2:A:738:ARG:NH2	2.28	0.48
2:A:1002:VAL:O	2:A:1003:ASN:CB	2.62	0.48
2:A:978:ASN:O	2:A:981:GLU:HG2	2.13	0.48
8:G:201:HOH:O	3:B:3:SIA:C7	2.62	0.47
2:A:890:ASP:O	2:A:894:GLU:HG3	2.15	0.47
8:G:202:HOH:O	3:B:2:NAG:C6	2.54	0.46
1:G:7:GLY:HA2	3:B:2:NAG:O6	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:676:LYS:HE2	8:A:1203:HOH:O	2.16	0.46
2:A:713:PHE:CE1	2:A:715[A]:ILE:HG12	2.51	0.46
2:A:985:TYR:HB2	2:A:1002:VAL:HB	2.00	0.44
2:A:713:PHE:HE1	2:A:715[A]:ILE:HG12	1.84	0.43
1:G:6:PRO:HD2	2:A:725:GLY:O	2.18	0.43
2:A:1002:VAL:O	2:A:1003:ASN:CG	2.61	0.43
6:A:1103:EPE:H102	6:A:1103:EPE:H62	1.50	0.42
8:G:201:HOH:O	3:B:3:SIA:H7	2.20	0.42
2:A:587:LYS:HD2	2:A:587:LYS:HA	1.60	0.41
2:A:493:HIS:CG	2:A:618:LYS:HG3	2.56	0.41
2:A:687:GLN:HB3	2:A:748:ALA:HB3	2.03	0.41
2:A:716:MET:HE3	2:A:734:VAL:HG22	2.04	0.40
2:A:539:ASP:OD1	2:A:539:ASP:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	3/5 (60%)	3 (100%)	0	0	100	100
2	A	516/530 (97%)	506 (98%)	10 (2%)	0	100	100
All	All	519/535 (97%)	509 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	2/2 (100%)	2 (100%)	0	100	100
2	A	449/458 (98%)	445 (99%)	4 (1%)	70	55
All	All	451/460 (98%)	447 (99%)	4 (1%)	70	55

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	587	LYS
2	A	699	TRP
2	A	859	GLU
2	A	952	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	493	HIS
2	A	703	ASN
2	A	712	ASN
2	A	799	ASN
2	A	935	ASN
2	A	995	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A2G	B	1	3,1	14,14,15	1.21	2 (14%)	17,19,21	1.50	3 (17%)
3	NAG	B	2	3	14,14,15	0.51	0	17,19,21	1.27	3 (17%)
3	SIA	B	3	3	20,20,21	0.80	0	21,28,31	1.34	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A2G	B	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	B	2	3	-	2/6/23/26	0/1/1/1
3	SIA	B	3	3	-	0/18/34/38	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1	A2G	C1-C2	2.83	1.56	1.52
3	B	1	A2G	O5-C1	-2.20	1.40	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1	A2G	O5-C5-C6	-4.21	99.47	107.66
3	B	1	A2G	C2-N2-C7	-2.92	118.99	122.90
3	B	2	NAG	O5-C1-C2	-2.44	107.52	111.29
3	B	3	SIA	O1A-C1-C2	-2.40	117.67	122.85
3	B	2	NAG	C6-C5-C4	-2.34	107.27	113.02
3	B	3	SIA	C9-C8-C7	-2.31	107.47	112.17
3	B	1	A2G	O6-C6-C5	-2.26	103.62	111.33
3	B	2	NAG	C3-C4-C5	2.26	114.33	110.23
3	B	3	SIA	O1B-C1-C2	2.23	118.51	112.71

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	2	NAG	C4-C5-C6-O6
3	B	2	NAG	O5-C5-C6-O6
3	B	1	A2G	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2	NAG	3	0
3	B	3	SIA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	1106	-	3,3,3	0.43	0	2,2,2	0.40	0
5	TLA	A	1102	4	9,9,9	1.34	1 (11%)	12,12,12	1.18	1 (8%)
7	EDO	A	1105	-	3,3,3	0.43	0	2,2,2	0.52	0
6	EPE	A	1103	-	15,15,15	1.89	2 (13%)	19,20,20	1.53	5 (26%)
7	EDO	A	1104	-	3,3,3	0.52	0	2,2,2	0.67	0
7	EDO	A	1108	-	3,3,3	0.46	0	2,2,2	0.80	0
7	EDO	A	1107	-	3,3,3	0.42	0	2,2,2	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	A	1106	-	-	0/1/1/1	-
5	TLA	A	1102	4	-	0/12/12/12	-
7	EDO	A	1105	-	-	0/1/1/1	-
6	EPE	A	1103	-	-	4/9/19/19	0/1/1/1
7	EDO	A	1104	-	-	0/1/1/1	-
7	EDO	A	1108	-	-	1/1/1/1	-
7	EDO	A	1107	-	-	1/1/1/1	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1103	EPE	C10-S	-6.13	1.69	1.77
6	A	1103	EPE	O2S-S	2.43	1.52	1.45
5	A	1102	TLA	O2-C2	2.13	1.46	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1103	EPE	C6-N1-C2	3.40	116.17	108.84
6	A	1103	EPE	C5-C6-N1	2.92	116.54	110.65
6	A	1103	EPE	C3-C2-N1	2.55	115.80	110.65
6	A	1103	EPE	O1S-S-C10	2.24	110.12	106.73
5	A	1102	TLA	O11-C1-C2	2.22	119.48	113.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1103	EPE	O2S-S-O1S	-2.13	106.89	113.82

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1103	EPE	C10-C9-N1-C6
6	A	1103	EPE	C8-C7-N4-C5
6	A	1103	EPE	S-C10-C9-N1
6	A	1103	EPE	C10-C9-N1-C2
7	A	1107	EDO	O1-C1-C2-O2
7	A	1108	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1102	TLA	2	0
6	A	1103	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	5/5 (100%)	2.49	3 (60%) 0 0	23, 34, 75, 79	0
2	A	511/530 (96%)	-0.32	15 (2%) 53 56	8, 18, 36, 65	7 (1%)
All	All	516/535 (96%)	-0.29	18 (3%) 47 49	8, 18, 37, 79	7 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	1003	ASN	5.7
2	A	494	MET	5.1
2	A	725	GLY	4.8
2	A	495	ALA	4.8
2	A	493	HIS	4.4
2	A	1002	VAL	3.8
1	G	8	GLY	3.7
1	G	6	PRO	3.4
1	G	7	GLY	3.3
2	A	724	GLY	3.2
2	A	723	SER	2.8
2	A	917	LYS	2.8
2	A	961	GLY	2.7
2	A	859	GLU	2.6
2	A	726	ALA	2.6
2	A	982	ASP	2.4
2	A	919	ASP	2.2
2	A	894	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	A2G	B	1	14/15	-	-	18,20,22,24	0
3	NAG	B	2	14/15	-	-	28,41,49,55	0
3	SIA	B	3	20/21	-	-	19,26,37,37	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	EDO	A	1108	4/4	0.85	0.13	37,37,37,43	0
7	EDO	A	1107	4/4	0.87	0.13	34,36,38,43	0
6	EPE	A	1103	15/15	0.89	0.21	25,65,86,86	0
7	EDO	A	1106	4/4	0.89	0.13	35,37,37,40	0
5	TLA	A	1102	10/10	0.93	0.08	26,32,36,37	0
7	EDO	A	1104	4/4	0.95	0.09	22,29,30,38	0
7	EDO	A	1105	4/4	0.97	0.07	24,29,31,32	0
4	ZN	A	1101	1/1	1.00	0.13	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.