



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 08:06 AM UTC

PDB ID : 6KDM / pdb_00006kdm
Title : HIV-1 reverse transcriptase with Q151M/Y115F/F116Y:DNA:entecavir
5'-triphosphate ternary complex
Authors : Yasutake, Y.; Hattori, S.I.; Tamura, N.; Maeda, K.
Deposited on : 2019-07-02
Resolution : 2.32 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 180053 | 7754 (2.34-2.30) |
| Clashscore | 190562 | 8383 (2.34-2.30) |
| Ramachandran outliers | 187476 | 8303 (2.34-2.30) |
| Sidechain outliers | 187428 | 8303 (2.34-2.30) |
| RSRZ outliers | 180081 | 7760 (2.34-2.30) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--------------------|
| 1 | A | 557 | 11% 82% 17% |
| 1 | C | 557 | 5% 89% 10% |
| 2 | B | 444 | 18% 77% 14% 9% |
| 2 | D | 444 | 8% 79% 11% 9% |
| 3 | E | 38 | 71% 16% 5% 8% |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 3 | F | 38 |  74% 21% 5% |

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 17780 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 reverse transcriptase p66 subunit.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 553 | Total | C | N | O | S | 0 | 1 | 0 |
| | | | 4503 | 2914 | 750 | 831 | 8 | | | |
| 1 | C | 553 | Total | C | N | O | S | 0 | 1 | 0 |
| | | | 4503 | 2915 | 750 | 829 | 9 | | | |

There are 14 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| A | -1 | MET | - | expression tag | UNP D3XFN5 |
| A | 0 | VAL | - | expression tag | UNP D3XFN5 |
| A | 115 | PHE | TYR | engineered mutation | UNP D3XFN5 |
| A | 116 | TYR | PHE | engineered mutation | UNP D3XFN5 |
| A | 151 | MET | GLN | engineered mutation | UNP D3XFN5 |
| A | 162 | SER | CYS | engineered mutation | UNP D3XFN5 |
| A | 280 | SER | CYS | engineered mutation | UNP D3XFN5 |
| C | -1 | MET | - | expression tag | UNP D3XFN5 |
| C | 0 | VAL | - | expression tag | UNP D3XFN5 |
| C | 115 | PHE | TYR | engineered mutation | UNP D3XFN5 |
| C | 116 | TYR | PHE | engineered mutation | UNP D3XFN5 |
| C | 151 | MET | GLN | engineered mutation | UNP D3XFN5 |
| C | 162 | SER | CYS | engineered mutation | UNP D3XFN5 |
| C | 280 | SER | CYS | engineered mutation | UNP D3XFN5 |

- Molecule 2 is a protein called HIV-1 RT p51 subunit.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 2 | B | 406 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 3347 | 2178 | 557 | 606 | 6 | | | |
| 2 | D | 406 | Total | C | N | O | S | 0 | 1 | 0 |
| | | | 3352 | 2182 | 557 | 606 | 7 | | | |

There are 36 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| B | -15 | MET | - | expression tag | UNP P12497 |
| B | -14 | ALA | - | expression tag | UNP P12497 |
| B | -13 | HIS | - | expression tag | UNP P12497 |
| B | -12 | HIS | - | expression tag | UNP P12497 |
| B | -11 | HIS | - | expression tag | UNP P12497 |
| B | -10 | HIS | - | expression tag | UNP P12497 |
| B | -9 | HIS | - | expression tag | UNP P12497 |
| B | -8 | HIS | - | expression tag | UNP P12497 |
| B | -7 | ALA | - | expression tag | UNP P12497 |
| B | -6 | LEU | - | expression tag | UNP P12497 |
| B | -5 | GLU | - | expression tag | UNP P12497 |
| B | -4 | VAL | - | expression tag | UNP P12497 |
| B | -3 | LEU | - | expression tag | UNP P12497 |
| B | -2 | PHE | - | expression tag | UNP P12497 |
| B | -1 | GLN | - | expression tag | UNP P12497 |
| B | 0 | GLY | - | expression tag | UNP P12497 |
| B | 162 | SER | CYS | engineered mutation | UNP P12497 |
| B | 280 | SER | CYS | engineered mutation | UNP P12497 |
| D | -15 | MET | - | expression tag | UNP P12497 |
| D | -14 | ALA | - | expression tag | UNP P12497 |
| D | -13 | HIS | - | expression tag | UNP P12497 |
| D | -12 | HIS | - | expression tag | UNP P12497 |
| D | -11 | HIS | - | expression tag | UNP P12497 |
| D | -10 | HIS | - | expression tag | UNP P12497 |
| D | -9 | HIS | - | expression tag | UNP P12497 |
| D | -8 | HIS | - | expression tag | UNP P12497 |
| D | -7 | ALA | - | expression tag | UNP P12497 |
| D | -6 | LEU | - | expression tag | UNP P12497 |
| D | -5 | GLU | - | expression tag | UNP P12497 |
| D | -4 | VAL | - | expression tag | UNP P12497 |
| D | -3 | LEU | - | expression tag | UNP P12497 |
| D | -2 | PHE | - | expression tag | UNP P12497 |
| D | -1 | GLN | - | expression tag | UNP P12497 |
| D | 0 | GLY | - | expression tag | UNP P12497 |
| D | 162 | SER | CYS | engineered mutation | UNP P12497 |
| D | 280 | SER | CYS | engineered mutation | UNP P12497 |

- Molecule 3 is a DNA chain called DNA/RNA (38-MER).


| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|---------|-------|
| | | | Total | C | N | O | P | | | |
| 3 | E | 35 | 718 | 339 | 128 | 216 | 35 | 0 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|---------|-------|
| | | | Total | C | N | O | P | | | |
| 3 | F | 38 | 777 | 369 | 140 | 231 | 37 | 0 | 0 | 0 |

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| | | | Total | C | O | | |
| 4 | A | 1 | 6 | 3 | 3 | 0 | 0 |
| 4 | B | 1 | 6 | 3 | 3 | 0 | 0 |
| 4 | E | 1 | 6 | 3 | 3 | 0 | 0 |
| 4 | D | 1 | 6 | 3 | 3 | 0 | 0 |
| 4 | D | 1 | 6 | 3 | 3 | 0 | 0 |
| 4 | D | 1 | 6 | 3 | 3 | 0 | 0 |

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| | | | Total | Mg | | |
| 5 | A | 1 | 1 | 1 | 0 | 0 |
| 5 | C | 1 | 1 | 1 | 0 | 0 |

- Molecule 6 is [[(1R,3S,5S)-3-(2-azanyl-6-oxidanylidene-3H-purin-9-yl)-2-methylidene-5-oxidanyl-cyclopentyl]methoxy-oxidanyl-phosphoryl l] phosphono hydrogen phosphate (CCD ID: ET9) (formula: C₁₂H₁₈N₅O₁₂P₃) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | |
|-----|-------|----------|-------|----|----|---|---------|---------|---|
| 6 | A | 1 | Total | C | N | O | P | 0 | 0 |
| | | | | 32 | 12 | 5 | 12 | | |
| 6 | C | 1 | Total | C | N | O | P | 0 | 0 |
| | | | | 32 | 12 | 5 | 12 | | |


- Molecule 7 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 7 | A | 133 | Total | O | 0 | 0 |
| | | | | 133 | | |
| 7 | B | 53 | Total | O | 0 | 0 |
| | | | | 53 | | |
| 7 | E | 42 | Total | O | 0 | 0 |
| | | | | 42 | | |
| 7 | C | 116 | Total | O | 0 | 0 |
| | | | | 116 | | |
| 7 | D | 93 | Total | O | 0 | 0 |
| | | | | 93 | | |
| 7 | F | 41 | Total | O | 0 | 0 |
| | | | | 41 | | |


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HIV-1 reverse transcriptase p66 subunit




- Molecule 1: HIV-1 reverse transcriptase p66 subunit





- Molecule 2: HIV-1 RT p51 subunit





• Molecule 2: HIV-1 RT p51 subunit



• Molecule 3: DNA/RNA (38-MER)



• Molecule 3: DNA/RNA (38-MER)



4 Data and refinement statistics i

| Property | Value | Source |
|---|---|------------------|
| Space group | H 3 | Depositor |
| Cell constants a, b, c, α , β , γ | 284.45Å 284.45Å 95.77Å 90.00° 90.00° 120.00° | Depositor |
| Resolution (Å) | 48.67 – 2.32 48.67 – 2.32 | Depositor EDS |
| % Data completeness (in resolution range) | 100.0 (48.67-2.32) 99.9 (48.67-2.32) | Depositor EDS |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 2.17 (at 2.32Å) | Xtrriage |
| Refinement program | PHENIX (1.14_3260: ???) | Depositor |
| R, R_{free} | 0.190 , 0.223 0.192 , 0.224 | Depositor DCC |
| R_{free} test set | 6352 reflections (5.08%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 44.1 | Xtrriage |
| Anisotropy | 0.113 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.35 , 48.3 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$ | Xtrriage |
| Estimated twinning fraction | 0.010 for h,-h-k,-l | Xtrriage |
| F_o, F_c correlation | 0.95 | EDS |
| Total number of atoms | 17780 | wwPDB-VP |
| Average B, all atoms (Å ²) | 58.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ET9, GOL, OMC, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.30 | 1/4624 (0.0%) | 0.49 | 7/6279 (0.1%) |
| 1 | C | 0.27 | 2/4624 (0.0%) | 0.40 | 2/6278 (0.0%) |
| 2 | B | 0.31 | 0/3441 | 0.42 | 2/4673 (0.0%) |
| 2 | D | 0.39 | 2/3449 (0.1%) | 0.43 | 2/4683 (0.0%) |
| 3 | E | 0.20 | 0/756 | 0.40 | 0/1165 |
| 3 | F | 0.19 | 0/823 | 0.39 | 0/1269 |
| All | All | 0.31 | 5/17717 (0.0%) | 0.43 | 13/24347 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | C | 0 | 2 |
| 2 | D | 0 | 1 |
| All | All | 0 | 3 |

All (5) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|--------|------|-------|-------|-------------|----------|
| 1 | C | 357[A] | MET | N-CA | 6.13 | 1.54 | 1.46 |
| 1 | C | 357[B] | MET | N-CA | 6.13 | 1.54 | 1.46 |
| 2 | D | 235 | HIS | C-O | -5.55 | 1.17 | 1.24 |
| 1 | A | 447 | ASN | C-O | -5.48 | 1.17 | 1.24 |
| 2 | D | 235 | HIS | CA-C | -5.19 | 1.46 | 1.52 |

The worst 5 of 13 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|--------|------|--------|--------|-------------|----------|
| 1 | A | 67 | ASP | N-CA-C | -10.88 | 93.60 | 109.59 |
| 1 | A | 68 | SER | N-CA-C | 7.45 | 119.40 | 111.28 |
| 1 | A | 452 | LEU | N-CA-C | 6.91 | 121.31 | 110.32 |
| 1 | C | 357[A] | MET | N-CA-C | 6.63 | 119.65 | 107.99 |
| 1 | C | 357[B] | MET | N-CA-C | 6.63 | 119.65 | 107.99 |

There are no chirality outliers.

All (3) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|--------|------|-----------|
| 1 | C | 357[A] | MET | Mainchain |
| 1 | C | 357[B] | MET | Mainchain |
| 2 | D | 425 | LEU | Mainchain |

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 4503 | 0 | 4557 | 65 | 0 |
| 1 | C | 4503 | 0 | 4562 | 40 | 0 |
| 2 | B | 3347 | 0 | 3379 | 45 | 0 |
| 2 | D | 3352 | 0 | 3388 | 45 | 0 |
| 3 | E | 718 | 0 | 397 | 8 | 0 |
| 3 | F | 777 | 0 | 432 | 8 | 0 |
| 4 | A | 6 | 0 | 8 | 0 | 0 |
| 4 | B | 6 | 0 | 8 | 0 | 0 |
| 4 | D | 18 | 0 | 24 | 2 | 0 |
| 4 | E | 6 | 0 | 8 | 3 | 0 |
| 5 | A | 1 | 0 | 0 | 0 | 0 |
| 5 | C | 1 | 0 | 0 | 0 | 0 |
| 6 | A | 32 | 0 | 0 | 2 | 0 |
| 6 | C | 32 | 0 | 0 | 0 | 0 |
| 7 | A | 133 | 0 | 0 | 4 | 0 |
| 7 | B | 53 | 0 | 0 | 0 | 0 |
| 7 | C | 116 | 0 | 0 | 1 | 0 |
| 7 | D | 93 | 0 | 0 | 0 | 0 |
| 7 | E | 42 | 0 | 0 | 0 | 0 |
| 7 | F | 41 | 0 | 0 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| All | All | 17780 | 0 | 16763 | 201 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 201 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 1:A:450:THR:HG22 | 1:A:451:LYS:H | 1.19 | 1.06 |
| 2:D:425:LEU:H | 2:D:425:LEU:CD1 | 1.78 | 0.95 |
| 2:D:320:ASP:H | 2:D:343:GLN:HE22 | 1.10 | 0.92 |
| 1:A:450:THR:HG22 | 1:A:451:LYS:N | 1.77 | 0.91 |
| 1:A:222:GLN:HE22 | 1:A:228:LEU:H | 1.19 | 0.90 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|---------|----------|-------------|-----|
| 1 | A | 552/557 (99%) | 536 (97%) | 15 (3%) | 1 (0%) | 43 | 53 |
| 1 | C | 552/557 (99%) | 535 (97%) | 15 (3%) | 2 (0%) | 30 | 37 |
| 2 | B | 402/444 (90%) | 383 (95%) | 18 (4%) | 1 (0%) | 43 | 53 |
| 2 | D | 403/444 (91%) | 388 (96%) | 15 (4%) | 0 | 100 | 100 |
| All | All | 1909/2002 (95%) | 1842 (96%) | 63 (3%) | 4 (0%) | 43 | 53 |

All (4) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 136 | ASN |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | C | 137 | ASN |
| 2 | B | 282 | LEU |
| 1 | C | 133 | PRO |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|----|
| 1 | A | 493/494 (100%) | 490 (99%) | 3 (1%) | 78 | 88 |
| 1 | C | 493/494 (100%) | 491 (100%) | 2 (0%) | 84 | 91 |
| 2 | B | 365/400 (91%) | 359 (98%) | 6 (2%) | 55 | 72 |
| 2 | D | 366/400 (92%) | 362 (99%) | 4 (1%) | 65 | 80 |
| All | All | 1717/1788 (96%) | 1702 (99%) | 15 (1%) | 70 | 83 |

5 of 15 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 356 | ARG |
| 2 | D | 424 | LYS |
| 2 | B | 422 | LEU |
| 2 | D | 425 | LEU |
| 2 | D | 212 | TRP |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | D | 175 | ASN |
| 2 | D | 182 | GLN |
| 2 | D | 367 | GLN |
| 2 | B | 161 | GLN |
| 2 | B | 151 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 3 | OMC | E | 4 | 3 | 19,22,23 | 2.85 | 8 (42%) | 25,31,34 | 0.79 | 0 |
| 3 | OMC | F | 2 | 3 | 19,22,23 | 2.83 | 8 (42%) | 25,31,34 | 0.75 | 0 |
| 3 | OMC | F | 4 | 3 | 19,22,23 | 2.87 | 8 (42%) | 25,31,34 | 0.66 | 0 |
| 3 | OMC | E | 2 | 3 | 19,22,23 | 2.81 | 8 (42%) | 25,31,34 | 0.81 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 3 | OMC | E | 4 | 3 | - | 0/9/27/28 | 0/2/2/2 |
| 3 | OMC | F | 2 | 3 | - | 0/9/27/28 | 0/2/2/2 |
| 3 | OMC | F | 4 | 3 | - | 0/9/27/28 | 0/2/2/2 |
| 3 | OMC | E | 2 | 3 | - | 0/9/27/28 | 0/2/2/2 |

The worst 5 of 32 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 3 | F | 4 | OMC | C2-N3 | 6.50 | 1.49 | 1.36 |
| 3 | E | 4 | OMC | C2-N3 | 6.39 | 1.49 | 1.36 |
| 3 | F | 2 | OMC | C2-N3 | 6.15 | 1.48 | 1.36 |
| 3 | E | 2 | OMC | C2-N3 | 6.06 | 1.48 | 1.36 |
| 3 | F | 2 | OMC | C6-C5 | 5.96 | 1.48 | 1.35 |

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 3 | E | 4 | OMC | 2 | 0 |
| 3 | F | 2 | OMC | 1 | 0 |
| 3 | F | 4 | OMC | 2 | 0 |
| 3 | E | 2 | OMC | 1 | 0 |

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|-------------|-----------------|-------------|------|-----------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 4 | GOL | E | 101 | - | 5,5,5 | 0.29 | 0 | 5,5,5 | 0.32 | 0 |
| 4 | GOL | D | 501 | - | 5,5,5 | 0.36 | 0 | 5,5,5 | 0.31 | 0 |
| 6 | ET9 | A | 603 | 5 | 33,34,34 | 2.29 | 7 (21%) | 37,54,54 | 1.93 | 9 (24%) |
| 4 | GOL | D | 503 | - | 5,5,5 | 0.54 | 0 | 5,5,5 | 0.31 | 0 |
| 4 | GOL | D | 502 | - | 5,5,5 | 0.38 | 0 | 5,5,5 | 0.27 | 0 |
| 6 | ET9 | C | 602 | 5 | 33,34,34 | 2.39 | 10 (30%) | 37,54,54 | 1.88 | 10 (27%) |
| 4 | GOL | A | 601 | - | 5,5,5 | 0.37 | 0 | 5,5,5 | 0.27 | 0 |
| 4 | GOL | B | 501 | - | 5,5,5 | 0.37 | 0 | 5,5,5 | 0.33 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|------------|---------|
| 4 | GOL | E | 101 | - | - | 2/4/4/4 | - |
| 4 | GOL | D | 501 | - | - | 0/4/4/4 | - |
| 6 | ET9 | A | 603 | 5 | - | 4/22/38/38 | 0/3/3/3 |
| 4 | GOL | D | 503 | - | - | 2/4/4/4 | - |
| 4 | GOL | D | 502 | - | - | 1/4/4/4 | - |
| 6 | ET9 | C | 602 | 5 | - | 4/22/38/38 | 0/3/3/3 |
| 4 | GOL | A | 601 | - | - | 0/4/4/4 | - |
| 4 | GOL | B | 501 | - | - | 1/4/4/4 | - |

The worst 5 of 17 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|-------|-------------|----------|
| 6 | C | 602 | ET9 | C4-N9 | -6.99 | 1.30 | 1.37 |
| 6 | A | 603 | ET9 | C4-N9 | -6.72 | 1.30 | 1.37 |
| 6 | C | 602 | ET9 | C5-C4 | 5.47 | 1.47 | 1.38 |
| 6 | A | 603 | ET9 | C5-C4 | 5.01 | 1.46 | 1.38 |
| 6 | A | 603 | ET9 | PB-O3A | 4.91 | 1.64 | 1.59 |

The worst 5 of 19 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 6 | A | 603 | ET9 | C2'-C3'-C4' | -5.74 | 95.89 | 103.62 |
| 6 | C | 602 | ET9 | C8-N9-C4 | 5.18 | 111.04 | 106.67 |
| 6 | A | 603 | ET9 | C8-N9-C4 | 4.14 | 110.16 | 106.67 |
| 6 | C | 602 | ET9 | O1A-PA-O3A | 4.04 | 118.19 | 107.27 |
| 6 | C | 602 | ET9 | C5-C4-N9 | -3.58 | 104.59 | 106.25 |

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|---------------|
| 4 | E | 101 | GOL | O1-C1-C2-O2 |
| 4 | E | 101 | GOL | O1-C1-C2-C3 |
| 4 | D | 503 | GOL | O1-C1-C2-C3 |
| 6 | A | 603 | ET9 | PB-O3B-PG-O1G |
| 6 | A | 603 | ET9 | PB-O3B-PG-O2G |


There are no ring outliers.

3 monomers are involved in 7 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 4 | E | 101 | GOL | 3 | 0 |
| 4 | D | 501 | GOL | 2 | 0 |
| 6 | A | 603 | ET9 | 2 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|-----------------|-----------------------|--------|
| 1 | A | 553/557 (99%) | 0.53 | 61 (11%) 10 12 | 26, 52, 94, 143 | 1 (0%) |
| 1 | C | 553/557 (99%) | 0.31 | 29 (5%) 33 35 | 27, 53, 96, 149 | 1 (0%) |
| 2 | B | 406/444 (91%) | 0.87 | 80 (19%) 3 3 | 31, 65, 129, 161 | 0 |
| 2 | D | 406/444 (91%) | 0.34 | 35 (8%) 16 18 | 25, 50, 91, 156 | 1 (0%) |
| 3 | E | 33/38 (86%) | -0.43 | 0 100 100 | 29, 48, 81, 125 | 0 |
| 3 | F | 36/38 (94%) | -0.16 | 0 100 100 | 35, 59, 101, 140 | 0 |
| All | All | 1987/2078 (95%) | 0.47 | 205 (10%) 12 13 | 25, 54, 110, 161 | 3 (0%) |

The worst 5 of 205 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | A | 135 | ILE | 6.6 |
| 2 | D | 360 | ALA | 6.5 |
| 2 | B | 88 | TRP | 6.5 |
| 2 | B | 360 | ALA | 6.4 |
| 1 | C | 135 | ILE | 6.2 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 3 | OMC | F | 2 | 21/22 | 0.97 | 0.08 | 39,45,50,56 | 0 |
| 3 | OMC | F | 4 | 21/22 | 0.97 | 0.07 | 30,35,40,48 | 0 |
| 3 | OMC | E | 4 | 21/22 | 0.98 | 0.05 | 25,29,35,40 | 0 |
| 3 | OMC | E | 2 | 21/22 | 0.98 | 0.06 | 29,34,40,42 | 0 |

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 4 | GOL | A | 601 | 6/6 | 0.87 | 0.17 | 49,58,60,61 | 0 |
| 4 | GOL | E | 101 | 6/6 | 0.92 | 0.15 | 63,70,71,72 | 0 |
| 4 | GOL | D | 501 | 6/6 | 0.93 | 0.13 | 46,49,49,51 | 0 |
| 4 | GOL | D | 502 | 6/6 | 0.93 | 0.14 | 41,46,47,48 | 0 |
| 4 | GOL | D | 503 | 6/6 | 0.93 | 0.17 | 55,59,61,61 | 0 |
| 6 | ET9 | C | 602 | 32/32 | 0.94 | 0.08 | 51,60,79,83 | 0 |
| 6 | ET9 | A | 603 | 32/32 | 0.95 | 0.08 | 48,57,68,68 | 0 |
| 4 | GOL | B | 501 | 6/6 | 0.95 | 0.11 | 41,47,49,50 | 0 |
| 5 | MG | C | 601 | 1/1 | 0.96 | 0.10 | 65,65,65,65 | 0 |
| 5 | MG | A | 602 | 1/1 | 0.98 | 0.08 | 55,55,55,55 | 0 |




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ET9 C 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ET9 A 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.