



Full wwPDB EM Validation Report ⓘ

Mar 25, 2026 – 02:49 PM UTC

PDB ID : 7KDP / pdb_00007kdp
EMDB ID : EMD-22828
Title : HCMV prefusion gB in complex with fusion inhibitor WAY-174865
Authors : Liu, Y.; Heim, P.K.; Che, Y.; Chi, X.; Qiu, X.; Han, S.; Dormitzer, P.R.;
Yang, X.
Deposited on : 2020-10-09
Resolution : 3.60 Å(reported)
Based on initial model : 5CXF

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : **FAILED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Q-score	25397	12797 (3.10 - 4.10)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 15844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	638	Total	C	N	O	S	0	0
			5107	3254	858	969	26		
1	B	638	Total	C	N	O	S	0	0
			5107	3254	858	969	26		
1	C	638	Total	C	N	O	S	0	0
			5107	3254	858	969	26		

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	A	1	Total	C	N	O	0
			14	8	1	5	
2	A	1	Total	C	N	O	0
			14	8	1	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total 14	C 8	N 1	O 5	0
2	A	1	Total 14	C 8	N 1	O 5	0
2	A	1	Total 14	C 8	N 1	O 5	0
2	A	1	Total 14	C 8	N 1	O 5	0
2	A	1	Total 14	C 8	N 1	O 5	0
2	A	1	Total 14	C 8	N 1	O 5	0
2	A	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	B	1	Total 14	C 8	N 1	O 5	0
2	C	1	Total 14	C 8	N 1	O 5	0
2	C	1	Total 14	C 8	N 1	O 5	0
2	C	1	Total 14	C 8	N 1	O 5	0
2	C	1	Total 14	C 8	N 1	O 5	0

Continued on next page...

[illegible]

- # WCY

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
4	A	1	Total	Ca	0
			1	1	

MolProbity failed to run properly - this section is therefore empty.

3 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C3	Depositor
Number of particles used	129837	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	18000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.319	Depositor
Minimum map value	-0.187	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.05	Depositor
Map size (Å)	385.28, 385.28, 385.28	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.376, 1.376, 1.376	Depositor

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 34 ligands modelled in this entry, 1 is monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	WCY	C	1101	-	36,36,36	0.11	0	51,53,53	0.44	0
2	NAG	A	1009	1	14,14,15	0.17	0	17,19,21	0.48	0
2	NAG	B	1104	1	14,14,15	0.44	0	17,19,21	0.54	0
2	NAG	B	1106	1	14,14,15	0.28	0	17,19,21	0.49	0
2	NAG	A	1004	1	14,14,15	0.22	0	17,19,21	0.43	0
2	NAG	C	1105	1	14,14,15	0.22	0	17,19,21	0.42	0
2	NAG	C	1108	1	14,14,15	0.29	0	17,19,21	0.53	0
2	NAG	C	1107	1	14,14,15	0.19	0	17,19,21	0.40	0
2	NAG	B	1102	1	14,14,15	0.33	0	17,19,21	0.51	0
2	NAG	A	1006	1	14,14,15	0.18	0	17,19,21	0.40	0
2	NAG	C	1111	1	14,14,15	0.21	0	17,19,21	0.47	0
2	NAG	A	1003	1	14,14,15	0.45	0	17,19,21	0.54	0
2	NAG	B	1107	1	14,14,15	0.18	0	17,19,21	0.39	0
3	WCY	B	1101	-	36,36,36	0.12	0	51,53,53	0.44	0
2	NAG	B	1103	1	14,14,15	0.70	1 (7%)	17,19,21	0.62	0
2	NAG	C	1106	1	14,14,15	0.27	0	17,19,21	0.49	0
2	NAG	B	1109	1	14,14,15	0.32	0	17,19,21	0.50	0
2	NAG	A	1005	1	14,14,15	0.28	0	17,19,21	0.49	0
2	NAG	C	1102	1	14,14,15	0.33	0	17,19,21	0.52	0
2	NAG	A	1002	1	14,14,15	0.69	1 (7%)	17,19,21	0.62	0
2	NAG	C	1103	1	14,14,15	0.69	1 (7%)	17,19,21	0.61	0
2	NAG	B	1111	1	14,14,15	0.20	0	17,19,21	0.47	0
3	WCY	A	1011	-	36,36,36	0.12	0	51,53,53	0.44	0
2	NAG	C	1110	1	14,14,15	0.17	0	17,19,21	0.49	0
2	NAG	C	1104	1	14,14,15	0.45	0	17,19,21	0.54	0
2	NAG	A	1007	1	14,14,15	0.30	0	17,19,21	0.52	0
2	NAG	A	1001	1	14,14,15	0.33	0	17,19,21	0.51	0
2	NAG	A	1010	1	14,14,15	0.21	0	17,19,21	0.47	0
2	NAG	B	1105	1	14,14,15	0.23	0	17,19,21	0.43	0
2	NAG	A	1008	1	14,14,15	0.32	0	17,19,21	0.49	0
2	NAG	B	1108	1	14,14,15	0.30	0	17,19,21	0.53	0
2	NAG	B	1110	1	14,14,15	0.17	0	17,19,21	0.48	0
2	NAG	C	1109	1	14,14,15	0.31	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	WCY	C	1101	-	-	2/32/32/32	0/3/3/3
2	NAG	A	1009	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1104	1	-	4/6/23/26	0/1/1/1
2	NAG	B	1106	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1004	1	-	2/6/23/26	0/1/1/1
2	NAG	C	1105	1	-	2/6/23/26	0/1/1/1
2	NAG	C	1108	1	-	2/6/23/26	0/1/1/1
2	NAG	C	1107	1	-	2/6/23/26	0/1/1/1
2	NAG	B	1102	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1006	1	-	2/6/23/26	0/1/1/1
2	NAG	C	1111	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1003	1	-	4/6/23/26	0/1/1/1
2	NAG	B	1107	1	-	2/6/23/26	0/1/1/1
3	WCY	B	1101	-	-	2/32/32/32	0/3/3/3
2	NAG	B	1103	1	-	2/6/23/26	0/1/1/1
2	NAG	C	1106	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1109	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1005	1	-	0/6/23/26	0/1/1/1
2	NAG	C	1102	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1002	1	-	2/6/23/26	0/1/1/1
2	NAG	C	1103	1	-	2/6/23/26	0/1/1/1
2	NAG	B	1111	1	-	0/6/23/26	0/1/1/1
3	WCY	A	1011	-	-	2/32/32/32	0/3/3/3
2	NAG	C	1110	1	-	0/6/23/26	0/1/1/1
2	NAG	C	1104	1	-	4/6/23/26	0/1/1/1
2	NAG	A	1007	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1001	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1010	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1105	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1008	1	-	2/6/23/26	0/1/1/1
2	NAG	B	1108	1	-	2/6/23/26	0/1/1/1
2	NAG	B	1110	1	-	0/6/23/26	0/1/1/1
2	NAG	C	1109	1	-	2/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1103	NAG	O5-C1	-2.47	1.39	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1103	NAG	O5-C1	-2.46	1.39	1.43
2	A	1002	NAG	O5-C1	-2.44	1.39	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (54) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1002	NAG	O5-C5-C6-O6
2	B	1103	NAG	O5-C5-C6-O6
2	C	1103	NAG	O5-C5-C6-O6
2	A	1003	NAG	O5-C5-C6-O6
2	B	1104	NAG	O5-C5-C6-O6
2	C	1104	NAG	O5-C5-C6-O6
2	A	1004	NAG	C4-C5-C6-O6
2	B	1105	NAG	C4-C5-C6-O6
2	C	1105	NAG	C4-C5-C6-O6
2	A	1002	NAG	C4-C5-C6-O6
2	B	1103	NAG	C4-C5-C6-O6
2	C	1103	NAG	C4-C5-C6-O6
2	A	1003	NAG	C4-C5-C6-O6
2	B	1104	NAG	C4-C5-C6-O6
2	C	1104	NAG	C4-C5-C6-O6
2	A	1006	NAG	O5-C5-C6-O6
2	B	1107	NAG	O5-C5-C6-O6
2	C	1107	NAG	O5-C5-C6-O6
2	A	1001	NAG	C4-C5-C6-O6
2	B	1102	NAG	C4-C5-C6-O6
2	C	1102	NAG	C4-C5-C6-O6
2	A	1004	NAG	O5-C5-C6-O6
2	B	1105	NAG	O5-C5-C6-O6
2	C	1105	NAG	O5-C5-C6-O6
2	A	1001	NAG	O5-C5-C6-O6
2	B	1102	NAG	O5-C5-C6-O6
2	C	1102	NAG	O5-C5-C6-O6
2	A	1006	NAG	C4-C5-C6-O6
2	B	1107	NAG	C4-C5-C6-O6
2	C	1107	NAG	C4-C5-C6-O6
2	A	1008	NAG	C4-C5-C6-O6
2	B	1109	NAG	C4-C5-C6-O6
2	C	1109	NAG	C4-C5-C6-O6

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	1109	NAG	O5-C5-C6-O6
2	A	1008	NAG	O5-C5-C6-O6
2	B	1109	NAG	O5-C5-C6-O6
2	A	1003	NAG	C3-C2-N2-C7
2	B	1104	NAG	C3-C2-N2-C7
2	C	1104	NAG	C3-C2-N2-C7
2	A	1007	NAG	C4-C5-C6-O6
2	C	1108	NAG	C4-C5-C6-O6
2	B	1108	NAG	C4-C5-C6-O6
3	A	1011	WCY	C1-C2-C3-C4
3	B	1101	WCY	C1-C2-C3-C4
3	C	1101	WCY	C1-C2-C3-C4
3	C	1101	WCY	C1-C2-C3-C8
3	A	1011	WCY	C1-C2-C3-C8
3	B	1101	WCY	C1-C2-C3-C8
2	C	1108	NAG	O5-C5-C6-O6
2	A	1007	NAG	O5-C5-C6-O6
2	A	1003	NAG	C1-C2-N2-C7
2	B	1104	NAG	C1-C2-N2-C7
2	C	1104	NAG	C1-C2-N2-C7
2	B	1108	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-22828. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

5.1 Orthogonal projections [i](#)


5.1.1 Primary map



X



Y




Z

The images above show the map projected in three orthogonal directions.

5.2 Central slices [i](#)

5.2.1 Primary map



X Index: 140



Y Index: 140




Z Index: 140

The images above show central slices of the map in three orthogonal directions.

5.3 Largest variance slices [i](#)


5.3.1 Primary map



X Index: 152



Y Index: 137




Z Index: 158


The images above show the largest variance slices of the map in three orthogonal directions.

5.4 Orthogonal standard-deviation projections (False-color) [i](#)


5.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

5.5 Orthogonal surface views [i](#)

5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


6.2 Volume estimate [i](#)



The volume at the recommended contour level is 138 nm³; this corresponds to an approximate mass of 125 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

6.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹

7 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

7.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.278 $\text{\AA}^{-1}$

7.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	3.59	3.89	3.62
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-22828 and PDB model 7KDP. Per-residue inclusion information can be found in section ?? on page ??.


8.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.05).

8.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7210	0.4670
A	0.7210	0.4690
B	0.7210	0.4660
C	0.7200	0.4660

