



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:27 AM UTC

PDB ID : 7KEV / pdb_00007kev
Title : PCSK9 in complex with a cyclic peptide LDLR disruptor
Authors : Spraggon, G.; Chopra, R.
Deposited on : 2020-10-12
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	124	
2	B	546	
3	C	14	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9 Propeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	1	0
			748	479	136	131	2			

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	471	Total	C	N	O	S	0	0	0
			3497	2154	645	667	31			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	474	ILE	VAL	variant	UNP Q8NBP7
B	670	GLU	GLY	variant	UNP Q8NBP7
B	693	HIS	-	expression tag	UNP Q8NBP7
B	694	HIS	-	expression tag	UNP Q8NBP7
B	695	HIS	-	expression tag	UNP Q8NBP7
B	696	HIS	-	expression tag	UNP Q8NBP7
B	697	HIS	-	expression tag	UNP Q8NBP7
B	698	HIS	-	expression tag	UNP Q8NBP7

- Molecule 3 is a protein called cyclic peptide LDLR disruptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	S	0	0	1
			124	85	18	20	1			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Ca 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total 5	O 5	0	0
5	B	12	Total 12	O 12	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proprotein convertase subtilisin/kexin type 9 Propeptide



- Molecule 2: Proprotein convertase subtilisin/kexin type 9



- Molecule 3: cyclic peptide LDLR disruptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.23Å 70.44Å 149.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.64 – 2.80 40.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	93.9 (40.64-2.80) 85.6 (40.64-2.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.226 , 0.280 0.224 , 0.284	Depositor DCC
R_{free} test set	785 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	46.7	Xtriage
Anisotropy	0.615	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4387	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ALO, ACE, NH2, WCM, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/768	0.58	0/1037
2	B	0.32	0/3554	0.50	0/4820
3	C	1.49	0/68	1.21	0/90
All	All	0.37	0/4390	0.53	0/5947

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	748	0	763	4	0
2	B	3497	0	3418	67	0
3	C	124	0	70	3	0
4	B	1	0	0	0	0
5	A	5	0	0	1	0
5	B	12	0	0	0	0
All	All	4387	0	4251	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:494:LYS:HG3	2:B:517:GLY:HA2	1.64	0.80
3:C:10:ARG:HG3	3:C:11:PRO:HD2	1.65	0.76
2:B:641:THR:HG1	2:B:671:ALA:N	1.87	0.71
2:B:558:VAL:HG21	2:B:606:LEU:HD23	1.76	0.67
2:B:375:CYS:HG	2:B:378:CYS:HG	0.69	0.65
2:B:302:GLN:HB2	2:B:331:PRO:HG2	1.79	0.65
2:B:310:VAL:HG22	2:B:408:LEU:HD13	1.80	0.63
2:B:641:THR:OG1	2:B:671:ALA:N	2.31	0.63
2:B:199:ARG:NH2	2:B:235:SER:O	2.33	0.62
2:B:589:VAL:HG13	2:B:638:LEU:HD21	1.81	0.62
2:B:283:LEU:HB3	2:B:309:VAL:HG13	1.82	0.62
2:B:275:GLN:HE21	2:B:283:LEU:HD22	1.64	0.60
2:B:468:THR:OG1	2:B:471:ALA:HB2	2.04	0.58
2:B:619:GLN:N	2:B:619:GLN:HE21	2.02	0.57
2:B:469:ARG:O	2:B:496:ARG:NH1	2.37	0.57
2:B:490:SER:HB2	2:B:520:VAL:HG12	1.86	0.57
2:B:372:SER:HB2	2:B:380:VAL:HB	1.85	0.56
2:B:242:ALA:HA	2:B:398:MET:HE2	1.86	0.56
2:B:306:ARG:HH12	2:B:479:PRO:HG2	1.70	0.56
2:B:375:CYS:CB	2:B:378:CYS:SG	2.95	0.55
2:B:391:HIS:O	2:B:395:ILE:HG13	2.08	0.54
2:B:186:ASP:N	2:B:186:ASP:OD1	2.40	0.54
2:B:645:LEU:HD23	2:B:657:ARG:O	2.09	0.53
2:B:275:GLN:NE2	2:B:283:LEU:HD22	2.24	0.53
2:B:638:LEU:HB2	2:B:673:THR:HB	1.92	0.51
2:B:196:ILE:HB	2:B:200:VAL:HG21	1.91	0.51
2:B:375:CYS:HB2	2:B:378:CYS:SG	2.50	0.51
2:B:465:SER:HB2	2:B:473:ALA:HB2	1.92	0.51
2:B:619:GLN:N	2:B:619:GLN:NE2	2.59	0.51
2:B:614:GLY:HA2	2:B:673:THR:HA	1.93	0.50
2:B:199:ARG:HH22	2:B:237:ARG:HG3	1.77	0.50
2:B:503:GLN:OE1	2:B:510:ARG:NH2	2.44	0.50
2:B:490:SER:O	2:B:645:LEU:HD12	2.12	0.50
2:B:186:ASP:HB2	2:B:253:LEU:HD12	1.94	0.49
2:B:478:ALA:HB3	2:B:481:GLU:HB2	1.94	0.49
2:B:334:ILE:HD11	2:B:408:LEU:HD11	1.94	0.49
2:B:323:CYS:HA	2:B:329:SER:HB3	1.95	0.48
2:B:319:ARG:HG3	2:B:352:GLY:HA3	1.95	0.48
3:C:10:ARG:HG3	3:C:11:PRO:CD	2.38	0.48
2:B:410:GLU:HA	2:B:528:LEU:HD11	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:554:GLN:OE1	2:B:554:GLN:N	2.44	0.48
1:A:82:LEU:HD23	1:A:140:VAL:HA	1.97	0.47
2:B:331:PRO:O	2:B:332:GLU:HB2	2.13	0.47
2:B:208:VAL:HG21	2:B:251:ARG:NH1	2.30	0.47
3:C:3:VAL:HG23	3:C:5:THR:H	1.79	0.47
2:B:272:ARG:NH1	2:B:275:GLN:OE1	2.49	0.46
2:B:613:HIS:NE2	2:B:615:ILE:HD11	2.30	0.46
2:B:628:GLU:OE1	2:B:628:GLU:N	2.46	0.46
2:B:644:VAL:HG23	2:B:657:ARG:O	2.16	0.46
2:B:499:ARG:HD2	2:B:501:GLU:OE2	2.15	0.46
2:B:645:LEU:HD23	2:B:645:LEU:H	1.81	0.44
1:A:151:ALA:HB2	2:B:253:LEU:HD13	2.00	0.44
2:B:658:SER:OG	2:B:672:VAL:HG21	2.17	0.44
2:B:229:HIS:O	2:B:233:VAL:HG23	2.18	0.44
2:B:565:HIS:CE1	2:B:597:HIS:HE2	2.36	0.43
2:B:532:ALA:HA	2:B:602:HIS:O	2.18	0.43
2:B:638:LEU:HA	2:B:638:LEU:HD23	1.79	0.43
2:B:620:GLU:OE1	2:B:621:GLN:HB2	2.18	0.43
2:B:415:LEU:HD23	2:B:415:LEU:HA	1.90	0.43
2:B:465:SER:OG	2:B:471:ALA:HB1	2.19	0.42
2:B:157:ASN:HB2	2:B:391:HIS:NE2	2.34	0.42
2:B:194:ARG:HA	2:B:194:ARG:HD3	1.86	0.42
2:B:343:ASP:O	2:B:424:ILE:HA	2.20	0.42
2:B:589:VAL:HG11	2:B:638:LEU:HD11	2.02	0.42
2:B:607:GLU:O	2:B:679:CYS:HA	2.20	0.41
2:B:455:LEU:HD13	2:B:529:LEU:HD13	2.01	0.41
1:A:108:LEU:HD12	1:A:108:LEU:HA	1.80	0.41
2:B:306:ARG:HG2	2:B:332:GLU:OE2	2.21	0.41
2:B:153:SER:O	2:B:153:SER:OG	2.29	0.41
1:A:132:GLU:HG3	5:A:204:HOH:O	2.21	0.40
2:B:185:LEU:HD23	2:B:252:VAL:HG11	2.04	0.40
2:B:648:TYR:OH	2:B:657:ARG:NH1	2.48	0.40
2:B:230:LEU:HD22	2:B:390:ALA:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/124 (73%)	86 (94%)	4 (4%)	1 (1%)	11	36
2	B	451/546 (83%)	428 (95%)	19 (4%)	4 (1%)	14	41
3	C	8/14 (57%)	6 (75%)	2 (25%)	0	100	100
All	All	550/684 (80%)	520 (94%)	25 (4%)	5 (1%)	14	41

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	GLU
2	B	186	ASP
2	B	585	PRO
2	B	280	VAL
2	B	604	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/104 (77%)	80 (100%)	0	100	100
2	B	379/437 (87%)	378 (100%)	1 (0%)	86	95
3	C	8/8 (100%)	5 (62%)	3 (38%)	0	0
All	All	467/549 (85%)	463 (99%)	4 (1%)	70	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	619	GLN
3	C	3	VAL
3	C	4	SER
3	C	11	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
2	B	278	GLN
2	B	344	GLN
2	B	557	HIS
2	B	621	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	WCM	C	7	3	23,24,25	1.82	8 (34%)	29,32,34	1.47	4 (13%)
3	WCM	C	8	3	23,24,25	1.63	4 (17%)	29,32,34	2.45	14 (48%)
3	ALO	C	6	3	5,6,7	1.56	1 (20%)	5,7,9	3.23	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	WCM	C	7	3	-	0/21/22/24	0/2/2/2
3	WCM	C	8	3	-	2/21/22/24	0/2/2/2
3	ALO	C	6	3	-	2/5/6/8	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	8	WCM	C11-N6	5.16	1.46	1.35
3	C	7	WCM	C11-N6	4.75	1.45	1.35
3	C	7	WCM	C72-C69	2.87	1.55	1.49
3	C	7	WCM	C68-C67	2.79	1.43	1.38
3	C	7	WCM	C71-C70	2.71	1.43	1.38
3	C	7	WCM	C73-C72	2.45	1.44	1.39
3	C	7	WCM	O12-C11	-2.34	1.18	1.22
3	C	8	WCM	C71-C70	2.23	1.42	1.38
3	C	7	WCM	C71-C66	2.19	1.43	1.38
3	C	8	WCM	O11-C14	2.14	1.28	1.20
3	C	8	WCM	O12-C11	-2.10	1.18	1.22
3	C	7	WCM	C77-C72	2.05	1.43	1.39
3	C	6	ALO	CB-CA	2.03	1.59	1.53

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	ALO	CG2-CB-CA	6.39	126.68	112.16
3	C	8	WCM	C67-C68-C69	-4.62	115.19	121.12
3	C	8	WCM	C76-C77-C72	-4.31	115.64	120.54
3	C	7	WCM	C13-C12-N6	4.17	116.29	110.61
3	C	8	WCM	C70-C69-C68	4.07	124.96	117.68
3	C	8	WCM	C13-C12-N6	-3.89	105.31	110.61
3	C	8	WCM	C68-C69-C72	-3.61	115.03	121.25
3	C	8	WCM	C77-C72-C73	3.60	124.12	117.68
3	C	6	ALO	OG1-CB-CA	-3.13	102.41	109.01
3	C	8	WCM	C65-C66-C67	-3.01	115.31	120.90
3	C	8	WCM	C10-C11-N6	-2.93	113.27	119.44
3	C	8	WCM	C73-C72-C69	-2.90	116.25	121.25
3	C	8	WCM	C75-C76-C77	2.73	123.60	120.24
3	C	7	WCM	C71-C66-C67	-2.71	114.21	118.23
3	C	7	WCM	C65-C10-C11	2.66	116.53	109.43
3	C	8	WCM	C71-C70-C69	-2.54	117.87	121.12
3	C	7	WCM	O12-C11-N6	-2.49	117.67	121.76
3	C	8	WCM	C71-C66-C67	2.24	121.56	118.23
3	C	8	WCM	C70-C71-C66	-2.18	118.13	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	8	WCM	C12-N6-C11	2.13	125.92	117.98

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	8	WCM	C10-C65-C66-C71
3	C	8	WCM	C10-C65-C66-C67
3	C	6	ALO	N-CA-CB-CG2
3	C	6	ALO	C-CA-CB-CG2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/124 (74%)	0.01	1 (1%) 78 70	33, 44, 53, 64	1 (1%)
2	B	471/546 (86%)	0.51	29 (6%) 26 20	39, 54, 82, 103	0
3	C	9/14 (64%)	0.59	0 100 100	48, 54, 68, 73	0
All	All	572/684 (83%)	0.43	30 (5%) 33 25	33, 50, 80, 103	1 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	612	GLU	4.2
2	B	467	PRO	3.6
2	B	466	GLY	3.2
2	B	568	VAL	3.0
2	B	204	ASP	3.0
2	B	520	VAL	3.0
2	B	589	VAL	2.7
2	B	537	HIS	2.6
2	B	212	ASP	2.6
2	B	220	ALA	2.6
2	B	332	GLU	2.5
2	B	671	ALA	2.5
2	B	541	PRO	2.5
2	B	591	HIS	2.5
2	B	641	THR	2.4
2	B	507	LEU	2.4
2	B	519	GLY	2.3
2	B	166	TYR	2.3
2	B	602	HIS	2.3
2	B	504	GLY	2.3
2	B	533	ASN	2.3
2	B	538	THR	2.3
2	B	211	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	593	GLU	2.1
2	B	550	VAL	2.1
1	A	108	LEU	2.1
2	B	640	GLY	2.1
2	B	548	THR	2.1
2	B	634	GLY	2.0
2	B	521	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ALO	C	6	7/8	0.87	0.09	45,48,50,51	0
3	WCM	C	7	23/24	0.94	0.09	39,45,51,54	0
3	WCM	C	8	23/24	0.94	0.10	42,51,56,60	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CA	B	801	1/1	0.81	0.13	79,79,79,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.