



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 02:23 PM UTC

PDB ID : 8KEC / pdb_00008kec
EMDB ID : EMD-37152
Title : Cyanophage A-1(L) tail fiber
Authors : Yu, R.C.; Li, Q.; Zhou, C.Z.
Deposited on : 2023-08-11
Resolution : 3.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





















Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8855 (3.40 - 4.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	
1	B	379	
1	C	379	
1	D	379	
1	E	379	







Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	379	
1	G	379	
1	H	379	
1	I	379	
1	J	379	
1	K	379	
1	L	379	
1	M	379	
1	N	379	
1	O	379	
1	P	379	
1	Q	379	
1	R	379	
2	a	462	
2	b	462	
2	c	462	
2	d	462	
2	e	462	
2	f	462	
2	g	462	
2	h	462	
2	i	462	
2	j	462	
2	k	462	
2	l	462	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	m	462	
2	n	462	
2	o	462	
2	p	462	
2	q	462	
2	r	462	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 87756 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Long tail fiber.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	379	Total	C	N	O	S	0	0
			2795	1744	491	558	2		
1	B	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	C	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	D	379	Total	C	N	O	S	0	0
			2795	1744	491	558	2		
1	E	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	F	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	G	379	Total	C	N	O	S	0	0
			2795	1744	491	558	2		
1	H	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	I	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	J	379	Total	C	N	O	S	0	0
			2795	1744	491	558	2		
1	K	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	L	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	M	379	Total	C	N	O	S	0	0
			2795	1744	491	558	2		
1	N	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	O	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		
1	P	379	Total	C	N	O	S	0	0
			2795	1744	491	558	2		
1	Q	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	378	Total	C	N	O	S	0	0
			2787	1739	490	557	1		


- Molecule 2 is a protein called short tail fiber.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	n	280	Total	C	N	O	S	0	0
			2091	1333	331	425	2		
2	i	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	j	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	k	280	Total	C	N	O	S	0	0
			2091	1333	331	425	2		
2	l	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	m	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	o	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	p	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	q	280	Total	C	N	O	S	0	0
			2091	1333	331	425	2		
2	r	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	a	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	b	280	Total	C	N	O	S	0	0
			2091	1333	331	425	2		
2	c	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	d	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	e	280	Total	C	N	O	S	0	0
			2091	1333	331	425	2		
2	f	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	g	279	Total	C	N	O	S	0	0
			2083	1328	330	424	1		
2	h	280	Total	C	N	O	S	0	0
			2091	1333	331	425	2		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Long tail fiber




- Molecule 1: Long tail fiber




- Molecule 1: Long tail fiber






• Molecule 1: Long tail fiber




• Molecule 1: Long tail fiber




• Molecule 1: Long tail fiber






- Molecule 1: Long tail fiber

Chain G: 73% 26% .



- Molecule 1: Long tail fiber


Chain H: 72% 26% .



- Molecule 1: Long tail fiber


Chain I: 71% 27% .






- Molecule 1: Long tail fiber

Chain J: 73% 25% .



- Molecule 1: Long tail fiber


Chain K: 72% 26% .



- Molecule 1: Long tail fiber


Chain L: 72% 27% .






- Molecule 1: Long tail fiber

Chain M: 71% 29%




- Molecule 1: Long tail fiber

Chain N: 71% 28%




- Molecule 1: Long tail fiber

Chain O: 74% 24%




- Molecule 1: Long tail fiber

Chain P: 72% 27%




- Molecule 1: Long tail fiber

Chain Q: 72% 26%





- Molecule 1: Long tail fiber


Chain R: 69% 29%



• Molecule 2: short tail fiber




• Molecule 2: short tail fiber



VAL PHE LEU ASN TYR VAL VAL VAL GLY GLY SER TYR VAL VAL LEU GLU ASP VAL ARG PHE SER GLY ILE GLN LEU GLY


• Molecule 2: short tail fiber

Chain l: 47% 12% 40%




• Molecule 2: short tail fiber

Chain m: 42% 16% 40%




• Molecule 2: short tail fiber



Chain o: 47% 12% 40%




• Molecule 2: short tail fiber




• Molecule 2: short tail fiber






• Molecule 2: short tail fiber




• Molecule 2: short tail fiber





• Molecule 2: short tail fiber



- Molecule 2: short tail fiber



- Molecule 2: short tail fiber



PHE
ALA
THR
VAL
GLY
GLY
THR
PRO
THR
SER
TYR
PHE
ASP
ALA
LEU
PHE
VAL
GLY
LEU
SER
GLY
GLY
GLY
VAL
GLU
ASN
PHE
LEU
THR
ARG
SER
GLN
SER
VAL
TYR
PRO
SER
PHE
GLY
TYR
THR
LEU
SER
TRP
VAL
GLY
ILE
LEU
THR
PHE
ASN
THR
GLY
GLN
ASN
VAL
PHE
LEU
ASN

TYR
GLN
VAL
ASN
ALA
VAL
GLY
GLY
GLY
SER
TYR
SER
VAL
VAL
LEU
GLU
ASP
VAL
ARG
PHE
SER
GLY
ILE
GLN
LEU
GLY

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	41062	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.032	Depositor
Minimum map value	-0.015	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.006	Depositor
Map size ($\text{\AA}$)	479.36002, 479.36002, 479.36002	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.07, 1.07, 1.07	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/2859	0.30	0/3916
1	B	0.21	0/2851	0.36	0/3906
1	C	0.16	0/2851	0.32	0/3906
1	D	0.17	0/2859	0.30	0/3916
1	E	0.19	0/2851	0.37	0/3906
1	F	0.17	0/2851	0.32	0/3906
1	G	0.17	0/2859	0.30	0/3916
1	H	0.17	0/2851	0.32	0/3906
1	I	0.17	0/2851	0.32	0/3906
1	J	0.18	0/2859	0.32	0/3916
1	K	0.19	0/2851	0.34	0/3906
1	L	0.17	0/2851	0.32	0/3906
1	M	0.17	0/2859	0.31	0/3916
1	N	0.23	0/2851	0.39	0/3906
1	O	0.17	0/2851	0.33	0/3906
1	P	0.17	0/2859	0.30	0/3916
1	Q	0.17	0/2851	0.33	0/3906
1	R	0.17	0/2851	0.33	0/3906
2	a	0.20	0/2122	0.50	4/2914 (0.1%)
2	b	0.22	0/2130	0.57	6/2924 (0.2%)
2	c	0.19	0/2122	0.43	1/2914 (0.0%)
2	d	0.21	0/2122	0.51	3/2914 (0.1%)
2	e	0.22	0/2130	0.55	5/2924 (0.2%)
2	f	0.19	0/2122	0.43	1/2914 (0.0%)
2	g	0.22	0/2122	0.55	7/2914 (0.2%)
2	h	0.23	0/2130	0.55	5/2924 (0.2%)
2	i	0.21	0/2122	0.46	3/2914 (0.1%)
2	j	0.22	0/2122	0.52	1/2914 (0.0%)
2	k	0.22	0/2130	0.55	1/2924 (0.0%)
2	l	0.20	0/2122	0.44	1/2914 (0.0%)
2	m	0.30	1/2122 (0.0%)	0.59	5/2914 (0.2%)
2	n	0.26	0/2130	0.63	3/2924 (0.1%)
2	o	0.22	0/2122	0.46	1/2914 (0.0%)
2	p	0.26	1/2122 (0.0%)	0.54	4/2914 (0.1%)

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
2	q	0.21	0/2130	0.51	3/2924 (0.1%)
2	r	0.21	0/2122	0.47	4/2914 (0.1%)
All	All	0.20	2/89610 (0.0%)	0.42	58/122880 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	p	81	PRO	N-CD	6.02	1.56	1.47
2	m	187	LEU	C-N	5.48	1.38	1.33

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	a	79	VAL	N-CA-C	-9.85	99.32	110.05
2	e	79	VAL	N-CA-C	-9.82	99.35	110.05
2	g	79	VAL	N-CA-C	-9.36	99.84	110.05
2	b	79	VAL	N-CA-C	-9.26	99.95	110.05
2	h	79	VAL	N-CA-C	-8.43	100.86	110.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2795	0	2708	92	0
1	B	2787	0	2696	81	0
1	C	2787	0	2696	84	0
1	D	2795	0	2708	101	0
1	E	2787	0	2696	93	0
1	F	2787	0	2696	83	0
1	G	2795	0	2708	94	0
1	H	2787	0	2696	87	0
1	I	2787	0	2696	85	0
1	J	2795	0	2708	100	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	K	2787	0	2696	87	0
1	L	2787	0	2696	86	0
1	M	2795	0	2708	108	0
1	N	2787	0	2696	93	0
1	O	2787	0	2696	81	0
1	P	2795	0	2708	102	0
1	Q	2787	0	2696	89	0
1	R	2787	0	2696	96	0
2	a	2083	0	2082	64	0
2	b	2091	0	2094	69	0
2	c	2083	0	2082	62	0
2	d	2083	0	2082	66	0
2	e	2091	0	2094	66	0
2	f	2083	0	2082	48	0
2	g	2083	0	2082	64	0
2	h	2091	0	2094	53	0
2	i	2083	0	2082	54	0
2	j	2083	0	2082	66	0
2	k	2091	0	2094	73	0
2	l	2083	0	2082	61	0
2	m	2083	0	2082	78	0
2	n	2091	0	2094	91	0
2	o	2083	0	2082	54	0
2	p	2083	0	2082	53	0
2	q	2091	0	2094	54	0
2	r	2083	0	2082	49	0
All	All	87756	0	86148	1984	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 1984 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:N:45:GLU:HB2	1:N:48:ASN:OD1	1.34	1.26
1:J:39:ASP:OD1	1:J:50:LYS:HD2	1.43	1.18
2:n:223:VAL:HG11	2:m:221:LYS:NZ	1.58	1.16
2:k:198:SER:HB3	2:k:212:VAL:CG2	1.76	1.15
1:P:39:ASP:OD1	1:P:50:LYS:HD2	1.53	1.08

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/311 (100%)	302 (97%)	9 (3%)	37	58
1	B	310/311 (100%)	295 (95%)	15 (5%)	23	48
1	C	310/311 (100%)	296 (96%)	14 (4%)	24	49
1	D	311/311 (100%)	304 (98%)	7 (2%)	44	63
1	E	310/311 (100%)	299 (96%)	11 (4%)	32	54
1	F	310/311 (100%)	297 (96%)	13 (4%)	26	50
1	G	311/311 (100%)	304 (98%)	7 (2%)	44	63
1	H	310/311 (100%)	297 (96%)	13 (4%)	26	50
1	I	310/311 (100%)	297 (96%)	13 (4%)	26	50
1	J	311/311 (100%)	302 (97%)	9 (3%)	37	58
1	K	310/311 (100%)	296 (96%)	14 (4%)	24	49
1	L	310/311 (100%)	296 (96%)	14 (4%)	24	49
1	M	311/311 (100%)	302 (97%)	9 (3%)	37	58
1	N	310/311 (100%)	300 (97%)	10 (3%)	34	56
1	O	310/311 (100%)	299 (96%)	11 (4%)	32	54
1	P	311/311 (100%)	303 (97%)	8 (3%)	40	60
1	Q	310/311 (100%)	296 (96%)	14 (4%)	24	49
1	R	310/311 (100%)	300 (97%)	10 (3%)	34	56
2	a	234/384 (61%)	224 (96%)	10 (4%)	26	49
2	b	235/384 (61%)	221 (94%)	14 (6%)	17	44
2	c	234/384 (61%)	225 (96%)	9 (4%)	29	52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	d	234/384 (61%)	222 (95%)	12 (5%)	21	46
2	e	235/384 (61%)	228 (97%)	7 (3%)	36	57
2	f	234/384 (61%)	221 (94%)	13 (6%)	19	45
2	g	234/384 (61%)	226 (97%)	8 (3%)	32	55
2	h	235/384 (61%)	227 (97%)	8 (3%)	32	55
2	i	234/384 (61%)	224 (96%)	10 (4%)	26	49
2	j	234/384 (61%)	219 (94%)	15 (6%)	16	42
2	k	235/384 (61%)	216 (92%)	19 (8%)	11	35
2	l	234/384 (61%)	221 (94%)	13 (6%)	19	45
2	m	234/384 (61%)	221 (94%)	13 (6%)	19	45
2	n	235/384 (61%)	215 (92%)	20 (8%)	10	33
2	o	234/384 (61%)	217 (93%)	17 (7%)	13	39
2	p	234/384 (61%)	224 (96%)	10 (4%)	26	49
2	q	235/384 (61%)	220 (94%)	15 (6%)	16	42
2	r	234/384 (61%)	224 (96%)	10 (4%)	26	49
All	All	9804/12510 (78%)	9380 (96%)	424 (4%)	27	49

5 of 424 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	j	30	THR
2	m	193	VAL
2	f	64	THR
2	j	110	GLN
2	k	212	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 152 such sidechains are listed below:

Mol	Chain	Res	Type
2	k	109	ASN
2	e	109	ASN
2	m	160	ASN
2	r	238	ASN
2	g	160	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-37152. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 224




Y Index: 224




Z Index: 224


6.2.2 Raw map



X Index: 224



Y Index: 224




Z Index: 224


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 204




Y Index: 204




Z Index: 221


6.3.2 Raw map



X Index: 204



Y Index: 244




Z Index: 221


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.006. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 2387 nm³; this corresponds to an approximate mass of 2157 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.90	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.19	3.77	3.25

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.19 differs from the reported value 3.9 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-37152 and PDB model 8KEC. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.006 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.006).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.006) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8160	 0.5100
A	 0.8520	 0.5260
B	 0.8570	 0.5360
C	 0.8530	 0.5380
D	 0.8460	 0.5260
E	 0.8530	 0.5350
F	 0.8480	 0.5370
G	 0.8540	 0.5260
H	 0.8540	 0.5350
I	 0.8500	 0.5380
J	 0.8490	 0.5260
K	 0.8580	 0.5370
L	 0.8540	 0.5390
M	 0.8460	 0.5250
N	 0.8530	 0.5350
O	 0.8490	 0.5390
P	 0.8540	 0.5260
Q	 0.8550	 0.5370
R	 0.8500	 0.5380
a	 0.7620	 0.4730
b	 0.7540	 0.4660
c	 0.7950	 0.5070
d	 0.7670	 0.4710
e	 0.7510	 0.4660
f	 0.7930	 0.5060
g	 0.7650	 0.4740
h	 0.7450	 0.4620
i	 0.7940	 0.5030
j	 0.7680	 0.4710
k	 0.7470	 0.4630
l	 0.7900	 0.5030
m	 0.7630	 0.4720
n	 0.7240	 0.4420
o	 0.7870	 0.5000
p	 0.7620	 0.4720



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
q	 0.7490	 0.4660
r	 0.7930	 0.5060