



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 02:49 AM UTC

PDB ID : 5KF7 / pdb_00005kf7
Title : Structure of proline utilization A from Sinorhizobium meliloti complexed with L-tetrahydrofuroic acid and NAD⁺ in space group P3121
Authors : Tanner, J.J.
Deposited on : 2016-06-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1235	

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 9756 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional protein PutA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1211	9004	5663	1614	1694	33	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	expression tag	UNP F7X6I3
A	0	MET	-	expression tag	UNP F7X6I3

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is TETRAHYDROFURAN-2-CARBOXYLIC ACID (CCD ID: TFB) (formula: C₅H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	5	3		

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 7 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total 1	Mg 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	604	Total 604	O 604	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional protein PutA



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.84Å 128.84Å 150.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	62.39 – 1.90 62.39 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (62.39-1.90) 99.8 (62.39-1.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 1.90Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, R_{free}	0.174 , 0.203 0.172 , 0.201	Depositor DCC
R_{free} test set	5640 reflections (2.97%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.414	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9756	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, PGE, PG4, FAD, 1PE, MG, TFB, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/9170	0.46	0/12479

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9004	0	9032	38	1
2	A	53	0	31	3	0
3	A	44	0	26	2	0
4	A	8	0	7	0	0
5	A	10	0	14	2	0
6	A	13	0	18	0	0
7	A	14	0	16	2	0
8	A	5	0	0	0	0
9	A	1	0	0	0	0
10	A	604	0	0	4	0
All	All	9756	0	9144	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1079:GLY:HA2	5:A:2004:PGE:H2	1.66	0.76
1:A:844:CYS:SG	3:A:2002:NAD:C4N	2.81	0.69
1:A:873:ILE:HG13	1:A:883:VAL:HB	1.78	0.65
1:A:288[B]:ARG:NE	10:A:2101:HOH:O	2.33	0.60
1:A:56:GLN:O	1:A:61:ARG:NH2	2.36	0.58
1:A:85:GLU:O	1:A:89:GLN:HG2	2.02	0.58
1:A:144:ALA:HB3	1:A:159:VAL:HG11	1.90	0.54
1:A:759:LEU:HD13	1:A:769:LEU:HD21	1.91	0.52
1:A:650:MET:O	1:A:654:LEU:HG	2.11	0.50
1:A:793:GLN:HG2	10:A:2190:HOH:O	2.13	0.49
1:A:155:LEU:HD22	1:A:491:LEU:HD21	1.96	0.47
1:A:1079:GLY:HA2	5:A:2004:PGE:C2	2.39	0.46
1:A:421:ALA:HB1	2:A:2001:FAD:C8	2.46	0.46
1:A:460:VAL:HG22	1:A:466:LEU:HD12	1.97	0.46
7:A:2007:1PE:H242	7:A:2007:1PE:H232	1.55	0.45
1:A:266:LEU:HD13	1:A:319:LEU:HD13	1.99	0.45
1:A:272:ARG:HB3	1:A:277:GLN:HG3	1.99	0.44
1:A:918:THR:HB	1:A:923:PHE:CD1	2.53	0.44
1:A:1226:ASN:HB3	1:A:1229:LEU:HB2	2.00	0.44
1:A:448:CYS:HB2	1:A:453:GLY:HA3	2.00	0.43
1:A:58:LYS:HD2	1:A:61:ARG:NH1	2.33	0.43
1:A:197:THR:O	1:A:207:ARG:HD2	2.18	0.43
1:A:130:TRP:H	1:A:130:TRP:CD1	2.37	0.43
1:A:187:ALA:O	1:A:191:MET:HG3	2.19	0.43
1:A:1056:TRP:CD1	1:A:1142:PRO:HD3	2.54	0.43
1:A:931:LEU:HB2	1:A:949:ILE:HG22	2.00	0.42
1:A:449:LEU:HD23	1:A:473:TYR:HB3	2.00	0.42
1:A:498:SER:O	1:A:502:ARG:HG3	2.19	0.42
1:A:782:GLY:O	1:A:811:THR:HA	2.19	0.42
1:A:984:HIS:C	1:A:984:HIS:CD2	2.97	0.42
2:A:2001:FAD:H4'	2:A:2001:FAD:H1'1	1.83	0.42
1:A:1026:LEU:HD23	1:A:1038:PRO:HG2	2.01	0.42
2:A:2001:FAD:H1'1	2:A:2001:FAD:H9	1.86	0.42
1:A:706:TRP:CE3	1:A:707:ASN:HA	2.54	0.42
1:A:785:GLU:OE1	1:A:785:GLU:N	2.48	0.42
1:A:367:ARG:HA	1:A:419:GLN:HB2	2.02	0.41
1:A:601:VAL:HG12	7:A:2006:1PE:H231	2.01	0.41
3:A:2002:NAD:H8A	10:A:2623:HOH:O	2.18	0.41
1:A:186:MET:O	1:A:190:MET:HG3	2.20	0.40
1:A:338:VAL:HG22	1:A:367:ARG:HB3	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:369:VAL:HG12	1:A:421:ALA:HB3	2.02	0.40
1:A:892:LYS:NZ	10:A:2120:HOH:O	2.50	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:THR:N	1:A:748:GLU:OE1[6_655]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1208/1235 (98%)	1188 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	908/951 (96%)	899 (99%)	9 (1%)	68	70

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	SER
1	A	194	GLN
1	A	258	GLU
1	A	342	TYR
1	A	483	LEU
1	A	788	ARG
1	A	855	ASP
1	A	934	LEU
1	A	1147	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	419	GLN
1	A	450	HIS
1	A	685	GLN
1	A	872	HIS
1	A	1140	ASN
1	A	1226	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	1PE	A	2006	-	6,6,15	0.64	0	5,5,14	0.76	0
5	PGE	A	2004	-	9,9,9	0.64	0	8,8,8	0.90	0
3	NAD	A	2002	9	46,48,48	2.21	8 (17%)	64,73,73	1.73	10 (15%)
7	1PE	A	2007	-	6,6,15	0.46	0	5,5,14	0.87	0
2	FAD	A	2001	-	58,58,58	2.33	17 (29%)	85,89,89	1.76	18 (21%)
6	PG4	A	2005	-	12,12,12	0.65	0	11,11,11	0.70	0
8	SO4	A	2008	-	4,4,4	0.17	0	6,6,6	0.23	0
4	TFB	A	2003	-	8,8,8	1.48	2 (25%)	7,10,10	1.70	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	1PE	A	2006	-	-	2/4/4/13	-
5	PGE	A	2004	-	-	3/7/7/7	-
3	NAD	A	2002	9	-	4/30/62/62	0/5/5/5
7	1PE	A	2007	-	-	2/4/4/13	-
2	FAD	A	2001	-	-	4/34/50/50	0/6/6/6
6	PG4	A	2005	-	-	2/10/10/10	-
4	TFB	A	2003	-	-	3/4/11/11	0/1/1/1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	PA-O3P	-8.49	1.50	1.59
3	A	2002	NAD	PA-O3	-7.99	1.50	1.59
2	A	2001	FAD	O4-C4	7.78	1.38	1.23
3	A	2002	NAD	C2N-N1N	6.94	1.42	1.35
3	A	2002	NAD	C7N-N7N	5.66	1.43	1.33
2	A	2001	FAD	O2-C2	5.40	1.35	1.24
2	A	2001	FAD	C4X-N5	4.33	1.40	1.30
2	A	2001	FAD	C6A-N6A	4.28	1.45	1.34
3	A	2002	NAD	C6A-N6A	3.77	1.43	1.34
2	A	2001	FAD	P-O3P	3.48	1.63	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	C2-N1	3.31	1.44	1.36
3	A	2002	NAD	C6N-N1N	2.83	1.41	1.35
4	A	2003	TFB	CB-CA	-2.81	1.46	1.52
2	A	2001	FAD	C10-N1	2.66	1.38	1.33
2	A	2001	FAD	O2'-C2'	-2.52	1.38	1.43
3	A	2002	NAD	PA-O5B	-2.52	1.49	1.59
3	A	2002	NAD	C5N-C4N	2.27	1.42	1.38
4	A	2003	TFB	OXT-C	2.26	1.28	1.22
2	A	2001	FAD	C5A-C4A	2.26	1.43	1.39
2	A	2001	FAD	O4'-C4'	-2.25	1.38	1.43
2	A	2001	FAD	PA-O2A	-2.22	1.45	1.55
2	A	2001	FAD	P-O1P	2.19	1.58	1.50
3	A	2002	NAD	PA-O2A	-2.15	1.45	1.55
2	A	2001	FAD	PA-O5B	-2.13	1.51	1.59
2	A	2001	FAD	O4B-C4B	-2.12	1.40	1.45
2	A	2001	FAD	O3'-C3'	-2.12	1.37	1.43
2	A	2001	FAD	C4A-N9A	2.06	1.42	1.37

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2002	NAD	N3A-C2A-N1A	-5.62	120.07	128.58
2	A	2001	FAD	N3A-C2A-N1A	-5.45	120.33	128.58
2	A	2001	FAD	C5A-C4A-N3A	-5.18	119.58	126.72
3	A	2002	NAD	C5A-C4A-N3A	-4.58	120.41	126.72
2	A	2001	FAD	O2P-P-O3P	-4.44	95.27	107.27
3	A	2002	NAD	C5A-N7A-C8A	4.25	110.12	103.45
2	A	2001	FAD	N3A-C4A-N9A	4.13	134.18	127.17
2	A	2001	FAD	C5A-N7A-C8A	3.86	109.51	103.45
3	A	2002	NAD	N3A-C4A-N9A	3.73	133.51	127.17
2	A	2001	FAD	C4-C4X-N5	3.73	123.36	118.21
3	A	2002	NAD	N9A-C8A-N7A	-3.69	108.69	113.94
2	A	2001	FAD	C2A-N3A-C4A	3.61	120.66	111.83
3	A	2002	NAD	C2A-N3A-C4A	3.44	120.23	111.83
3	A	2002	NAD	C4A-C5A-N7A	-3.33	106.77	110.58
2	A	2001	FAD	O3P-P-O1P	3.04	119.85	110.70
2	A	2001	FAD	C4A-C5A-N7A	-3.02	107.13	110.58
4	A	2003	TFB	OB-C-CA	2.99	120.49	112.71
2	A	2001	FAD	C4-N3-C2	-2.85	120.57	125.64
3	A	2002	NAD	O2N-PN-O3	2.82	114.91	107.27
2	A	2001	FAD	N9A-C8A-N7A	-2.81	109.94	113.94
2	A	2001	FAD	C10-C4X-N5	-2.70	119.29	124.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FAD	C4X-C10-N10	2.66	120.29	116.48
2	A	2001	FAD	O2-C2-N1	-2.66	117.39	121.80
2	A	2001	FAD	C4X-C4-N3	2.51	119.64	113.25
3	A	2002	NAD	C4A-N9A-C8A	2.47	108.33	105.74
2	A	2001	FAD	O5'-P-O1P	2.40	118.45	108.94
2	A	2001	FAD	O2P-P-O5'	-2.17	97.75	107.57
4	A	2003	TFB	CB-CA-C	-2.14	109.56	113.00
2	A	2001	FAD	O4-C4-C4X	-2.12	120.94	126.53
3	A	2002	NAD	O2A-PA-O5B	-2.00	98.49	107.57

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2001	FAD	P-O3P-PA-O5B
2	A	2001	FAD	N10-C1'-C2'-O2'
2	A	2001	FAD	N10-C1'-C2'-C3'
2	A	2001	FAD	C1'-C2'-C3'-C4'
7	A	2007	1PE	C23-C13-OH4-C24
7	A	2007	1PE	OH4-C13-C23-OH3
5	A	2004	PGE	O2-C3-C4-O3
5	A	2004	PGE	O1-C1-C2-O2
4	A	2003	TFB	OXT-C-CA-OA
3	A	2002	NAD	PN-O3-PA-O5B
6	A	2005	PG4	C4-C3-O2-C2
5	A	2004	PGE	C4-C3-O2-C2
3	A	2002	NAD	C2B-C1B-N9A-C8A
3	A	2002	NAD	C4D-C5D-O5D-PN
4	A	2003	TFB	OB-C-CA-CB
4	A	2003	TFB	OXT-C-CA-CB
6	A	2005	PG4	C3-C4-O3-C5
3	A	2002	NAD	O4B-C1B-N9A-C8A
7	A	2006	1PE	OH4-C13-C23-OH3
7	A	2006	1PE	C23-C13-OH4-C24

There are no ring outliers.

5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	2006	1PE	1	0
5	A	2004	PGE	2	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2002	NAD	2	0
7	A	2007	1PE	1	0
2	A	2001	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1211/1235 (98%)	-0.04	53 (4%)	39 41	10, 29, 58, 94	3 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	ASP	5.7
1	A	129	ASN	5.7
1	A	1227	ALA	4.7
1	A	1230	MET	4.6
1	A	1231	ALA	4.5
1	A	1229	LEU	4.5
1	A	14	ALA	4.3
1	A	128	GLY	3.9
1	A	481	THR	3.9
1	A	1228	SER	3.5
1	A	1223	ALA	3.5
1	A	483	LEU	3.4
1	A	482	LEU	3.4
1	A	134	LEU	3.2
1	A	1225	GLY	3.2
1	A	464	GLY	3.1
1	A	912	ILE	2.9
1	A	499	PHE	2.8
1	A	1222	ALA	2.8
1	A	1147	LEU	2.8
1	A	83	GLY	2.8
1	A	480	GLU	2.7
1	A	508	VAL	2.7
1	A	493	ASN	2.7
1	A	905	LEU	2.7
1	A	78	LYS	2.7
1	A	800	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	75	LEU	2.6
1	A	486	LEU	2.6
1	A	1064	ALA	2.6
1	A	195	PHE	2.5
1	A	215	GLY	2.5
1	A	506	PRO	2.5
1	A	1224	GLY	2.5
1	A	1226	ASN	2.4
1	A	510	ILE	2.4
1	A	485	TYR	2.4
1	A	463	ARG	2.4
1	A	904	GLY	2.4
1	A	513	LEU	2.4
1	A	77	GLY	2.4
1	A	155	LEU	2.3
1	A	490	LEU	2.2
1	A	133	HIS	2.2
1	A	138	ARG	2.2
1	A	479	HIS	2.1
1	A	949	ILE	2.1
1	A	156	THR	2.1
1	A	466	LEU	2.1
1	A	72	ILE	2.0
1	A	507	LYS	2.0
1	A	223	LEU	2.0
1	A	15	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no oligosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	1PE	A	2006	7/16	0.89	0.13	48,50,54,59	0
5	PGE	A	2004	10/10	0.91	0.12	36,41,51,53	0
7	1PE	A	2007	7/16	0.91	0.15	20,37,46,50	0
4	TFB	A	2003	8/8	0.93	0.10	32,37,42,43	0
6	PG4	A	2005	13/13	0.93	0.11	22,30,53,54	0
2	FAD	A	2001	53/53	0.94	0.08	19,31,42,45	0
9	MG	A	2009	1/1	0.94	0.15	37,37,37,37	0
3	NAD	A	2002	44/44	0.97	0.05	15,19,25,29	0
8	SO4	A	2008	5/5	0.99	0.04	21,21,22,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around NAD A 2002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.