



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 03:48 AM EDT

PDB ID : 7KFL / pdb_00007kfl
Title : Crystal structure of the cargo-binding domain from the plant class XI myosin (MyoXIk)
Authors : Turowski, V.R.; Ruiz, D.M.; Nascimento, A.F.Z.; Millan, C.; Sammito, M.D.; Juanhuix, J.; Cremonesi, A.S.; Uson, I.; Giuseppe, P.O.; Murakami, M.T.
Deposited on : 2020-10-14
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	3% 85% 5% 10%
1	B	395	6% 78% 7% 15%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5619 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	0	0
			2850	1812	491	532	15			
1	B	336	Total	C	N	O	S	0	0	0
			2707	1728	462	502	15			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	37	Total	O	0	0
			37	37		
2	B	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Myosin-17



• Molecule 1: Myosin-17



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	147.06Å 54.33Å 113.43Å 90.00° 96.60° 90.00°	Depositor
Resolution (Å)	47.36 – 2.35 47.36 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.36-2.35) 99.3 (47.36-2.35)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.232 , 0.273 0.235 , 0.275	Depositor DCC
R_{free} test set	1875 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	56.2	Xtriage
Anisotropy	0.428	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5619	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/2900	1.55	0/3923
1	B	0.99	0/2755	1.54	0/3727
All	All	0.99	0/5655	1.54	0/7650

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2850	0	2877	11	0
1	B	2707	0	2733	9	0
2	A	37	0	0	0	0
2	B	25	0	0	0	0
All	All	5619	0	5610	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1121:LYS:HE2	1:A:1121:LYS:HA	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1418:LYS:N	1:A:1419:PRO:CD	2.76	0.48
1:A:1296:SER:O	1:A:1297:ARG:C	2.57	0.46
1:A:1283:SER:HB3	1:A:1284:PRO:HD3	1.98	0.46
1:A:1377:VAL:HG12	1:A:1410:VAL:HG21	1.98	0.46
1:A:1118:LEU:HB3	1:A:1233:LEU:CD1	2.45	0.45
1:A:1190:LEU:HD22	1:A:1190:LEU:O	2.17	0.44
1:B:1418:LYS:N	1:B:1419:PRO:CD	2.81	0.44
1:B:1138:CYS:O	1:B:1142:LYS:HG3	2.17	0.44
1:B:1415:ILE:O	1:B:1418:LYS:NZ	2.43	0.43
1:A:1118:LEU:HB3	1:A:1233:LEU:HD13	1.99	0.43
1:A:1254:TYR:N	1:A:1255:PRO:CD	2.82	0.42
1:B:1418:LYS:HE2	1:B:1446:TYR:CE2	2.56	0.41
1:A:1377:VAL:CG1	1:A:1410:VAL:HG21	2.50	0.41
1:B:1190:LEU:O	1:B:1194:ARG:HG3	2.21	0.41
1:B:1254:TYR:HB2	1:B:1255:PRO:HD3	2.01	0.41
1:B:1292:ALA:N	1:B:1293:PRO:CD	2.83	0.41
1:B:1437:GLN:N	1:B:1437:GLN:OE1	2.53	0.41
1:B:1141:TYR:OH	1:B:1195:THR:HG21	2.21	0.41
1:A:1342:LEU:HD23	1:A:1342:LEU:HA	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/395 (89%)	338 (97%)	12 (3%)	0	100	100
1	B	328/395 (83%)	314 (96%)	14 (4%)	0	100	100
All	All	678/790 (86%)	652 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/347 (92%)	307 (96%)	11 (4%)	32	42
1	B	303/347 (87%)	292 (96%)	11 (4%)	31	41
All	All	621/694 (90%)	599 (96%)	22 (4%)	32	42

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1111	LYS
1	A	1118	LEU
1	A	1190	LEU
1	A	1233	LEU
1	A	1244	LEU
1	A	1250	VAL
1	A	1297	ARG
1	A	1352	SER
1	A	1429	GLU
1	A	1437	GLN
1	A	1499	MET
1	B	1112	GLN
1	B	1113	GLN
1	B	1116	GLN
1	B	1135	VAL
1	B	1177	VAL
1	B	1233	LEU
1	B	1250	VAL
1	B	1283	SER
1	B	1364	LEU
1	B	1369	CYS
1	B	1483	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1109	ASN
1	A	1193	GLN
1	A	1405	HIS
1	A	1417	GLN
1	B	1249	GLN
1	B	1331	ASN
1	B	1405	HIS
1	B	1416	HIS
1	B	1417	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/395 (90%)	0.50	12 (3%) 48 55	47, 61, 101, 117	0
1	B	336/395 (85%)	0.70	22 (6%) 25 28	48, 68, 112, 131	0
All	All	692/790 (87%)	0.60	34 (4%) 35 41	47, 64, 107, 131	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1244	LEU	3.9
1	A	1241	LEU	3.3
1	B	1172	PRO	3.3
1	B	1499	MET	3.3
1	B	1112	GLN	3.1
1	B	1319	TRP	3.0
1	B	1472	SER	3.0
1	B	1233	LEU	3.0
1	A	1105	GLN	2.8
1	B	1254	TYR	2.7
1	A	1499	MET	2.7
1	B	1146	HIS	2.7
1	B	1495	ILE	2.6
1	A	1247	LEU	2.5
1	A	1131	GLY	2.5
1	A	1106	LYS	2.4
1	B	1295	THR	2.4
1	B	1150	PHE	2.3
1	B	1235	PHE	2.3
1	B	1492	VAL	2.3
1	B	1241	LEU	2.3
1	B	1152	VAL	2.3
1	B	1485	ASP	2.2
1	B	1252	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1236	LEU	2.1
1	B	1145	LEU	2.1
1	B	1257	LEU	2.1
1	B	1236	LEU	2.1
1	A	1427	THR	2.0
1	A	1122	CYS	2.0
1	A	1423	LEU	2.0
1	B	1469	THR	2.0
1	A	1108	LEU	2.0
1	B	1430	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.