



Full wwPDB EM Validation Report ⓘ

Mar 10, 2026 – 01:12 AM UTC

PDB ID : 7KFR / pdb_00007kfr
EMDB ID : EMD-22854
Title : Adeno-Associated Virus (AAV-DJ) - cryo-EM structure at 1.56 Angstrom Resolution
Authors : Xie, Q.; Yoshioka, C.K.; Chapman, M.S.
Deposited on : 2020-10-14
Resolution : 1.56 Å(reported)
Based on initial model : 5UF6

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Q-score	-	25397	300 (1.09 - 2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8954 atoms, of which 4496 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	A	521	8158	2646	3968	721	809	14	8	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	262	ASN	-	insertion	UNP Q6JC41
A	264	THR	GLN	conflict	UNP Q6JC41
A	266	GLY	-	insertion	UNP Q6JC41
A	268	SER	ALA	conflict	UNP Q6JC41
A	273	ALA	HIS	conflict	UNP Q6JC41
A	314	SER	ASN	conflict	UNP Q6JC41
A	329	GLU	ASP	conflict	UNP Q6JC41
A	332	LYS	THR	conflict	UNP Q6JC41
A	342	ILE	VAL	conflict	UNP Q6JC41
A	374	ILE	VAL	conflict	UNP Q6JC41
A	412	GLN	THR	conflict	UNP Q6JC41
A	414	THR	SER	conflict	UNP Q6JC41
A	451	GLN	ASN	conflict	UNP Q6JC41
A	453	THR	PRO	conflict	UNP Q6JC41
A	454	GLY	SER	conflict	UNP Q6JC41
A	458	ASN	THR	conflict	UNP Q6JC41
A	459	THR	GLN	conflict	UNP Q6JC41
A	460	GLN	SER	conflict	UNP Q6JC41
A	461	THR	ARG	conflict	UNP Q6JC41
A	463	GLY	GLN	conflict	UNP Q6JC41
A	467	GLY	ALA	conflict	UNP Q6JC41
A	469	PRO	ALA	conflict	UNP Q6JC41
A	470	ASN	SER	conflict	UNP Q6JC41
A	471	THR	ASP	conflict	UNP Q6JC41
A	472	MET	ILE	conflict	UNP Q6JC41
A	473	ALA	ARG	conflict	UNP Q6JC41
A	474	ASN	ASP	conflict	UNP Q6JC41
A	476	ALA	SER	conflict	UNP Q6JC41

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	477	LYS	ARG	conflict	UNP Q6JC41
A	587	ARG	GLY	conflict	UNP Q6JC41
A	590	ARG	THR	conflict	UNP Q6JC41
A	658	ASP	ASN	conflict	UNP Q6JC41
A	660	PRO	SER	conflict	UNP Q6JC41
A	664	ASN	SER	conflict	UNP Q6JC41
A	665	GLN	ALA	conflict	UNP Q6JC41
A	666	SER	ALA	conflict	UNP Q6JC41
A	668	LEU	PHE	conflict	UNP Q6JC41
A	669	ASN	ALA	conflict	UNP Q6JC41
A	707	TYR	ASN	conflict	UNP Q6JC41
A	710	THR	VAL	conflict	UNP Q6JC41
A	711	SER	ASN	conflict	UNP Q6JC41
A	715	ALA	THR	conflict	UNP Q6JC41
A	717	ASN	ASP	conflict	UNP Q6JC41
A	719	GLU	ASN	conflict	UNP Q6JC41

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
2	A	3	Total Mg 3 3	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	AltConf
3	A	265	Total H O 793 528 265	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Capsid protein VP1



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	48209	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	30	Depositor
Minimum defocus (nm)	-800	Depositor
Maximum defocus (nm)	-2600	Depositor
Magnification	154400	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.372	Depositor
Minimum map value	-0.177	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.018	Depositor
Recommended contour level	0.064	Depositor
Map size ($\text{\AA}$)	306.30002, 306.30002, 306.30002	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.5105, 0.5105, 0.5105	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.67	33/4337 (0.8%)	1.17	5/5911 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	VAL	CA-CB	6.98	1.63	1.54
1	A	457	THR	CA-CB	6.77	1.62	1.52
1	A	420	VAL	CA-CB	6.74	1.62	1.54
1	A	237	ASP	CA-CB	6.67	1.62	1.54
1	A	329	GLU	CA-CB	6.28	1.63	1.53
1	A	639	PHE	CG-CD1	6.19	1.51	1.38
1	A	221	VAL	CA-C	5.88	1.60	1.52
1	A	334	ILE	CA-CB	5.87	1.61	1.54
1	A	345	PHE	CG-CD1	5.75	1.50	1.38
1	A	219	ASP	CA-C	5.67	1.60	1.52
1	A	231	ASP	CA-CB	5.52	1.62	1.53
1	A	536	PHE	CG-CD2	5.51	1.50	1.38
1	A	400	PHE	C-N	5.51	1.40	1.33
1	A	422	PHE	CG-CD1	5.43	1.50	1.38
1	A	680	VAL	C-O	-5.43	1.18	1.24
1	A	312	ARG	NE-CZ	5.36	1.39	1.33
1	A	531	ASP	N-CA	5.34	1.53	1.46
1	A	353	PRO	N-CD	5.32	1.55	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	608	TRP	C-O	-5.30	1.17	1.23
1	A	440	ILE	CA-CB	5.28	1.61	1.54
1	A	663	PHE	CG-CD2	5.27	1.50	1.38
1	A	687	GLU	C-O	-5.23	1.17	1.23
1	A	329	GLU	CA-C	5.21	1.60	1.53
1	A	413	PHE	CG-CD1	5.19	1.49	1.38
1	A	220	GLY	N-CA	5.18	1.50	1.45
1	A	218	ALA	C-N	5.15	1.39	1.33
1	A	344	VAL	CA-CB	5.14	1.61	1.54
1	A	329	GLU	N-CA	5.12	1.54	1.46
1	A	352	LEU	C-N	5.11	1.39	1.33
1	A	279	THR	C-O	-5.07	1.19	1.24
1	A	320	ILE	CA-CB	5.02	1.59	1.54
1	A	682	VAL	CA-CB	5.02	1.61	1.54
1	A	712	VAL	CA-CB	5.01	1.61	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	ASP	CA-CB-CG	6.63	119.23	112.60
1	A	356	LEU	N-CA-C	-5.74	102.40	110.50
1	A	712	VAL	CB-CA-C	5.43	118.65	111.21
1	A	639	PHE	CB-CG-CD2	5.39	129.87	120.70
1	A	290	HIS	CA-CB-CG	5.28	119.08	113.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	510	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4190	3968	3980	10	0
2	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	265	528	0	0	0
All	All	4458	4496	3980	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:PHE:CE2	1:A:374[B]:ILE:HD13	2.43	0.53
1:A:314:SER:OG	1:A:683:GLU:HB3	2.12	0.49
1:A:521:PRO:O	1:A:540:GLY:HA2	2.15	0.47
1:A:622[B]:LYS:HB2	1:A:644:PRO:HG3	1.96	0.47
1:A:616:GLN:HE22	1:A:727:PRO:HA	1.81	0.45
1:A:485:TYR:O	1:A:525:MET:HE1	2.18	0.43
1:A:521:PRO:HA	1:A:539:SER:O	2.18	0.43
1:A:249:LEU:HB2	1:A:374[A]:ILE:HD12	2.01	0.42
1:A:424:SER:HB2	1:A:426:TYR:CE2	2.56	0.41
1:A:427:ALA:O	1:A:734:THR:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	527/521 (101%)	511 (97%)	15 (3%)	1 (0%)	43 24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	231	ASP

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-22854. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 300




Y Index: 300




Z Index: 300


6.2.2 Raw map



X Index: 300



Y Index: 300




Z Index: 300


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 259




Y Index: 341




Z Index: 259


6.3.2 Raw map



X Index: 429



Y Index: 171




Z Index: 429


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.064. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_22854_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 301 nm³; this corresponds to an approximate mass of 272 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of $0.641 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.641 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	1.56	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	1.66	1.85	1.68


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-22854 and PDB model 7KFR. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.064 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.064).

9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.064) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7800	0.8380
A	0.7920	0.8380

