



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:31 PM UTC

PDB ID : 3KG8 / pdb_00003kg8
Title : Dehydratase domain from CurJ module of Curacin polyketide synthase
Authors : Akey, D.L.; Smith, J.L.
Deposited on : 2009-10-28
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4577 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CurJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	0	0	0
			2288	1472	377	431	8			
1	B	277	Total	C	N	O	S	0	0	0
			2215	1430	364	414	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	938	SER	-	expression tag	UNP Q6DNE3
A	939	ASN	-	expression tag	UNP Q6DNE3
A	940	ALA	-	expression tag	UNP Q6DNE3
B	938	SER	-	expression tag	UNP Q6DNE3
B	939	ASN	-	expression tag	UNP Q6DNE3
B	940	ALA	-	expression tag	UNP Q6DNE3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	60	Total	O	0	0
			60	60		
2	B	14	Total	O	0	0
			14	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CurJ



• Molecule 1: CurJ



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.42Å 70.34Å 174.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.69 – 2.45 43.69 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.7 (43.69-2.45) 97.7 (43.69-2.45)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.45Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.195 , 0.258 0.193 , 0.254	Depositor DCC
R_{free} test set	1115 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtriage
Anisotropy	0.379	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 57.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4577	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	1/2341 (0.0%)	0.80	1/3176 (0.0%)
1	B	0.53	0/2266	0.82	1/3074 (0.0%)
All	All	0.58	1/4607 (0.0%)	0.81	2/6250 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1080	ASN	CG-ND2	7.76	1.49	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1042	PHE	N-CA-C	5.81	118.01	109.24
1	A	1167	LEU	N-CA-C	5.37	117.14	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	0	2250	17	0
1	B	2215	0	2185	26	0
2	A	60	0	0	0	0
2	B	14	0	0	0	0
All	All	4577	0	4435	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1068:TRP:CZ3	1:B:957:LEU:HD11	2.34	0.62
1:B:1115:LEU:O	1:B:1120:ARG:NH2	2.34	0.61
1:B:1149:GLN:O	1:B:1190:SER:HB3	2.05	0.56
1:B:981:TYR:CZ	1:B:1112:GLN:HG3	2.41	0.55
1:A:947:PRO:HG3	1:A:1147:LYS:HB3	1.88	0.55
1:B:1107:HIS:O	1:B:1111:ARG:HG2	2.06	0.55
1:A:1135:GLN:NE2	1:A:1195:LEU:HD11	2.24	0.53
1:A:1136:MET:HE2	1:A:1157:SER:HB3	1.90	0.51
1:B:1108:LEU:HD11	1:B:1163:ILE:HD13	1.93	0.50
1:B:1174:ASN:OD1	1:B:1174:ASN:C	2.53	0.50
1:A:1064:HIS:CG	1:A:1065:ILE:N	2.81	0.49
1:A:1089:GLU:H	1:A:1089:GLU:CD	2.19	0.49
1:B:951:GLN:HG2	1:B:964:GLU:HB3	1.95	0.48
1:A:983:GLN:NE2	1:A:1243:LEU:HD13	2.28	0.48
1:B:1009:GLN:HB2	1:B:1079:ALA:HB2	1.95	0.47
1:B:1123:GLU:HG2	1:B:1135:GLN:HE21	1.79	0.47
1:A:1045:ILE:HG22	1:A:1070:VAL:HG22	1.96	0.47
1:A:1135:GLN:HE21	1:A:1195:LEU:HD11	1.80	0.47
1:B:988:GLY:HA2	1:B:991:HIS:CD2	2.50	0.47
1:B:979:ILE:HG13	1:B:1114:HIS:HB2	1.97	0.47
1:A:1137:LYS:HA	1:A:1194:ASP:O	2.16	0.46
1:B:1122:ILE:HG12	1:B:1136:MET:HE2	1.98	0.46
1:A:1088:LEU:HD21	1:A:1197:TRP:CD1	2.50	0.46
1:B:975:LEU:HD22	1:B:986:VAL:HG22	1.98	0.45
1:B:1040:PHE:O	1:B:1076:LEU:N	2.41	0.44
1:B:1238:ASN:O	1:B:1241:ILE:HG12	2.17	0.44
1:B:999:ALA:HB2	1:B:1076:LEU:HD21	1.99	0.44
1:B:963:PHE:HZ	1:B:1002:THR:CG2	2.31	0.43
1:A:1014:LEU:O	1:A:1072:ALA:HA	2.18	0.43
1:B:1108:LEU:HB3	1:B:1113:ILE:HB	2.00	0.43
1:B:1095:CYS:O	1:B:1126:TRP:HB3	2.19	0.42
1:B:1137:LYS:HA	1:B:1194:ASP:O	2.19	0.42
1:B:1213:LYS:HD2	1:B:1231:GLY:HA2	2.01	0.42
1:A:1013:ILE:HA	1:A:1073:THR:O	2.19	0.42
1:A:1045:ILE:HD13	1:B:957:LEU:HD13	2.02	0.42
1:B:1158:CYS:SG	1:B:1218:LEU:HD13	2.59	0.42
1:A:1243:LEU:HD23	1:A:1243:LEU:HA	1.89	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1132:VAL:HG23	1:B:1165:LEU:HD21	2.01	0.41
1:A:1064:HIS:CG	1:A:1065:ILE:H	2.38	0.41
1:A:1172:ASN:HB3	1:A:1175:GLU:HB2	2.03	0.41
1:B:981:TYR:CE1	1:B:1112:GLN:HG3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/308 (91%)	272 (97%)	9 (3%)	0	100	100
1	B	269/308 (87%)	254 (94%)	15 (6%)	0	100	100
All	All	550/616 (89%)	526 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/275 (93%)	238 (93%)	18 (7%)	14	18
1	B	248/275 (90%)	236 (95%)	12 (5%)	23	34
All	All	504/550 (92%)	474 (94%)	30 (6%)	17	26

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	951	GLN
1	A	959	LYS
1	A	960	GLU
1	A	1006	THR
1	A	1007	GLU
1	A	1050	SER
1	A	1051	LEU
1	A	1064	HIS
1	A	1065	ILE
1	A	1075	LYS
1	A	1089	GLU
1	A	1137	LYS
1	A	1146	THR
1	A	1162	ILE
1	A	1168	ASP
1	A	1204	LYS
1	A	1213	LYS
1	A	1241	ILE
1	B	950	ASN
1	B	1006	THR
1	B	1021	ILE
1	B	1023	GLU
1	B	1083	GLN
1	B	1098	LYS
1	B	1120	ARG
1	B	1127	LEU
1	B	1135	GLN
1	B	1138	VAL
1	B	1162	ILE
1	B	1174	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	950	ASN
1	A	1228	GLN
1	B	991	HIS
1	B	1092	GLN
1	B	1106	GLN
1	B	1107	HIS
1	B	1135	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1228	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	287/308 (93%)	-0.06	6 (2%) 63 65	26, 39, 66, 87	0
1	B	277/308 (89%)	0.32	5 (1%) 67 69	28, 56, 71, 86	3 (1%)
All	All	564/616 (91%)	0.12	11 (1%) 65 66	26, 49, 71, 87	3 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1051	LEU	4.8
1	B	1204	LYS	4.3
1	B	1207	GLN	4.3
1	B	959	LYS	4.2
1	B	1079	ALA	2.6
1	A	1244	MET	2.6
1	A	1079	ALA	2.5
1	A	1242	LEU	2.2
1	A	1083	GLN	2.2
1	A	1087	PRO	2.1
1	B	1008	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.