



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:38 PM UTC

PDB ID : 4KGG / pdb_00004kgg
Title : Crystal structure of light mutant2 and dcr3 complex
Authors : Liu, W.; Bonanno, J.B.; Zhan, C.; Kumar, P.R.; Toro, R.; Nathenson, S.G.;
Almo, S.C.; Atoms-to-Animals: The Immune Function Network (IFN); New
York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2013-04-29
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	174	
1	D	174	
2	A	158	
2	B	158	
3	E	4	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	2	 100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4747 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor receptor superfamily member 6B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	165	Total	C	N	O	S	0	0	0
			1244	753	237	236	18			
1	D	163	Total	C	N	O	S	0	0	0
			1233	747	235	233	18			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	196	THR	-	expression tag	UNP O95407
C	197	GLY	-	expression tag	UNP O95407
C	198	HIS	-	expression tag	UNP O95407
C	199	HIS	-	expression tag	UNP O95407
C	200	HIS	-	expression tag	UNP O95407
C	201	HIS	-	expression tag	UNP O95407
C	202	HIS	-	expression tag	UNP O95407
C	203	HIS	-	expression tag	UNP O95407
D	196	THR	-	expression tag	UNP O95407
D	197	GLY	-	expression tag	UNP O95407
D	198	HIS	-	expression tag	UNP O95407
D	199	HIS	-	expression tag	UNP O95407
D	200	HIS	-	expression tag	UNP O95407
D	201	HIS	-	expression tag	UNP O95407
D	202	HIS	-	expression tag	UNP O95407
D	203	HIS	-	expression tag	UNP O95407


- Molecule 2 is a protein called Tumor necrosis factor ligand superfamily member 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	141	Total	C	N	O	S	0	0	0
			1080	691	185	201	3			
2	B	139	Total	C	N	O	S	0	0	0
			1065	680	183	199	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	SER	ARG	engineered mutation	UNP O43557
A	196	ASN	VAL	engineered mutation	UNP O43557
A	198	PHE	TRP	engineered mutation	UNP O43557
A	214	GLU	LYS	engineered mutation	UNP O43557
B	195	SER	ARG	engineered mutation	UNP O43557
B	196	ASN	VAL	engineered mutation	UNP O43557
B	198	PHE	TRP	engineered mutation	UNP O43557
B	214	GLU	LYS	engineered mutation	UNP O43557

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Mg	0	0
			1	1		
5	D	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	9	Total 9	O 9	0	0
6	D	8	Total 8	O 8	0	0
6	A	17	Total 17	O 17	0	0
6	B	11	Total 11	O 11	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor receptor superfamily member 6B

Chain C: 




- Molecule 1: Tumor necrosis factor receptor superfamily member 6B

Chain D: 



- Molecule 2: Tumor necrosis factor ligand superfamily member 14

Chain A: 



- Molecule 2: Tumor necrosis factor ligand superfamily member 14

Chain B: 



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  25% 75%

MAG1
MAG2
BGLA3
MAIN4

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	148.75Å 148.75Å 148.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.63 – 2.78 49.63 – 2.78	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.63-2.78) 99.8 (49.63-2.78)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.178 , 0.235 0.184 , 0.230	Depositor DCC
R_{free} test set	1401 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.036 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4747	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MG, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	1.12	0/1282	1.13	5/1751 (0.3%)
1	D	0.99	0/1271	1.11	2/1736 (0.1%)
2	A	1.17	1/1104 (0.1%)	1.21	7/1496 (0.5%)
2	B	1.20	1/1088 (0.1%)	1.19	4/1473 (0.3%)
All	All	1.12	2/4745 (0.0%)	1.16	18/6456 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	186	PRO	N-CA	-6.58	1.43	1.47
2	A	138	ALA	C-O	-5.52	1.17	1.23

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	196	ASN	N-CA-C	8.22	122.44	110.28
1	C	89	ARG	CG-CD-NE	7.34	128.16	112.00
2	A	110	GLY	CA-C-N	6.40	126.36	120.03
2	A	110	GLY	C-N-CA	6.40	126.36	120.03
2	A	196	ASN	N-CA-C	6.22	119.67	109.72
2	B	104	SER	N-CA-C	5.94	118.54	111.71
2	B	196	ASN	CB-CA-C	-5.75	99.82	109.65
2	A	218	ARG	CB-CG-CD	5.62	124.22	111.30
1	C	89	ARG	NE-CZ-NH2	5.59	124.23	119.20
2	A	206	VAL	CB-CA-C	-5.58	103.99	110.91
2	A	206	VAL	N-CA-C	5.48	116.48	108.58
1	C	83	ASN	N-CA-C	5.45	116.81	108.42
1	C	89	ARG	NE-CZ-NH1	-5.43	116.07	121.50
1	C	56	THR	N-CA-C	5.21	116.57	109.18
1	D	83	ASN	N-CA-C	5.17	116.14	108.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	118	LEU	N-CA-C	5.16	118.53	110.32
1	D	89	ARG	CG-CD-NE	5.02	123.05	112.00
2	B	206	VAL	CB-CA-C	-5.02	104.69	110.91

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1244	0	1133	9	0
1	D	1233	0	1123	18	0
2	A	1080	0	1061	6	0
2	B	1065	0	1043	8	0
3	E	50	0	43	5	0
4	F	28	0	25	1	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	17	0	0	1	0
6	B	11	0	0	0	0
6	C	9	0	0	2	0
6	D	8	0	0	1	0
All	All	4747	0	4428	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:109:HIS:CE1	3:E:4:MAN:H2	2.13	0.83
1:D:109:HIS:HE1	3:E:4:MAN:H2	1.41	0.83
2:A:228:ARG:HD2	6:A:312:HOH:O	1.79	0.81
1:D:81:PHE:CD1	3:E:1:NAG:H83	2.17	0.79
4:F:1:NAG:O3	4:F:2:NAG:O5	2.05	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:89:ARG:NH2	6:D:402:HOH:O	2.21	0.70
1:C:34:PRO:HG3	1:D:175:THR:HB	1.74	0.69
1:C:159:SER:OG	1:C:166:GLU:OE1	2.11	0.64
1:C:43:THR:OG1	1:C:45:GLU:HG3	2.04	0.56
1:D:81:PHE:CG	3:E:1:NAG:H83	2.40	0.56
6:C:408:HOH:O	2:B:218:ARG:HD3	2.05	0.56
1:D:46:ARG:HD2	1:D:46:ARG:H	1.73	0.54
1:D:40:ASP:HB2	1:D:47:LEU:HD11	1.89	0.53
1:D:179:LEU:HD22	1:D:194:THR:O	2.08	0.53
1:C:53:PRO:HD2	1:C:56:THR:OG1	2.09	0.53
1:D:52:CYS:HB3	1:D:56:THR:OG1	2.10	0.52
1:C:146:GLN:HG3	1:D:149:GLN:NE2	2.25	0.51
1:D:54:PRO:HG2	3:E:2:NAG:H2	1.93	0.51
1:D:43:THR:OG1	1:D:45:GLU:OE2	2.29	0.48
2:B:94:PRO:HA	2:B:124:ARG:O	2.14	0.48
2:A:93:ASN:HD21	2:A:239:MET:HG2	1.79	0.47
1:C:132:CYS:O	1:C:162:SER:HA	2.14	0.47
1:D:89:ARG:CG	1:D:89:ARG:HH21	2.28	0.47
1:C:89:ARG:NH2	6:C:403:HOH:O	2.49	0.45
1:C:53:PRO:O	1:C:56:THR:OG1	2.35	0.45
2:A:94:PRO:HG2	2:A:239:MET:HE3	1.98	0.45
2:A:186:PRO:HG2	2:A:197:TRP:CZ3	2.52	0.44
1:C:77:HIS:HB3	1:C:88:CYS:HB2	1.99	0.44
1:D:143:THR:HB	1:D:144:PRO:CD	2.48	0.44
1:D:119:PHE:HA	1:D:127:LEU:O	2.17	0.44
2:B:187:CYS:O	2:B:188:GLY:C	2.61	0.44
2:B:146:LYS:O	2:B:233:SER:HA	2.18	0.43
2:B:129:HIS:O	2:B:130:ASP:C	2.62	0.43
2:B:134:VAL:HG22	2:B:214:GLU:HG2	2.01	0.42
2:B:162:ILE:HD11	2:B:224:LEU:HD13	2.01	0.42
1:D:143:THR:HB	1:D:144:PRO:HD2	2.00	0.42
2:A:135:VAL:HG21	2:A:209:LEU:HB3	2.02	0.41
1:D:159:SER:OG	1:D:166:GLU:OE1	2.35	0.41
2:A:180:LEU:C	2:A:181:VAL:HG23	2.45	0.41
2:B:146:LYS:HA	2:B:201:SER:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	163/174 (94%)	156 (96%)	7 (4%)	0	100	100
1	D	161/174 (92%)	153 (95%)	8 (5%)	0	100	100
2	A	135/158 (85%)	128 (95%)	7 (5%)	0	100	100
2	B	133/158 (84%)	128 (96%)	5 (4%)	0	100	100
All	All	592/664 (89%)	565 (95%)	27 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	138/146 (94%)	128 (93%)	10 (7%)	13	35
1	D	137/146 (94%)	126 (92%)	11 (8%)	11	31
2	A	114/129 (88%)	109 (96%)	5 (4%)	25	56
2	B	112/129 (87%)	107 (96%)	5 (4%)	24	55
All	All	501/550 (91%)	470 (94%)	31 (6%)	16	42

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	42	GLU
1	C	45	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	64	ARG
1	C	89	ARG
1	C	98	ARG
1	C	117	THR
1	C	131	SER
1	C	162	SER
1	C	183	VAL
1	C	192	LEU
1	D	64	ARG
1	D	87	ARG
1	D	89	ARG
1	D	98	ARG
1	D	117	THR
1	D	125	PHE
1	D	163	SER
1	D	175	THR
1	D	183	VAL
1	D	192	LEU
1	D	194	THR
2	A	92	VAL
2	A	156	LEU
2	A	161	THR
2	A	218	ARG
2	A	239	MET
2	B	92	VAL
2	B	105	LEU
2	B	152	VAL
2	B	196	ASN
2	B	210	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	146	GLN
1	C	167	GLN
1	D	51	GLN
1	D	109	HIS
1	D	146	GLN
2	A	93	ASN
2	A	183	GLN
2	B	117	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	E	1	3,1	14,14,15	0.75	0	17,19,21	2.14	5 (29%)
3	NAG	E	2	3	14,14,15	0.73	0	17,19,21	1.21	2 (11%)
3	BMA	E	3	3	11,11,12	0.55	0	15,15,17	1.83	5 (33%)
3	MAN	E	4	3	11,11,12	0.57	0	15,15,17	2.68	3 (20%)
4	NAG	F	1	4,1	14,14,15	0.67	0	17,19,21	2.71	6 (35%)
4	NAG	F	2	4	14,14,15	0.92	0	17,19,21	1.34	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	-	2/2/19/22	0/1/1/1
4	NAG	F	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	F	2	4	-	5/6/23/26	0/1/1/1

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	4	MAN	C1-O5-C5	9.07	124.34	112.19
4	F	1	NAG	C1-O5-C5	7.96	122.86	112.19
3	E	1	NAG	C1-C2-N2	-5.16	102.30	110.43
4	F	1	NAG	C8-C7-N2	4.04	122.82	116.12
3	E	1	NAG	C1-O5-C5	3.95	117.48	112.19
4	F	1	NAG	O5-C1-C2	-3.88	105.28	111.29
3	E	3	BMA	O3-C3-C4	3.75	119.22	110.38
3	E	1	NAG	O5-C1-C2	-3.53	105.83	111.29
3	E	3	BMA	C1-C2-C3	-3.41	104.68	109.64
4	F	1	NAG	C6-C5-C4	-2.90	105.89	113.02
3	E	4	MAN	C2-C3-C4	2.71	115.63	110.86
3	E	3	BMA	C1-O5-C5	2.59	115.65	112.19
3	E	3	BMA	O2-C2-C3	2.45	115.23	110.15
3	E	2	NAG	O7-C7-N2	2.45	126.31	121.98
3	E	3	BMA	O3-C3-C2	2.39	114.93	110.05
4	F	1	NAG	O7-C7-C8	-2.39	117.80	122.05
4	F	1	NAG	C4-C3-C2	-2.31	107.64	111.02
3	E	1	NAG	C2-N2-C7	2.29	125.97	122.90
3	E	1	NAG	O7-C7-N2	2.24	125.93	121.98
4	F	2	NAG	O5-C1-C2	-2.13	108.00	111.29
3	E	2	NAG	C8-C7-N2	-2.12	112.61	116.12
3	E	4	MAN	O3-C3-C4	-2.03	105.58	110.38

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	F	2	NAG	C3-C2-N2-C7
3	E	4	MAN	C4-C5-C6-O6
4	F	1	NAG	C4-C5-C6-O6
3	E	4	MAN	O5-C5-C6-O6
4	F	2	NAG	O5-C5-C6-O6
4	F	1	NAG	C8-C7-N2-C2
4	F	1	NAG	O7-C7-N2-C2
4	F	1	NAG	O5-C5-C6-O6
3	E	1	NAG	C4-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
4	F	2	NAG	C4-C5-C6-O6
4	F	2	NAG	C8-C7-N2-C2
4	F	2	NAG	O7-C7-N2-C2


There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	4	MAN	2	0
3	E	2	NAG	1	0
4	F	1	NAG	1	0
4	F	2	NAG	1	0
3	E	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	C	165/174 (94%)	0.01	5 (3%)	52 45	25, 43, 74, 98	0
1	D	163/174 (93%)	0.59	14 (8%)	16 13	31, 61, 105, 129	0
2	A	141/158 (89%)	-0.30	6 (4%)	40 33	15, 28, 71, 115	0
2	B	139/158 (87%)	-0.38	6 (4%)	40 33	18, 31, 65, 87	0
All	All	608/664 (91%)	0.00	31 (5%)	33 27	15, 40, 83, 129	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	156	LEU	9.1
2	B	188	GLY	5.6
2	B	159	ALA	5.1
2	A	188	GLY	4.9
2	A	195	SER	4.8
2	A	155	PRO	4.7
1	D	195	SER	4.5
2	A	159	ALA	4.3
1	D	64	ARG	4.3
1	D	41	ALA	4.2
2	B	154	CYS	4.1
2	A	92	VAL	3.3
2	B	195	SER	3.3
1	D	43	THR	3.2
1	D	123	ALA	3.1
1	D	179	LEU	3.0
1	C	197	GLY	2.9
1	D	46	ARG	2.8
2	B	196	ASN	2.8
1	D	63	ARG	2.8
1	D	48	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	122	HIS	2.5
1	C	64	ARG	2.4
1	D	42	GLU	2.4
2	B	92	VAL	2.4
1	D	33	THR	2.2
1	C	146	GLN	2.2
1	C	42	GLU	2.2
1	D	87	ARG	2.1
1	D	194	THR	2.0
1	D	178	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	F	2	14/15	0.51	0.26	65,78,90,91	0
4	NAG	F	1	14/15	0.68	0.19	52,61,66,71	0
3	BMA	E	3	11/12	0.73	0.21	71,76,79,84	0
3	MAN	E	4	11/12	0.80	0.23	62,84,98,101	0
3	NAG	E	1	14/15	0.81	0.19	49,57,65,68	0
3	NAG	E	2	14/15	0.84	0.19	61,69,74,80	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	C	301	1/1	0.95	0.15	46,46,46,46	0
5	MG	D	301	1/1	0.97	0.10	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.