



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:40 PM UTC

PDB ID : 4KGR / pdb_00004kgr
Title : Backbone Modifications in the Protein GB1 Helix: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35
Authors : Reinert, Z.E.; Lengyel, G.A.; Horne, W.S.
Deposited on : 2013-04-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	57	
1	B	57	
1	C	57	
1	D	57	
1	E	57	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	57	 86%9% . .
1	G	57	 82%12%5%
1	H	57	 74%18%9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5980 atoms, of which 2249 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	57	Total	C	H	N	O	0	0	1
			720	276	280	68	96			
1	B	57	Total	C	H	N	O	0	0	1
			720	276	280	68	96			
1	C	57	Total	C	H	N	O	0	0	1
			720	276	280	68	96			
1	D	57	Total	C	H	N	O	0	1	1
			726	279	282	68	97			
1	E	57	Total	C	H	N	O	0	0	1
			720	276	280	68	96			
1	F	57	Total	C	H	N	O	0	1	1
			726	279	282	68	97			
1	G	57	Total	C	H	N	O	0	0	1
			720	276	280	68	96			
1	H	57	Total	C	H	N	O	0	0	1
			720	276	280	68	96			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	H	1	Total	C	H	O	0	0
			11	3	5	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	34	Total	O	0	0
			34	34		
3	B	25	Total	O	0	0
			25	25		
3	C	27	Total	O	0	0
			27	27		
3	D	19	Total	O	0	0
			19	19		
3	E	16	Total	O	0	0
			16	16		
3	F	22	Total	O	0	0
			22	22		
3	G	34	Total	O	0	0
			34	34		
3	H	20	Total	O	0	0
			20	20		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain A: 




- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain B: 




- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain C: 




- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain D: 




- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain E: 




- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain F: 86% 9% . .




- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain G: 82% 12% 5%



- Molecule 1: Streptococcal Protein GB1 Backbone Modified Variant: beta-3-Ala24, beta-3-Lys28, beta-3-Lys31, beta-3-Asn35

Chain H: 74% 18% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.16Å 81.17Å 52.06Å 90.00° 89.97° 90.00°	Depositor
Resolution (Å)	32.03 – 2.00 32.03 – 2.00	Depositor EDS
% Data completeness (in resolution range)	93.7 (32.03-2.00) 93.1 (32.03-2.00)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.24 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.158 , 0.192 0.160 , 0.163	Depositor DCC
R_{free} test set	1944 reflections (6.62%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 25.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.074 for l,k,-h 0.069 for -h,-k,l 0.479 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5980	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B3X, GOL, B3K, NH2, B3A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/406	0.85	0/547
1	B	0.63	0/406	0.75	0/547
1	C	0.70	0/406	0.79	0/547
1	D	0.75	0/413	0.73	0/557
1	E	0.69	0/406	0.85	0/547
1	F	0.67	0/413	0.79	0/557
1	G	0.71	0/406	0.77	0/547
1	H	0.74	0/406	0.76	0/547
All	All	0.70	0/3262	0.79	0/4396

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
1	B	0	8
1	C	0	8
1	D	0	8
1	E	0	8
1	F	0	8
1	G	0	8
1	H	0	8
All	All	0	64

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (64) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	24	B3A	Peptide,Mainchain
1	A	28	B3K	Peptide,Mainchain
1	A	31	B3K	Peptide,Mainchain
1	A	35	B3X	Peptide,Mainchain
1	B	24	B3A	Peptide,Mainchain
1	B	28	B3K	Peptide,Mainchain
1	B	31	B3K	Peptide,Mainchain
1	B	35	B3X	Peptide,Mainchain
1	C	24	B3A	Peptide,Mainchain
1	C	28	B3K	Peptide,Mainchain
1	C	31	B3K	Peptide,Mainchain
1	C	35	B3X	Peptide,Mainchain
1	D	24	B3A	Peptide,Mainchain
1	D	28	B3K	Peptide,Mainchain
1	D	31	B3K	Peptide,Mainchain
1	D	35	B3X	Peptide,Mainchain
1	E	24	B3A	Peptide,Mainchain
1	E	28	B3K	Peptide,Mainchain
1	E	31	B3K	Peptide,Mainchain
1	E	35	B3X	Peptide,Mainchain
1	F	24	B3A	Peptide,Mainchain
1	F	28	B3K	Peptide,Mainchain
1	F	31	B3K	Peptide,Mainchain
1	F	35	B3X	Peptide,Mainchain
1	G	24	B3A	Peptide,Mainchain
1	G	28	B3K	Peptide,Mainchain
1	G	31	B3K	Peptide,Mainchain
1	G	35	B3X	Peptide,Mainchain
1	H	24	B3A	Peptide,Mainchain
1	H	28	B3K	Peptide,Mainchain
1	H	31	B3K	Peptide,Mainchain
1	H	35	B3X	Peptide,Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	440	280	400	4	0
1	B	440	280	401	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	440	280	402	1	0
1	D	444	282	409	2	0
1	E	440	280	403	4	0
1	F	444	282	408	3	0
1	G	440	280	401	8	0
1	H	440	280	402	9	0
2	H	6	5	8	0	0
3	A	34	0	0	0	0
3	B	25	0	0	0	0
3	C	27	0	0	0	0
3	D	19	0	0	2	0
3	E	16	0	0	1	0
3	F	22	0	0	0	1
3	G	34	0	0	1	1
3	H	20	0	0	4	2
All	All	3731	2249	3234	34	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:4:LYS:NZ	1:G:15:GLU:HB2	1.77	0.99
1:G:4:LYS:HZ3	1:G:15:GLU:HB2	1.40	0.87
1:E:7:LEU:HD21	1:E:33:TYR:HE2	1.56	0.71
1:H:19:GLU:OE2	3:H:216:HOH:O	2.09	0.69
1:H:12:LEU:HD12	3:H:217:HOH:O	1.94	0.68
1:B:39:VAL:HG13	1:B:56:GLU:HG3	1.76	0.67
1:H:4:LYS:HE3	1:H:6:ILE:HD11	1.81	0.63
1:E:47:ASP:OD1	3:E:110:HOH:O	2.17	0.60
1:B:7:LEU:HD11	1:B:33:TYR:HE2	1.66	0.59
1:F:12:LEU:HD22	1:G:16:THR:CG2	2.36	0.55
1:D:21:VAL:HG23	3:D:105:HOH:O	2.06	0.54
1:H:53:THR:HG23	3:H:201:HOH:O	2.07	0.54
1:H:7:LEU:HD11	1:H:33:TYR:HE2	1.73	0.54
1:G:45:TYR:HE2	1:G:47:ASP:OD1	1.91	0.53
1:D:29:VAL:HG21	3:D:114:HOH:O	2.09	0.52
1:G:4:LYS:HZ1	1:G:15:GLU:HB2	1.69	0.51
1:H:7:LEU:HD11	1:H:33:TYR:CE2	2.46	0.50
1:G:4:LYS:CE	1:G:15:GLU:HB2	2.39	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:VAL:HG13	1:B:56:GLU:CG	2.41	0.50
1:G:32:GLN:NE2	3:G:120:HOH:O	2.45	0.49
1:F:39:VAL:HG13	1:F:56:GLU:HG3	1.95	0.49
1:C:31:B3K:O	1:C:35:B3X:HG3	2.14	0.48
1:B:7:LEU:HD11	1:B:33:TYR:CE2	2.48	0.48
1:A:12:LEU:C	1:A:12:LEU:HD23	2.38	0.47
1:E:4:LYS:HG2	1:E:51:THR:HG23	1.97	0.47
1:G:4:LYS:NZ	1:G:15:GLU:CB	2.65	0.46
1:A:4:LYS:HE3	1:A:15:GLU:CD	2.41	0.45
1:A:31:B3K:HD2	1:A:43:TRP:CZ2	2.53	0.44
1:H:31:B3K:HD2	1:H:43:TRP:CZ2	2.53	0.44
1:H:44:THR:HG1	1:H:53:THR:HG1	1.59	0.43
1:F:31:B3K:HF1	1:F:43:TRP:CD2	2.55	0.42
1:H:47:ASP:OD1	3:H:206:HOH:O	2.22	0.42
1:E:9:GLY:HA2	1:E:56:GLU:HB2	2.00	0.42
1:A:3:TYR:CD2	1:A:50:LYS:HB3	2.55	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:109:HOH:O	3:H:216:HOH:O[2_655]	2.07	0.13
3:F:110:HOH:O	3:H:215:HOH:O[2_555]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	51/57 (90%)	50 (98%)	1 (2%)	0	100	100
1	B	51/57 (90%)	50 (98%)	1 (2%)	0	100	100
1	C	51/57 (90%)	49 (96%)	2 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	52/57 (91%)	50 (96%)	2 (4%)	0	100	100
1	E	51/57 (90%)	50 (98%)	1 (2%)	0	100	100
1	F	52/57 (91%)	51 (98%)	1 (2%)	0	100	100
1	G	51/57 (90%)	49 (96%)	2 (4%)	0	100	100
1	H	51/57 (90%)	50 (98%)	1 (2%)	0	100	100
All	All	410/456 (90%)	399 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	43/43 (100%)	43 (100%)	0	100	100
1	B	43/43 (100%)	42 (98%)	1 (2%)	44	49
1	C	43/43 (100%)	43 (100%)	0	100	100
1	D	44/43 (102%)	43 (98%)	1 (2%)	44	49
1	E	43/43 (100%)	41 (95%)	2 (5%)	23	22
1	F	44/43 (102%)	44 (100%)	0	100	100
1	G	43/43 (100%)	43 (100%)	0	100	100
1	H	43/43 (100%)	40 (93%)	3 (7%)	14	10
All	All	346/344 (101%)	339 (98%)	7 (2%)	48	54

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	12	LEU
1	D	32	GLN
1	E	4	LYS
1	E	11	THR
1	H	7	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	12	LEU
1	H	15	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	32	GLN
1	D	32	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

32 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	B3A	G	24	1	5,5,6	0.83	0	5,5,7	2.25	1 (20%)
1	B3K	A	28	1	9,9,10	0.46	0	7,9,11	0.89	0
1	B3X	B	35	1	8,8,9	0.42	0	6,9,11	1.37	1 (16%)
1	B3K	D	28	1	9,9,10	0.58	0	7,9,11	0.94	0
1	B3K	E	31	1	9,9,10	0.47	0	7,9,11	1.34	1 (14%)
1	B3K	C	31	1	9,9,10	0.42	0	7,9,11	1.99	2 (28%)
1	B3K	H	31	1	9,9,10	0.43	0	7,9,11	1.02	0
1	B3X	H	35	1	8,8,9	0.78	0	6,9,11	1.42	1 (16%)
1	B3K	D	31	1	9,9,10	0.25	0	7,9,11	1.58	1 (14%)
1	B3K	F	28	1	9,9,10	0.37	0	7,9,11	1.20	1 (14%)
1	B3K	B	28	1	9,9,10	0.77	0	7,9,11	1.53	2 (28%)
1	B3K	F	31	1	9,9,10	0.58	0	7,9,11	1.43	2 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	B3A	A	24	1	5,5,6	0.87	0	5,5,7	2.07	1 (20%)
1	B3K	G	28	1	9,9,10	0.54	0	7,9,11	1.52	1 (14%)
1	B3A	B	24	1	5,5,6	0.83	0	5,5,7	1.29	0
1	B3K	G	31	1	9,9,10	0.43	0	7,9,11	1.03	0
1	B3A	E	24	1	5,5,6	0.74	0	5,5,7	1.64	1 (20%)
1	B3A	F	24	1	5,5,6	0.69	0	5,5,7	1.14	0
1	B3X	E	35	1	8,8,9	0.49	0	6,9,11	1.70	2 (33%)
1	B3X	D	35	1	8,8,9	0.41	0	6,9,11	1.44	1 (16%)
1	B3X	F	35	1	8,8,9	1.01	1 (12%)	6,9,11	1.37	1 (16%)
1	B3K	E	28	1	9,9,10	0.61	0	7,9,11	0.85	0
1	B3X	C	35	1	8,8,9	0.76	0	6,9,11	1.60	1 (16%)
1	B3X	G	35	1	8,8,9	0.51	0	6,9,11	2.06	2 (33%)
1	B3K	C	28	1	9,9,10	0.33	0	7,9,11	1.12	0
1	B3A	H	24	1	5,5,6	0.97	0	5,5,7	1.81	2 (40%)
1	B3A	C	24	1	5,5,6	1.09	0	5,5,7	1.63	1 (20%)
1	B3K	B	31	1	9,9,10	0.35	0	7,9,11	1.06	1 (14%)
1	B3X	A	35	1	8,8,9	0.61	0	6,9,11	0.90	1 (16%)
1	B3A	D	24	1	5,5,6	1.06	1 (20%)	5,5,7	1.74	2 (40%)
1	B3K	H	28	1	9,9,10	0.45	0	7,9,11	1.03	0
1	B3K	A	31	1	9,9,10	0.68	0	7,9,11	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	B3A	G	24	1	-	2/3/3/4	-
1	B3K	A	28	1	-	4/8/8/9	-
1	B3X	B	35	1	-	2/7/7/8	-
1	B3K	D	28	1	-	5/8/8/9	-
1	B3K	E	31	1	-	1/8/8/9	-
1	B3K	C	31	1	-	2/8/8/9	-
1	B3K	H	31	1	-	2/8/8/9	-
1	B3X	H	35	1	-	6/7/7/8	-
1	B3K	D	31	1	-	1/8/8/9	-
1	B3K	F	28	1	-	3/8/8/9	-
1	B3K	B	28	1	-	4/8/8/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	B3K	F	31	1	-	4/8/8/9	-
1	B3A	A	24	1	-	2/3/3/4	-
1	B3K	G	28	1	-	3/8/8/9	-
1	B3A	B	24	1	-	1/3/3/4	-
1	B3K	G	31	1	-	3/8/8/9	-
1	B3A	E	24	1	-	1/3/3/4	-
1	B3A	F	24	1	-	2/3/3/4	-
1	B3X	E	35	1	-	1/7/7/8	-
1	B3X	D	35	1	-	3/7/7/8	-
1	B3X	F	35	1	-	4/7/7/8	-
1	B3K	E	28	1	-	2/8/8/9	-
1	B3X	C	35	1	-	5/7/7/8	-
1	B3X	G	35	1	-	4/7/7/8	-
1	B3K	C	28	1	-	4/8/8/9	-
1	B3A	H	24	1	-	1/3/3/4	-
1	B3A	C	24	1	-	1/3/3/4	-
1	B3K	B	31	1	-	1/8/8/9	-
1	B3X	A	35	1	-	3/7/7/8	-
1	B3A	D	24	1	-	1/3/3/4	-
1	B3K	H	28	1	-	4/8/8/9	-
1	B3K	A	31	1	-	5/8/8/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	35	B3X	CB-CA	-2.38	1.50	1.53
1	D	24	B3A	CA-N	-2.07	1.43	1.49

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	24	B3A	CG-CA-CB	-4.65	105.91	111.86
1	C	31	B3K	CA-CB-C	4.41	119.30	112.17
1	A	24	B3A	CG-CA-CB	-4.12	106.59	111.86
1	G	35	B3X	CB-CA-CG	3.57	115.86	110.82
1	G	28	B3K	CA-CB-C	3.47	117.77	112.17
1	C	35	B3X	CB-CA-CG	-3.20	106.30	110.82
1	D	35	B3X	CB-CA-CG	-3.11	106.43	110.82
1	D	31	B3K	CA-CB-C	3.08	117.15	112.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	28	B3K	CA-CB-C	-3.06	107.24	112.17
1	C	24	B3A	CA-CB-C	2.97	119.38	113.29
1	G	35	B3X	CA-CB-C	-2.92	107.46	112.17
1	E	24	B3A	CA-CB-C	2.86	119.14	113.29
1	B	35	B3X	CB-CA-CG	-2.83	106.83	110.82
1	H	24	B3A	CA-CB-C	2.81	119.05	113.29
1	E	35	B3X	CA-CB-C	2.73	116.57	112.17
1	F	31	B3K	CA-CB-C	2.72	116.57	112.17
1	F	35	B3X	CA-CB-C	2.72	116.56	112.17
1	D	24	B3A	CG-CA-CB	-2.59	108.55	111.86
1	F	28	B3K	CA-CB-C	2.50	116.20	112.17
1	H	24	B3A	CG-CA-CB	-2.46	108.71	111.86
1	H	35	B3X	CB-CA-CG	2.44	114.26	110.82
1	B	28	B3K	O-C-CB	-2.40	118.41	125.38
1	E	31	B3K	CA-CB-C	2.31	115.90	112.17
1	D	24	B3A	CA-CB-C	2.29	117.99	113.29
1	C	31	B3K	CG-CA-N	2.26	116.05	109.16
1	A	35	B3X	O-C-CB	-2.11	119.24	125.38
1	E	35	B3X	CB-CA-CG	-2.10	107.85	110.82
1	F	31	B3K	O-C-CB	-2.08	119.33	125.38
1	B	31	B3K	O-C-CB	-2.07	119.34	125.38

There are no chirality outliers.

All (87) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	28	B3K	CB-CA-CG-CD
1	A	31	B3K	N-CA-CG-CD
1	A	31	B3K	CB-CA-CG-CD
1	A	35	B3X	N-CA-CG-CD
1	C	28	B3K	CB-CA-CG-CD
1	C	35	B3X	N-CA-CG-CD
1	D	28	B3K	CD-CE-CF-NZ
1	F	35	B3X	N-CA-CG-CD
1	F	35	B3X	CB-CA-CG-CD
1	H	35	B3X	N-CA-CG-CD
1	H	35	B3X	CB-CA-CG-CD
1	F	31	B3K	CG-CD-CE-CF
1	B	28	B3K	CG-CD-CE-CF
1	C	35	B3X	NE2-CD-CG-CA
1	C	28	B3K	CG-CD-CE-CF
1	G	35	B3X	CB-CA-CG-CD

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	G	31	B3K	CG-CD-CE-CF
1	C	35	B3X	OE1-CD-CG-CA
1	D	35	B3X	NE2-CD-CG-CA
1	H	35	B3X	NE2-CD-CG-CA
1	C	28	B3K	CD-CE-CF-NZ
1	B	28	B3K	CG-CA-CB-C
1	G	35	B3X	CG-CA-CB-C
1	D	35	B3X	OE1-CD-CG-CA
1	H	35	B3X	OE1-CD-CG-CA
1	A	24	B3A	N-CA-CB-C
1	G	24	B3A	N-CA-CB-C
1	A	24	B3A	O-C-CB-CA
1	A	28	B3K	O-C-CB-CA
1	A	31	B3K	O-C-CB-CA
1	A	35	B3X	O-C-CB-CA
1	B	24	B3A	O-C-CB-CA
1	B	31	B3K	O-C-CB-CA
1	B	35	B3X	O-C-CB-CA
1	C	28	B3K	O-C-CB-CA
1	C	35	B3X	O-C-CB-CA
1	D	24	B3A	O-C-CB-CA
1	D	28	B3K	N-CA-CG-CD
1	D	28	B3K	CB-CA-CG-CD
1	E	24	B3A	O-C-CB-CA
1	E	28	B3K	O-C-CB-CA
1	E	31	B3K	O-C-CB-CA
1	F	24	B3A	O-C-CB-CA
1	F	31	B3K	N-CA-CG-CD
1	F	31	B3K	CB-CA-CG-CD
1	F	31	B3K	O-C-CB-CA
1	F	35	B3X	O-C-CB-CA
1	G	31	B3K	CB-CA-CG-CD
1	G	31	B3K	O-C-CB-CA
1	G	35	B3X	O-C-CB-CA
1	H	24	B3A	O-C-CB-CA
1	H	28	B3K	O-C-CB-CA
1	H	31	B3K	O-C-CB-CA
1	H	35	B3X	O-C-CB-CA
1	F	28	B3K	CD-CE-CF-NZ
1	A	35	B3X	N-CA-CB-C
1	B	28	B3K	N-CA-CB-C
1	G	35	B3X	N-CA-CB-C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	H	31	B3K	CD-CE-CF-NZ
1	A	28	B3K	CG-CD-CE-CF
1	H	28	B3K	CG-CD-CE-CF
1	B	28	B3K	O-C-CB-CA
1	C	31	B3K	O-C-CB-CA
1	D	28	B3K	O-C-CB-CA
1	D	31	B3K	O-C-CB-CA
1	D	35	B3X	O-C-CB-CA
1	E	35	B3X	O-C-CB-CA
1	G	24	B3A	O-C-CB-CA
1	G	28	B3K	O-C-CB-CA
1	A	31	B3K	CE-CD-CG-CA
1	G	28	B3K	CD-CE-CF-NZ
1	A	28	B3K	N-CA-CB-C
1	B	35	B3X	N-CA-CB-C
1	D	28	B3K	N-CA-CB-C
1	E	28	B3K	N-CA-CB-C
1	F	28	B3K	N-CA-CB-C
1	F	35	B3X	N-CA-CB-C
1	G	28	B3K	N-CA-CB-C
1	H	28	B3K	N-CA-CB-C
1	H	35	B3X	N-CA-CB-C
1	C	31	B3K	CG-CD-CE-CF
1	H	28	B3K	CD-CE-CF-NZ
1	F	24	B3A	N-CA-CB-C
1	A	31	B3K	CG-CD-CE-CF
1	C	24	B3A	O-C-CB-CA
1	F	28	B3K	O-C-CB-CA
1	C	35	B3X	CB-CA-CG-CD

There are no ring outliers.

5 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	31	B3K	1	0
1	H	31	B3K	1	0
1	F	31	B3K	1	0
1	C	35	B3X	1	0
1	A	31	B3K	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	H	101	-	5,5,5	0.23	0	5,5,5	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	H	101	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	101	GOL	C1-C2-C3-O3
2	H	101	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	52/57 (91%)	-1.62	0 100 100	14, 20, 36, 48	0
1	B	52/57 (91%)	-1.59	0 100 100	14, 22, 43, 51	0
1	C	52/57 (91%)	-1.64	0 100 100	14, 21, 35, 52	0
1	D	52/57 (91%)	-1.56	0 100 100	14, 24, 47, 56	1 (1%)
1	E	52/57 (91%)	-1.57	0 100 100	12, 21, 49, 72	0
1	F	52/57 (91%)	-1.61	0 100 100	12, 19, 34, 48	1 (1%)
1	G	52/57 (91%)	-1.58	0 100 100	15, 22, 38, 54	0
1	H	52/57 (91%)	-1.61	0 100 100	12, 22, 39, 49	0
All	All	416/456 (91%)	-1.60	0 100 100	12, 22, 44, 72	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	B3X	G	35	9/10	0.98	0.05	21,35,63,68	0
1	B3K	A	28	10/11	0.99	0.03	14,26,45,54	0
1	B3K	A	31	10/11	0.99	0.03	12,26,31,37	0
1	B3X	A	35	9/10	0.99	0.03	11,19,40,40	0
1	B3A	B	24	6/7	0.99	0.03	8,15,17,18	0
1	B3K	B	28	10/11	0.99	0.03	14,32,48,53	0
1	B3X	B	35	9/10	0.99	0.04	21,30,48,48	0
1	B3A	C	24	6/7	0.99	0.02	12,16,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	B3K	C	31	10/11	0.99	0.02	17,26,45,54	0
1	B3X	C	35	9/10	0.99	0.03	18,30,42,44	0
1	B3A	D	24	6/7	0.99	0.03	16,21,26,28	0
1	B3K	D	28	10/11	0.99	0.06	16,53,99,101	0
1	B3X	D	35	9/10	0.99	0.03	22,26,44,44	0
1	B3X	E	35	9/10	0.99	0.02	13,16,27,30	0
1	B3A	F	24	6/7	0.99	0.03	11,15,21,25	0
1	B3X	F	35	9/10	0.99	0.03	11,25,42,51	0
1	B3K	G	28	10/11	0.99	0.04	21,29,37,41	0
1	B3A	A	24	6/7	0.99	0.03	9,11,16,17	0
1	B3A	H	24	6/7	0.99	0.02	12,17,26,31	0
1	B3K	H	28	10/11	0.99	0.03	14,24,28,30	0
1	B3K	H	31	10/11	0.99	0.03	13,23,36,40	0
1	B3X	H	35	9/10	0.99	0.03	17,31,58,58	0
1	B3K	F	31	10/11	1.00	0.03	16,22,31,33	0
1	B3K	B	31	10/11	1.00	0.02	17,26,31,32	0
1	B3A	G	24	6/7	1.00	0.03	16,19,21,25	0
1	B3A	E	24	6/7	1.00	0.02	13,18,26,27	0
1	B3K	G	31	10/11	1.00	0.02	9,23,39,40	0
1	B3K	E	28	10/11	1.00	0.02	11,17,25,29	0
1	B3K	E	31	10/11	1.00	0.02	12,15,19,25	0
1	B3K	C	28	10/11	1.00	0.03	10,28,49,59	0
1	B3K	D	31	10/11	1.00	0.02	11,22,34,37	0
1	B3K	F	28	10/11	1.00	0.02	12,18,36,40	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	H	101	6/6	0.99	0.03	21,27,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.