



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:14 PM UTC

PDB ID : 3KHJ / pdb_00003khj
Title : C. parvum inosine monophosphate dehydrogenase bound by inhibitor C64
Authors : MacPherson, I.S.; Hedstrom, L.K.
Deposited on : 2009-10-30
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	
1	B	361	
1	C	361	
1	D	361	
1	E	361	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	361	 <p>3% 72% 10% • 17%</p>
1	G	361	 <p>3% 68% 14% • 17%</p>
1	H	361	 <p>3% 79% 11% • 9%</p>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 18502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inosine-5-monophosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	302	2221	1404	374	431	12	0	0	0
1	B	318	2330	1471	390	454	15	0	1	0
1	C	306	2255	1422	383	438	12	0	1	0
1	D	321	2322	1463	390	454	15	0	0	0
1	E	298	2162	1369	365	416	12	0	0	0
1	F	300	2198	1390	363	433	12	0	1	0
1	G	300	2188	1379	370	428	11	0	0	0
1	H	328	2355	1484	397	460	14	0	0	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q5CPK7
A	-1	SER	-	expression tag	UNP Q5CPK7
A	0	HIS	-	expression tag	UNP Q5CPK7
A	90	SER	-	linker	UNP Q5CPK7
A	91	GLY	-	linker	UNP Q5CPK7
A	92	GLY	-	linker	UNP Q5CPK7
B	-2	GLY	-	expression tag	UNP Q5CPK7
B	-1	SER	-	expression tag	UNP Q5CPK7
B	0	HIS	-	expression tag	UNP Q5CPK7
B	90	SER	-	linker	UNP Q5CPK7
B	91	GLY	-	linker	UNP Q5CPK7
B	92	GLY	-	linker	UNP Q5CPK7
C	-2	GLY	-	expression tag	UNP Q5CPK7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	SER	-	expression tag	UNP Q5CPK7
C	0	HIS	-	expression tag	UNP Q5CPK7
C	90	SER	-	linker	UNP Q5CPK7
C	91	GLY	-	linker	UNP Q5CPK7
C	92	GLY	-	linker	UNP Q5CPK7
D	-2	GLY	-	expression tag	UNP Q5CPK7
D	-1	SER	-	expression tag	UNP Q5CPK7
D	0	HIS	-	expression tag	UNP Q5CPK7
D	90	SER	-	linker	UNP Q5CPK7
D	91	GLY	-	linker	UNP Q5CPK7
D	92	GLY	-	linker	UNP Q5CPK7
E	-2	GLY	-	expression tag	UNP Q5CPK7
E	-1	SER	-	expression tag	UNP Q5CPK7
E	0	HIS	-	expression tag	UNP Q5CPK7
E	90	SER	-	linker	UNP Q5CPK7
E	91	GLY	-	linker	UNP Q5CPK7
E	92	GLY	-	linker	UNP Q5CPK7
F	-2	GLY	-	expression tag	UNP Q5CPK7
F	-1	SER	-	expression tag	UNP Q5CPK7
F	0	HIS	-	expression tag	UNP Q5CPK7
F	90	SER	-	linker	UNP Q5CPK7
F	91	GLY	-	linker	UNP Q5CPK7
F	92	GLY	-	linker	UNP Q5CPK7
G	-2	GLY	-	expression tag	UNP Q5CPK7
G	-1	SER	-	expression tag	UNP Q5CPK7
G	0	HIS	-	expression tag	UNP Q5CPK7
G	90	SER	-	linker	UNP Q5CPK7
G	91	GLY	-	linker	UNP Q5CPK7
G	92	GLY	-	linker	UNP Q5CPK7
H	-2	GLY	-	expression tag	UNP Q5CPK7
H	-1	SER	-	expression tag	UNP Q5CPK7
H	0	HIS	-	expression tag	UNP Q5CPK7
H	90	SER	-	linker	UNP Q5CPK7
H	91	GLY	-	linker	UNP Q5CPK7
H	92	GLY	-	linker	UNP Q5CPK7

- Molecule 2 is INOSINIC ACID (CCD ID: IMP) (formula: C₁₀H₁₃N₄O₈P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 23	10	4	8	1	0	0
2	B	1	Total 23	10	4	8	1	0	0
2	C	1	Total 23	10	4	8	1	0	0
2	D	1	Total 23	10	4	8	1	0	0
2	E	1	Total 23	10	4	8	1	0	0
2	F	1	Total 23	10	4	8	1	0	0
2	G	1	Total 23	10	4	8	1	0	0
2	H	1	Total 23	10	4	8	1	0	0

- Molecule 3 is N-(4-bromophenyl)-2-[2-(1,3-thiazol-2-yl)-1H-benzimidazol-1-yl]acetamide (CCD ID: C64) (formula: C₁₈H₁₃BrN₄OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	Br	C	N	O			S
3	B	1	Total	Br	C	N	O	S	0	0
			25	1	18	4	1	1		
3	D	1	Total	Br	C	N	O	S	0	0
			25	1	18	4	1	1		
3	H	1	Total	Br	C	N	O	S	0	0
			25	1	18	4	1	1		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
4	D	1	Total	C O	0	0
			4	2 2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	37	Total O 37 37	0	0
5	B	38	Total O 38 38	0	0
5	C	41	Total O 41 41	0	0
5	D	40	Total O 40 40	0	0
5	E	13	Total O 13 13	0	0
5	F	17	Total O 17 17	0	0
5	G	9	Total O 9 9	0	0
5	H	13	Total O 13 13	0	0


- Molecule 1: Inosine-5-monophosphate dehydrogenase




- Molecule 1: Inosine-5-monophosphate dehydrogenase



- Molecule 1: Inosine-5-monophosphate dehydrogenase







- Molecule 1: Inosine-5-monophosphate dehydrogenase





- Molecule 1: Inosine-5-monophosphate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.48Å 166.14Å 101.29Å 90.00° 105.14° 90.00°	Depositor
Resolution (Å)	42.13 – 2.80 42.13 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.1 (42.13-2.80) 99.4 (42.13-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.15 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.224 , 0.266 0.222 , 0.263	Depositor DCC
R_{free} test set	3303 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	50.7	Xtrriage
Anisotropy	0.063	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	18502	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMP, C64, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2243	0.82	2/3025 (0.1%)
1	B	0.44	0/2356	0.80	1/3176 (0.0%)
1	C	0.45	0/2279	0.82	0/3072
1	D	0.44	0/2345	0.81	2/3165 (0.1%)
1	E	0.46	0/2181	0.83	2/2941 (0.1%)
1	F	0.46	0/2223	0.83	1/3004 (0.0%)
1	G	0.45	0/2209	0.80	1/2980 (0.0%)
1	H	0.46	0/2380	0.81	0/3216
All	All	0.45	0/18216	0.82	9/24579 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	231	GLN	N-CA-C	6.29	117.81	111.07
1	E	53	VAL	N-CA-C	5.50	115.96	111.62
1	D	231	GLN	N-CA-C	5.32	116.76	111.07
1	B	231	GLN	N-CA-C	5.23	116.98	111.28
1	A	54	THR	N-CA-C	5.21	117.14	108.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2221	0	2315	17	0
1	B	2330	0	2417	23	0
1	C	2255	0	2349	21	0
1	D	2322	0	2382	20	0
1	E	2162	0	2252	9	0
1	F	2198	0	2262	24	0
1	G	2188	0	2241	20	0
1	H	2355	0	2380	20	0
2	A	23	0	11	0	0
2	B	23	0	11	0	0
2	C	23	0	11	0	0
2	D	23	0	11	0	0
2	E	23	0	11	0	0
2	F	23	0	11	0	0
2	G	23	0	11	0	0
2	H	23	0	11	0	0
3	B	25	0	13	2	0
3	D	25	0	13	0	0
3	H	25	0	13	0	0
4	D	4	0	3	0	0
5	A	37	0	0	1	0
5	B	38	0	0	0	0
5	C	41	0	0	0	0
5	D	40	0	0	0	0
5	E	13	0	0	0	0
5	F	17	0	0	0	0
5	G	9	0	0	0	0
5	H	13	0	0	0	0
All	All	18502	0	18728	143	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 143 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:231:GLN:HE21	1:C:231:GLN:HA	1.45	0.82
1:C:50:MET:HB2	1:C:53:VAL:HG12	1.64	0.79
1:D:143:VAL:HG11	1:D:171:ASN:HB3	1.66	0.77
1:C:77:MET:HE1	1:C:147:GLU:HG3	1.67	0.76
1:C:242:ALA:HB3	1:C:249:ILE:HD11	1.67	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/361 (82%)	288 (97%)	10 (3%)	0	100	100
1	B	315/361 (87%)	304 (96%)	10 (3%)	1 (0%)	36	66
1	C	303/361 (84%)	290 (96%)	13 (4%)	0	100	100
1	D	317/361 (88%)	307 (97%)	10 (3%)	0	100	100
1	E	292/361 (81%)	278 (95%)	14 (5%)	0	100	100
1	F	297/361 (82%)	285 (96%)	11 (4%)	1 (0%)	36	66
1	G	294/361 (81%)	279 (95%)	12 (4%)	3 (1%)	12	38
1	H	324/361 (90%)	300 (93%)	23 (7%)	1 (0%)	36	66
All	All	2440/2888 (84%)	2331 (96%)	103 (4%)	6 (0%)	43	72

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	91	GLY
1	H	91	GLY
1	G	212	GLY
1	G	268	GLY
1	B	275	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/295 (82%)	231 (95%)	12 (5%)	22	55
1	B	253/295 (86%)	244 (96%)	9 (4%)	31	66
1	C	246/295 (83%)	229 (93%)	17 (7%)	14	41
1	D	248/295 (84%)	236 (95%)	12 (5%)	23	56
1	E	233/295 (79%)	224 (96%)	9 (4%)	28	64
1	F	239/295 (81%)	230 (96%)	9 (4%)	29	64
1	G	234/295 (79%)	215 (92%)	19 (8%)	11	33
1	H	247/295 (84%)	235 (95%)	12 (5%)	22	55
All	All	1943/2360 (82%)	1844 (95%)	99 (5%)	21	54

5 of 99 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	277	ILE
1	G	39	VAL
1	F	27	ARG
1	F	341	MET
1	G	137	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	204	ASN
1	G	231	GLN
1	F	82	ASN
1	H	168	HIS
1	G	20	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IMP	F	808	-	25,25,25	1.25	1 (4%)	37,38,38	2.11	9 (24%)
2	IMP	G	807	-	25,25,25	1.23	1 (4%)	37,38,38	2.17	10 (27%)
2	IMP	H	806	-	25,25,25	1.22	1 (4%)	37,38,38	2.05	9 (24%)
2	IMP	B	804	-	25,25,25	1.23	1 (4%)	37,38,38	2.07	9 (24%)
4	ACT	D	401	-	3,3,3	0.82	0	3,3,3	1.34	0
3	C64	H	903	-	27,28,28	1.65	6 (22%)	37,39,39	1.30	5 (13%)
2	IMP	E	805	-	25,25,25	1.25	1 (4%)	37,38,38	2.12	9 (24%)
2	IMP	C	803	-	25,25,25	1.23	1 (4%)	37,38,38	2.13	10 (27%)
2	IMP	D	802	-	25,25,25	1.24	1 (4%)	37,38,38	2.06	9 (24%)
3	C64	D	902	-	27,28,28	1.64	6 (22%)	37,39,39	1.33	5 (13%)
3	C64	B	901	-	27,28,28	1.64	6 (22%)	37,39,39	1.29	5 (13%)
2	IMP	A	801	-	25,25,25	1.26	1 (4%)	37,38,38	2.10	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMP	F	808	-	-	1/10/26/26	0/3/3/3
2	IMP	G	807	-	-	3/10/26/26	0/3/3/3
2	IMP	H	806	-	-	0/10/26/26	0/3/3/3
2	IMP	B	804	-	-	0/10/26/26	0/3/3/3
3	C64	H	903	-	-	1/12/12/12	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMP	E	805	-	-	3/10/26/26	0/3/3/3
2	IMP	C	803	-	-	0/10/26/26	0/3/3/3
2	IMP	D	802	-	-	0/10/26/26	0/3/3/3
3	C64	D	902	-	-	1/12/12/12	0/4/4/4
3	C64	B	901	-	-	1/12/12/12	0/4/4/4
2	IMP	A	801	-	-	0/10/26/26	0/3/3/3

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	903	C64	C10-C11	5.75	1.49	1.40
3	D	902	C64	C10-C11	5.74	1.49	1.40
3	B	901	C64	C10-C11	5.66	1.49	1.40
2	A	801	IMP	C2-N3	4.83	1.39	1.30
2	F	808	IMP	C2-N3	4.82	1.39	1.30

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	803	IMP	C5-C6-N1	6.24	119.75	110.78
2	G	807	IMP	C5-C6-N1	6.22	119.72	110.78
2	A	801	IMP	C5-C6-N1	6.16	119.63	110.78
2	E	805	IMP	C5-C6-N1	6.03	119.45	110.78
2	F	808	IMP	C5-C6-N1	6.02	119.43	110.78

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	807	IMP	O4'-C4'-C5'-O5'
2	G	807	IMP	C3'-C4'-C5'-O5'
2	E	805	IMP	C5'-O5'-P-O1P
2	E	805	IMP	O4'-C4'-C5'-O5'
3	D	902	C64	N3-C6-C8-S



There are no ring outliers.


1 monomer is involved in 2 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	901	C64	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/361 (83%)	-0.04	6 (1%) 65 56	28, 38, 55, 68	0
1	B	318/361 (88%)	0.01	2 (0%) 85 80	22, 40, 59, 63	1 (0%)
1	C	306/361 (84%)	-0.06	8 (2%) 57 47	22, 37, 55, 64	1 (0%)
1	D	321/361 (88%)	-0.03	6 (1%) 66 57	28, 38, 58, 70	0
1	E	298/361 (82%)	0.49	12 (4%) 42 33	50, 64, 86, 88	0
1	F	300/361 (83%)	0.37	3 (1%) 79 72	39, 62, 76, 77	1 (0%)
1	G	300/361 (83%)	0.64	12 (4%) 42 33	57, 73, 93, 94	0
1	H	328/361 (90%)	0.63	11 (3%) 48 39	52, 66, 84, 85	0
All	All	2473/2888 (85%)	0.25	60 (2%) 59 49	22, 54, 83, 94	3 (0%)

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	379	SER	5.2
1	F	218	ILE	4.8
1	C	331	ILE	4.8
1	G	379	SER	4.5
1	C	379	SER	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	D	401	4/4	0.67	0.28	95,95,95,95	0
2	IMP	G	807	23/23	0.76	0.14	104,105,105,105	0
2	IMP	E	805	23/23	0.80	0.14	79,81,82,82	0
3	C64	H	903	25/25	0.86	0.26	51,52,52,52	25
2	IMP	F	808	23/23	0.87	0.11	56,59,61,61	0
3	C64	D	902	25/25	0.87	0.26	50,50,50,51	25
2	IMP	A	801	23/23	0.88	0.12	57,61,62,62	0
3	C64	B	901	25/25	0.90	0.21	54,54,54,54	25
2	IMP	C	803	23/23	0.93	0.09	44,49,50,50	0
2	IMP	B	804	23/23	0.94	0.08	46,48,49,49	0
2	IMP	H	806	23/23	0.94	0.08	51,52,52,53	0
2	IMP	D	802	23/23	0.94	0.07	39,41,41,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IMP G 807:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IMP E 805:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C64 H 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IMP F 808:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C64 D 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IMP A 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C64 B 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IMP C 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IMP B 804:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IMP H 806:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.