



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 07:24 PM UTC

PDB ID : 7KID / pdb_00007kid
Title : PRMT5:MEP50 Complexed with 5,5-Bicyclic Inhibitor Compound 72
Authors : Palte, R.L.
Deposited on : 2020-10-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	14% 89% 8% .
2	B	350	49% 83% 6% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein arginine N-methyltransferase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	625	Total	C	N	O	S	0	3	0
			5086	3251	873	937	25			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP O14744
A	-6	ASP	-	expression tag	UNP O14744
A	-5	TYR	-	expression tag	UNP O14744
A	-4	LYS	-	expression tag	UNP O14744
A	-3	ASP	-	expression tag	UNP O14744
A	-2	ASP	-	expression tag	UNP O14744
A	-1	ASP	-	expression tag	UNP O14744
A	0	ASP	-	expression tag	UNP O14744
A	1	LYS	-	expression tag	UNP O14744

- Molecule 2 is a protein called Methylosome protein 50.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	311	Total	C	N	O	S	0	1	0
			2353	1475	403	460	15			

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	MET	-	initiating methionine	UNP Q9BQA1
B	-6	HIS	-	expression tag	UNP Q9BQA1
B	-5	HIS	-	expression tag	UNP Q9BQA1
B	-4	HIS	-	expression tag	UNP Q9BQA1
B	-3	HIS	-	expression tag	UNP Q9BQA1
B	-2	HIS	-	expression tag	UNP Q9BQA1
B	-1	HIS	-	expression tag	UNP Q9BQA1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	0	HIS	-	expression tag	UNP Q9BQA1
B	1	HIS	-	expression tag	UNP Q9BQA1

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is (1S,2R,3aR,4S,6aR)-4-[(2-amino-3,5-difluoroquinolin-7-yl)methyl]-2-(4-amino-5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydropentalene-1,6a(1H)-diol (CCD ID: WFV) (formula: C₂₄H₂₃F₃N₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			35	24	3	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	223	Total	O	0	0
			223	223		
5	B	37	Total	O	0	0
			37	37		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein arginine N-methyltransferase 5



- Molecule 2: Methylosome protein 50



S306	F307	L308	N309	H310	S311	L312	L313	T314	T315	V316	G317	W318	D319	H320	Q321	V322	V326	V327	F328	THR	GLU	PRO	LEU	PRO	ALA	PRO	GLY	PRO	ALA	SER	VAL	THR	GLU
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	103.68Å 139.18Å 178.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.79 – 2.50 34.79 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.2 (34.79-2.50) 98.2 (34.79-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.259 , 0.279 0.273 , 0.300	Depositor DCC
R_{free} test set	2257 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.891	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 32.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7750	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: WFFV, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.16	1/5228 (0.0%)	0.98	0/7111
2	B	1.22	4/2410 (0.2%)	0.95	0/3294
All	All	1.18	5/7638 (0.1%)	0.97	0/10405

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	264	SER	C-N	6.31	1.39	1.33
2	B	245	THR	CA-C	5.92	1.60	1.52
2	B	147	ILE	CA-C	5.90	1.60	1.52
2	B	268	VAL	CA-C	5.89	1.59	1.52
1	A	627	ASN	CA-C	5.53	1.59	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5086	0	4970	23	0
2	B	2353	0	2266	5	0
3	A	16	0	24	0	0
4	A	35	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	223	0	0	0	0
5	B	37	0	0	0	0
All	All	7750	0	7260	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:MET:HG2	1:A:389:TYR:HB2	1.88	0.55
1:A:152:TRP:HB3	1:A:205:ALA:HB2	1.89	0.54
1:A:199:SER:HB3	1:A:202:ILE:HD12	1.93	0.51
1:A:445:LEU:HD11	1:A:611:VAL:HB	1.92	0.51
1:A:405:GLN:HA	1:A:409:TRP:HB2	1.92	0.50
1:A:301:ALA:HB1	1:A:505:ARG:HG2	1.93	0.50
1:A:369:GLY:N	1:A:370:PRO:HD3	2.26	0.50
1:A:493:ASP:HB3	1:A:496:ALA:HB2	1.94	0.49
1:A:432:ILE:HG12	1:A:457:LEU:HD13	1.95	0.49
1:A:567:ILE:HG21	1:A:579:TRP:HB2	1.94	0.49
1:A:369:GLY:H	1:A:370:PRO:HD3	1.78	0.48
1:A:603:TRP:HB2	1:A:614:GLU:HB2	1.96	0.48
1:A:221:TRP:HB3	1:A:226:ILE:HD11	1.95	0.48
1:A:371:LEU:HD13	1:A:433:VAL:HG12	1.98	0.46
1:A:53:GLU:HA	1:A:102[B]:ASN:HD21	1.80	0.46
2:B:96:VAL:HB	2:B:104:GLU:HB2	1.99	0.45
1:A:324:TYR:CB	1:A:368:ARG:HD2	2.47	0.45
1:A:553:GLY:HA3	1:A:582:ILE:HG22	1.98	0.45
1:A:476:SER:HB2	1:A:512:LEU:HD21	1.99	0.43
1:A:465:PRO:HB3	1:A:559:LEU:HD23	2.00	0.43
1:A:441:ALA:HB2	1:A:555:PHE:HB2	1.99	0.43
2:B:184:LEU:HB3	2:B:218:LEU:HD13	1.99	0.43
2:B:99:ASP:HA	2:B:127:ILE:HG23	2.00	0.43
1:A:421:ARG:HD2	1:A:449:CYS:HA	2.00	0.43
2:B:27:MET:HE3	2:B:31:LEU:HD21	2.00	0.42
1:A:558:VAL:HA	1:A:564:THR:HG22	2.02	0.41
1:A:361:LEU:HD13	1:A:375:SER:HB3	2.01	0.41
2:B:284:ASP:HB3	2:B:290:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	626/645 (97%)	605 (97%)	20 (3%)	1 (0%)	43	63
2	B	310/350 (89%)	298 (96%)	11 (4%)	1 (0%)	36	55
All	All	936/995 (94%)	903 (96%)	31 (3%)	2 (0%)	43	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	147	ILE
1	A	441	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	561/570 (98%)	553 (99%)	8 (1%)	59	81
2	B	265/298 (89%)	258 (97%)	7 (3%)	40	68
All	All	826/868 (95%)	811 (98%)	15 (2%)	51	77

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
1	A	177	GLU
1	A	235	ILE
1	A	594	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	622	CYS
1	A	629	THR
1	A	634	THR
1	A	635	ILE
2	B	19	LEU
2	B	249	LEU
2	B	257	CYS
2	B	280	LEU
2	B	283	LEU
2	B	298	ASP
2	B	300	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN
1	A	79	ASN
1	A	128	GLN
1	A	263	GLN
1	A	270	ASN
1	A	271	HIS
1	A	282	GLN
1	A	313	GLN
1	A	381	GLN
1	A	394	ASN
1	A	396	ASN
1	A	508	ASN
2	B	22	ASN
2	B	69	ASN
2	B	78	GLN
2	B	178	HIS
2	B	256	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	WFV	A	705	-	36,40,40	0.75	1 (2%)	51,63,63	1.50	4 (7%)
3	EDO	A	703	-	3,3,3	0.47	0	2,2,2	0.41	0
3	EDO	A	704	-	3,3,3	0.47	0	2,2,2	0.37	0
3	EDO	A	702	-	3,3,3	0.55	0	2,2,2	0.34	0
3	EDO	A	701	-	3,3,3	0.50	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	WFV	A	705	-	-	4/8/38/38	0/6/6/6
3	EDO	A	703	-	-	0/1/1/1	-
3	EDO	A	704	-	-	0/1/1/1	-
3	EDO	A	702	-	-	0/1/1/1	-
3	EDO	A	701	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	705	WFV	C21-C22	3.12	1.42	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	705	WFV	C-C4-C3	7.43	107.36	102.59
4	A	705	WFV	C16-C17-C18	-5.77	121.07	124.15
4	A	705	WFV	C6-C5-N	-3.10	106.41	108.67
4	A	705	WFV	C12-C11-C1	2.03	106.41	102.70

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	705	WFV	C4-C3-N-C5
4	A	705	WFV	C2-C3-N-C8
4	A	705	WFV	C2-C3-N-C5
4	A	705	WFV	C4-C3-N-C8

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	625/645 (96%)	1.06	92 (14%) 6 4	17, 40, 72, 93	3 (0%)
2	B	311/350 (88%)	2.11	172 (55%) 0 0	26, 58, 79, 114	1 (0%)
All	All	936/995 (94%)	1.41	264 (28%) 1 1	17, 47, 75, 114	4 (0%)

All (264) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	PHE	5.9
2	B	210	ALA	5.6
2	B	19	LEU	5.5
1	A	296	ALA	5.4
2	B	248	VAL	4.8
1	A	176	THR	4.7
1	A	146	HIS	4.7
2	B	211	PRO	4.6
2	B	86	LEU	4.5
1	A	294	PRO	4.5
2	B	21	PRO	4.4
2	B	29	ARG	4.4
2	B	41	ALA	4.4
2	B	245	THR	4.1
2	B	20	PRO	4.0
2	B	308	LEU	4.0
2	B	94	ILE	4.0
2	B	45	GLY	3.9
1	A	281	LEU	3.8
2	B	114	LEU	3.8
2	B	95	LEU	3.8
1	A	529	MET	3.7
2	B	243	LYS	3.7
2	B	303	ALA	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	209	SER	3.6
2	B	108	LEU	3.6
1	A	526	ARG	3.6
2	B	23	ALA	3.6
2	B	290	LEU	3.6
1	A	292	PRO	3.6
2	B	31	LEU	3.5
2	B	25	ALA	3.5
2	B	270	PHE	3.5
2	B	106	TRP	3.5
1	A	243	PHE	3.4
2	B	28	GLU	3.4
2	B	30	GLN	3.4
1	A	16	SER	3.3
1	A	280	TYR	3.3
2	B	18	ASN	3.3
2	B	22	ASN	3.3
1	A	175	HIS	3.3
1	A	357	ASN	3.3
2	B	239	LEU	3.3
2	B	116	VAL	3.3
2	B	66	ALA	3.3
2	B	67	ALA	3.3
1	A	275	LYS	3.3
1	A	528	PRO	3.3
2	B	307	PRO	3.3
2	B	149	ILE	3.3
1	A	454	GLN	3.2
2	B	302	ASP	3.2
1	A	293	PRO	3.2
1	A	316[A]	MET	3.2
2	B	72	PHE	3.2
2	B	274	LEU	3.2
1	A	273	SER	3.2
2	B	103	VAL	3.2
2	B	32	GLU	3.1
2	B	43	LEU	3.1
2	B	36	TYR	3.1
2	B	315	THR	3.1
2	B	208	CYS	3.1
2	B	322	VAL	3.1
2	B	136	SER	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	87	THR	3.1
1	A	299	LEU	3.1
1	A	266	ILE	3.1
2	B	115	ILE	3.1
2	B	62	LYS	3.0
2	B	27	MET	3.0
2	B	298	ASP	3.0
2	B	319	ASP	3.0
1	A	145	GLY	3.0
2	B	83	VAL	3.0
1	A	456	PHE	3.0
2	B	73	CYS	3.0
2	B	306	SER	3.0
1	A	50	PHE	3.0
2	B	47	SER	3.0
1	A	356	THR	2.9
1	A	346	LEU	2.9
2	B	68	PRO	2.9
2	B	99	ASP	2.9
1	A	249	MET	2.9
2	B	76	GLY	2.9
1	A	147	HIS	2.9
2	B	58	LEU	2.9
2	B	60	LEU	2.9
2	B	313	LEU	2.9
1	A	255	PHE	2.9
2	B	135	SER	2.8
1	A	51	LYS	2.8
2	B	44	LEU	2.8
2	B	312	LEU	2.8
2	B	317	GLY	2.8
1	A	349	VAL	2.8
1	A	315	LEU	2.8
1	A	33	VAL	2.8
2	B	300	VAL	2.8
2	B	278	CYS	2.8
2	B	220	TRP	2.8
2	B	281	ALA	2.8
1	A	169	GLU	2.8
2	B	112	GLU	2.8
2	B	310	HIS	2.8
2	B	263	PHE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	24	PRO	2.7
2	B	181	SER	2.7
2	B	179	LYS	2.7
2	B	301	ARG	2.7
2	B	88	TRP	2.7
1	A	412	GLN	2.7
1	A	462	VAL	2.7
2	B	96	VAL	2.7
2	B	98	SER	2.7
2	B	225	SER	2.7
2	B	314	THR	2.7
1	A	272	HIS	2.7
2	B	266	HIS	2.7
1	A	383	ASP	2.6
2	B	126	ASP	2.6
2	B	246	SER	2.6
1	A	329	LYS	2.6
2	B	164	ARG	2.6
2	B	77	VAL	2.6
1	A	379	ALA	2.6
2	B	214	LEU	2.6
2	B	141	VAL	2.6
2	B	267	SER	2.6
2	B	39	ASP	2.6
2	B	49	LEU	2.6
2	B	223	GLN	2.6
2	B	268	VAL	2.6
1	A	271	HIS	2.6
2	B	244	SER	2.6
2	B	227	VAL	2.5
1	A	378	ALA	2.5
1	A	267	THR	2.5
2	B	265	PRO	2.5
1	A	361	LEU	2.5
2	B	271	LEU	2.5
2	B	287	LEU	2.5
2	B	113	THR	2.5
1	A	242	GLY	2.5
1	A	177	GLU	2.5
2	B	59	TRP	2.5
1	A	231	LEU	2.5
2	B	61	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	316	VAL	2.5
1	A	530	ILE	2.5
2	B	275	SER	2.5
2	B	311	SER	2.5
1	A	431	ILE	2.4
2	B	134	LEU	2.4
1	A	245	VAL	2.4
1	A	240	LYS	2.4
1	A	428	LYS	2.4
2	B	109	ASP	2.4
1	A	399	VAL	2.4
2	B	241	ASP	2.4
1	A	260	LEU	2.4
1	A	301	ALA	2.4
2	B	118	LYS	2.4
2	B	259	THR	2.4
2	B	82	GLY	2.4
2	B	178	HIS	2.4
1	A	287	LEU	2.3
2	B	42	LEU	2.3
2	B	256	GLN	2.3
2	B	63	ASP	2.3
2	B	318	TRP	2.3
1	A	637	LEU	2.3
2	B	180	ASP	2.3
1	A	300	PHE	2.3
2	B	26	CYS	2.3
1	A	286	TYR	2.3
2	B	240	VAL	2.3
2	B	282	VAL	2.3
2	B	165	ALA	2.3
1	A	368	ARG	2.3
2	B	218	LEU	2.3
2	B	230	PHE	2.3
1	A	353	GLU	2.3
1	A	350	PRO	2.3
2	B	222	PRO	2.3
2	B	132	SER	2.3
1	A	26	ILE	2.3
2	B	261	LEU	2.3
1	A	276	GLU	2.3
2	B	170	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	297	ARG	2.3
2	B	219	ALA	2.2
2	B	304	THR	2.2
1	A	369	GLY	2.2
2	B	101	GLY	2.2
1	A	58	PRO	2.2
2	B	328	PRO	2.2
2	B	78	GLN	2.2
2	B	320	HIS	2.2
1	A	458	LYS	2.2
1	A	235	ILE	2.2
2	B	92	ARG	2.2
2	B	326	VAL	2.2
1	A	289	GLN	2.2
1	A	256	ARG	2.2
1	A	404	TRP	2.2
2	B	128	VAL	2.2
2	B	224	GLN	2.2
2	B	242	THR	2.2
2	B	38	SER	2.2
2	B	291	PHE	2.2
1	A	413	VAL	2.1
2	B	89	VAL	2.1
1	A	210	ALA	2.1
2	B	84	ALA	2.1
2	B	193	LEU	2.1
2	B	207	GLY	2.1
1	A	283	TYR	2.1
2	B	228	PHE	2.1
1	A	336	GLN	2.1
1	A	59	ALA	2.1
2	B	55	ALA	2.1
1	A	284	LEU	2.1
2	B	206	ILE	2.1
1	A	385	ARG	2.1
2	B	182	VAL	2.1
2	B	321	GLN	2.1
1	A	409	TRP	2.1
2	B	203	ALA	2.1
1	A	270	ASN	2.1
1	A	72	LEU	2.1
2	B	130	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	212	GLY	2.1
1	A	291	ARG	2.1
1	A	274	GLU	2.1
2	B	121	LYS	2.1
2	B	53[A]	CYS	2.1
2	B	100	SER	2.1
2	B	144	SER	2.1
2	B	64	PRO	2.1
1	A	544	VAL	2.1
2	B	131	VAL	2.1
2	B	33	ALA	2.1
2	B	234	ASN	2.1
1	A	348	ARG	2.1
1	A	285	GLU	2.1
2	B	117	SER	2.1
2	B	204	SER	2.1
2	B	119	PHE	2.0
2	B	258	VAL	2.0
2	B	309	ASN	2.0
2	B	104	GLU	2.0
1	A	241	LYS	2.0
1	A	297	TYR	2.0
1	A	307	TYR	2.0
2	B	74	SER	2.0
1	A	358	VAL	2.0
2	B	133	VAL	2.0
2	B	173	VAL	2.0
2	B	75	ALA	2.0
2	B	54	TRP	2.0
1	A	244	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	703	4/4	0.78	0.20	48,48,48,48	0
3	EDO	A	701	4/4	0.80	0.16	40,40,41,41	0
3	EDO	A	704	4/4	0.80	0.18	57,57,57,57	0
3	EDO	A	702	4/4	0.88	0.16	36,36,36,36	0
4	WV	A	705	35/35	0.89	0.12	29,32,33,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.