



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:52 AM UTC

PDB ID : 5KJK / pdb_00005kjk
Title : SMYD2 in complex with AZ370
Authors : Ferguson, A.
Deposited on : 2016-06-20
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BU3	A	502	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3888 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-lysine methyltransferase SMYD2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	1	0
			3456	2192	590	643	31			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	165	GLU	GLY	engineered mutation	UNP Q9NRG4

- Molecule 2 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: $C_{15}H_{22}N_6O_5S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

- Molecule 3 is (R,R)-2,3-BUTANEDIOL (CCD ID: BU3) (formula: $C_4H_{10}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	4	2		
3	A	1	Total	C	O	0	0
			6	4	2		
3	A	1	Total	C	O	0	0
			6	4	2		
3	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 4 is 2-[2-[1-[2-(3,4-dichlorophenyl)ethyl]azetidin-3-yl]oxyphenyl]- {N}-(3-pyrrolidin-1-ylpropyl)pyridine-4-carboxamide (CCD ID: 6T1) (formula: C₃₀H₃₄Cl₂N₄O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			38	30	2	4	2		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Zn	0	0
			3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	340	Total	O	0	0
			340	340		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-lysine methyltransferase SMYD2



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	155.87Å 155.87Å 52.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	110.21 – 1.93 110.21 – 1.93	Depositor EDS
% Data completeness (in resolution range)	95.5 (110.21-1.93) 95.5 (110.21-1.93)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 1.92Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.178 , 0.203 (Not available) , 0.201	Depositor DCC
R_{free} test set	2296 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3888	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, ZN, 6T1, BU3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/3538 (0.0%)	1.24	7/4769 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	342	MET	SD-CE	-6.99	1.62	1.79

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	GLU	CB-CG-CD	5.67	122.24	112.60
1	A	189	GLU	CB-CG-CD	5.61	122.14	112.60
1	A	52	CYS	CA-C-N	5.43	127.56	120.28
1	A	52	CYS	C-N-CA	5.43	127.56	120.28
1	A	60	GLU	N-CA-C	-5.26	102.74	110.52
1	A	134	PHE	CA-CB-CG	5.22	119.02	113.80
1	A	56	PHE	CA-CB-CG	5.17	118.97	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3456	0	3396	34	0
2	A	27	0	22	0	0
3	A	24	0	40	7	0
4	A	38	0	0	1	0
5	A	3	0	0	0	0
6	A	340	0	0	1	0
All	All	3888	0	3458	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ARG:HH12	3:A:502:BU3:H42	1.12	1.14
1:A:255:ARG:HH12	3:A:502:BU3:C4	1.76	0.99
1:A:207:HIS:HD2	1:A:240:TYR:H	1.10	0.97
1:A:420:HIS:HD2	1:A:422:TYR:H	1.19	0.90
1:A:48:ARG:HD3	6:A:601:HOH:O	1.76	0.86
1:A:207:HIS:CD2	1:A:240:TYR:H	1.95	0.85
1:A:255:ARG:NH1	3:A:502:BU3:H42	1.93	0.81
1:A:269:THR:HG23	1:A:271:ASP:H	1.54	0.71
1:A:265:GLN:O	1:A:269:THR:HG22	1.97	0.63
1:A:207:HIS:HD2	1:A:240:TYR:N	1.91	0.62
1:A:356:TRP:CG	3:A:503:BU3:H12	2.35	0.62
1:A:420:HIS:CD2	1:A:422:TYR:H	2.10	0.61
1:A:212:ASN:HD21	1:A:237:PHE:H	1.47	0.61
1:A:248:GLU:HA	3:A:502:BU3:H2	1.85	0.58
1:A:318:LEU:HD11	1:A:353:MET:HE1	1.85	0.58
1:A:248:GLU:HG3	3:A:502:BU3:H41	1.87	0.56
1:A:212:ASN:ND2	1:A:237:PHE:H	2.04	0.55
1:A:158:HIS:HD2	1:A:159:PHE:CZ	2.27	0.52
1:A:340:LEU:HD21	1:A:369:PRO:HB2	1.91	0.52
1:A:209:CYS:HB2	1:A:264:CYS:SG	2.50	0.52
1:A:158:HIS:HD2	1:A:159:PHE:CE2	2.28	0.51
1:A:137:HIS:HD2	1:A:140:LYS:NZ	2.08	0.50
1:A:308:ALA:HB1	1:A:312:LYS:HG3	1.95	0.49
1:A:187:GLU:HB3	1:A:191:LEU:HA	1.97	0.47
1:A:54:TYR:HB2	1:A:82:ASP:HB2	1.95	0.47
1:A:288:ALA:O	1:A:292:ARG:HG2	2.15	0.46
1:A:248:GLU:CG	3:A:502:BU3:H41	2.47	0.43
1:A:350:CYS:HA	1:A:353:MET:HE2	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:PRO:HD3	1:A:337:VAL:HG21	2.02	0.42
1:A:350:CYS:HA	1:A:353:MET:CE	2.49	0.42
1:A:158:HIS:CD2	1:A:159:PHE:CZ	3.07	0.41
1:A:294:MET:HA	1:A:294:MET:HE3	2.03	0.41
1:A:207:HIS:HE1	1:A:209:CYS:SG	2.44	0.41
1:A:180:ASN:OD1	4:A:506:6T1:C22	2.69	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/429 (100%)	421 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/379 (100%)	358 (94%)	22 (6%)	18	6

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	35	SER
1	A	77	GLU
1	A	94	VAL
1	A	141	LEU
1	A	164	LEU
1	A	173	VAL
1	A	212	ASN
1	A	253	ARG
1	A	281	LYS
1	A	287	LYS
1	A	294	MET
1	A	303	GLU
1	A	306	ARG
1	A	319	GLU
1	A	340	LEU
1	A	370	TYR
1	A	372	LYS
1	A	388	LEU
1	A	428	GLN
1	A	431	GLU
1	A	433	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	79	GLN
1	A	137	HIS
1	A	150	GLN
1	A	157	HIS
1	A	158	HIS
1	A	169	ASN
1	A	207	HIS
1	A	212	ASN
1	A	397	HIS
1	A	420	HIS
1	A	428	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SAM	A	501	-	27,29,29	0.52	0	34,42,42	0.45	0
3	BU3	A	505	-	4,5,5	1.18	0	6,6,6	0.49	0
3	BU3	A	502	-	4,5,5	1.13	0	6,6,6	0.96	0
4	6T1	A	506	-	42,42,42	0.51	1 (2%)	49,57,57	0.38	0
3	BU3	A	504	-	4,5,5	1.13	0	6,6,6	0.48	0
3	BU3	A	503	-	4,5,5	1.15	0	6,6,6	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAM	A	501	-	-	3/17/33/33	0/3/3/3
3	BU3	A	505	-	-	4/4/4/4	-
3	BU3	A	502	-	-	4/4/4/4	-
4	6T1	A	506	-	-	5/22/39/39	0/5/5/5
3	BU3	A	504	-	-	0/4/4/4	-
3	BU3	A	503	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	506	6T1	C22-N4	-2.79	1.46	1.49

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	SAM	CB-CG-SD-CE
2	A	501	SAM	CB-CG-SD-C5'
3	A	502	BU3	O5-C2-C3-O6
3	A	502	BU3	C1-C2-C3-O6
3	A	502	BU3	O5-C2-C3-C4
3	A	505	BU3	O5-C2-C3-O6
3	A	505	BU3	C1-C2-C3-C4
4	A	506	6T1	C21-C20-O2-C4
3	A	505	BU3	C1-C2-C3-O6
3	A	505	BU3	O5-C2-C3-C4
4	A	506	6T1	C3-C4-O2-C20
4	A	506	6T1	C5-C4-O2-C20
3	A	502	BU3	C1-C2-C3-C4
4	A	506	6T1	N4-C23-C24-C25
2	A	501	SAM	C-CA-CB-CG
4	A	506	6T1	C22-C20-O2-C4


There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	BU3	6	0
4	A	506	6T1	1	0
3	A	503	BU3	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/429 (100%)	0.07	10 (2%) 61 68	19, 37, 65, 108	1 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	GLY	4.2
1	A	433	HIS	3.9
1	A	282	LEU	2.8
1	A	6	LEU	2.7
1	A	283	SER	2.5
1	A	291	ILE	2.3
1	A	284	ASP	2.3
1	A	301	VAL	2.3
1	A	292	ARG	2.2
1	A	365	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BU3	A	505	6/6	0.73	0.33	81,82,83,83	0
3	BU3	A	504	6/6	0.75	0.32	84,84,85,86	0
3	BU3	A	503	6/6	0.82	0.21	64,66,68,69	0
3	BU3	A	502	6/6	0.87	0.20	40,42,53,54	0
4	6T1	A	506	38/38	0.93	0.10	24,31,50,52	0
2	SAM	A	501	27/27	0.98	0.06	28,30,34,37	0
5	ZN	A	507	1/1	0.99	0.03	36,36,36,36	0
5	ZN	A	508	1/1	0.99	0.03	35,35,35,35	0
5	ZN	A	509	1/1	0.99	0.03	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 6T1 A 506:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SAM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.