



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:57 AM UTC

PDB ID : 3KK7 / pdb_00003kk7
Title : Crystal structure of Putative cell invasion protein with MAC/Perforin domain (NP_812351.1) from BACTERIOIDES THETA IOTAOMICRON VPI-5482 at 2.46 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2009-11-04
Resolution : 2.46 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	541	
1	B	541	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative cell invasion protein with MAC/Perforin domain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	502	Total	C	N	O	S	Se	0	4	0
			4034	2584	641	796	2	11			
1	B	499	Total	C	N	O	S	Se	0	5	0
			4012	2566	637	796	2	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8A267
B	0	GLY	-	expression tag	UNP Q8A267

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	114	Total 114	O 114	0	0
5	B	125	Total 125	O 125	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.40Å 127.20Å 138.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.39 – 2.46 49.39 – 2.46	Depositor EDS
% Data completeness (in resolution range)	97.2 (49.39-2.46) 97.6 (49.39-2.46)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.5.0053, PHENIX	Depositor
R, R_{free}	0.209 , 0.252 (Not available) , 0.246	Depositor DCC
R_{free} test set	2514 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 30.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	8306	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/4124	0.87	1/5566 (0.0%)
1	B	0.82	0/4105	0.88	1/5546 (0.0%)
All	All	0.81	0/8229	0.87	2/11112 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	496	MSE	N-CA-C	5.80	118.67	109.50
1	B	412	ASP	N-CA-C	-5.37	102.09	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4034	0	3931	44	0
1	B	4012	0	3889	41	0
2	A	1	0	0	0	0
3	A	8	0	12	0	0
3	B	4	0	6	0	0
4	B	8	0	14	0	0
5	A	114	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	125	0	0	4	0
All	All	8306	0	7852	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387[A]:GLU:OE1	1:B:551:GLN:NE2	2.07	0.86
1:B:289:LEU:HD23	1:B:337:LEU:HB2	1.61	0.81
1:A:384:GLN:HG3	1:A:484:LEU:HD23	1.67	0.75
1:A:414:VAL:HG11	1:A:426:ILE:HG23	1.68	0.74
1:A:387[B]:GLU:OE1	1:B:198[B]:ARG:NH1	2.23	0.70
1:B:412:ASP:OD2	1:B:429:ASN:ND2	2.25	0.68
1:A:478:PRO:HG2	1:A:487:GLN:HE21	1.58	0.68
1:A:293:THR:HG21	1:A:301:ILE:CD1	2.25	0.66
1:B:520:TYR:OH	1:B:536:VAL:HG11	1.98	0.62
1:B:166:HIS:C	1:B:167:ASN:HD22	2.08	0.62
1:A:275:PRO:O	1:A:276:THR:HG22	2.00	0.62
1:B:167:ASN:HD22	1:B:167:ASN:N	1.97	0.61
1:A:520:TYR:OH	1:A:536:VAL:HG11	2.00	0.61
1:A:547:THR:HG22	5:A:709:HOH:O	2.00	0.61
1:A:293:THR:HG21	1:A:301:ILE:HD12	1.82	0.61
1:A:450:LEU:HD23	1:A:454:ASN:ND2	2.16	0.60
1:B:404:SER:HB2	5:B:934:HOH:O	2.00	0.60
1:A:414:VAL:HG11	1:A:426:ILE:CG2	2.32	0.60
1:A:413:ILE:HD12	1:A:452:LYS:HD3	1.86	0.58
1:B:515:MSE:HE2	5:B:921:HOH:O	2.04	0.58
1:A:241:ALA:HB2	1:A:312:ILE:HG22	1.86	0.58
1:A:185:ASN:HB3	5:A:746:HOH:O	2.04	0.57
1:B:167:ASN:N	1:B:167:ASN:ND2	2.51	0.57
1:B:165:VAL:O	1:B:167:ASN:ND2	2.39	0.55
1:B:117:THR:HG23	1:B:184:VAL:HB	1.89	0.55
1:A:293:THR:HG21	1:A:301:ILE:HD11	1.91	0.51
1:B:443:ASN:OD1	1:B:452:LYS:NZ	2.44	0.51
1:A:175:VAL:HG13	1:A:260:MSE:HE2	1.93	0.51
1:A:276:THR:HG23	1:A:276:THR:O	2.12	0.49
1:B:375:ARG:HA	1:B:484:LEU:HD11	1.95	0.49
1:A:70:THR:HA	1:A:188:LEU:CD2	2.42	0.49
1:A:450:LEU:HD23	1:A:454:ASN:HD22	1.76	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115:LEU:O	1:B:116:ARG:HG2	2.12	0.49
1:B:501:ASN:HD22	1:B:546:MSE:SE	2.47	0.48
1:A:258:LYS:O	1:A:262:VAL:HG23	2.12	0.48
1:A:387[A]:GLU:OE2	1:B:198[A]:ARG:NH1	2.35	0.48
1:B:252:GLN:CG	1:B:254[B]:ASP:OD1	2.62	0.48
1:B:199:LYS:HE3	5:B:925:HOH:O	2.14	0.48
1:A:168:THR:O	1:A:168:THR:CG2	2.61	0.48
1:A:199:LYS:NZ	5:A:709:HOH:O	2.42	0.48
1:A:223:GLN:HG2	5:A:735:HOH:O	2.13	0.48
1:A:70:THR:HA	1:A:188:LEU:HD21	1.96	0.47
1:A:384:GLN:HG3	1:A:484:LEU:CD2	2.38	0.47
1:B:181:ILE:HG23	1:B:270:TRP:CZ3	2.50	0.47
1:A:501:ASN:HD22	1:A:546:MSE:SE	2.47	0.47
1:B:89:PHE:HA	1:B:352:MSE:HE2	1.96	0.47
1:B:211:ASP:OD1	1:B:215:ASN:ND2	2.45	0.47
1:A:168:THR:O	1:A:168:THR:HG22	2.15	0.46
1:A:457:LYS:O	1:A:461:SER:HB2	2.14	0.46
1:A:478:PRO:CG	1:A:487:GLN:HE21	2.24	0.46
1:B:459:GLU:OE1	1:B:462:LYS:NZ	2.44	0.46
1:A:378:ASP:OD2	1:A:484:LEU:HD22	2.15	0.46
1:A:415:PRO:HG2	1:A:427:PHE:HB2	1.98	0.46
1:B:45:ASN:OD1	1:B:45:ASN:C	2.59	0.45
1:A:478:PRO:HG2	1:A:487:GLN:NE2	2.27	0.45
1:B:221:LEU:C	1:B:221:LEU:HD23	2.42	0.45
1:B:36:THR:O	1:B:36:THR:CG2	2.66	0.43
1:A:547:THR:HA	1:A:550:TYR:CZ	2.53	0.43
1:A:491:VAL:HA	1:A:516:TYR:CE2	2.54	0.43
1:B:427:PHE:CD1	1:B:491:VAL:HG21	2.53	0.43
1:B:481:GLN:CD	1:B:483:THR:O	2.62	0.42
1:B:293:THR:HG21	1:B:298:SER:HA	2.01	0.42
1:B:293:THR:OG1	1:B:301:ILE:HD12	2.19	0.42
1:A:496:MSE:HE2	1:A:509:ILE:HG22	2.02	0.42
1:A:508:TYR:CZ	1:A:546:MSE:HE1	2.53	0.42
1:B:165:VAL:HG22	1:B:166:HIS:O	2.18	0.42
1:A:241:ALA:CB	1:A:312:ILE:HG22	2.49	0.42
1:B:85:ILE:HB	1:B:88:ASP:HB2	2.01	0.42
1:B:116:ARG:N	5:B:918:HOH:O	2.50	0.41
1:B:378:ASP:OD2	1:B:484:LEU:HD21	2.20	0.41
1:A:76:SER:HA	1:A:228:PHE:O	2.20	0.41
1:A:94:TYR:HB2	1:A:225:TYR:CE2	2.56	0.41
1:A:203:ASP:OD1	1:A:503:ASN:ND2	2.44	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:PHE:O	1:B:165:VAL:C	2.64	0.41
1:B:70:THR:HA	1:B:188:LEU:CD2	2.50	0.41
1:B:381:MSE:HE3	1:B:381:MSE:HB3	1.99	0.41
1:B:394:ILE:HG13	1:B:466:LEU:HD11	2.01	0.41
1:B:481:GLN:NE2	1:B:483:THR:O	2.54	0.41
1:B:293:THR:HG22	1:B:294:LYS:O	2.21	0.40
1:A:161:LYS:NZ	5:A:813:HOH:O	2.54	0.40
1:B:175:VAL:HG13	1:B:260:MSE:HE2	2.04	0.40
1:A:436:SER:O	1:A:437:ASP:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/541 (92%)	478 (96%)	20 (4%)	0	100	100
1	B	498/541 (92%)	479 (96%)	18 (4%)	1 (0%)	43	54
All	All	996/1082 (92%)	957 (96%)	38 (4%)	1 (0%)	48	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	169	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/473 (94%)	436 (98%)	10 (2%)	45	60
1	B	443/473 (94%)	433 (98%)	10 (2%)	44	59
All	All	889/946 (94%)	869 (98%)	20 (2%)	44	60

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	GLU
1	A	167	ASN
1	A	276	THR
1	A	414	VAL
1	A	428	SER
1	A	448	THR
1	A	466	LEU
1	A	469	LYS
1	A	484	LEU
1	A	525	TYR
1	B	114	GLU
1	B	167	ASN
1	B	185	ASN
1	B	207	GLU
1	B	248	THR
1	B	291	ILE
1	B	296	GLU
1	B	298	SER
1	B	441	LYS
1	B	483	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	ASN
1	A	157	GLN
1	A	185	ASN
1	A	435	GLN
1	A	477	ASN
1	A	487	GLN
1	B	342	GLN
1	B	386	GLN
1	B	481	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	605	-	3,3,3	0.45	0	2,2,2	0.30	0
4	MPD	B	601	-	7,7,7	0.47	0	9,10,10	1.32	0
3	EDO	B	604	-	3,3,3	0.40	0	2,2,2	0.65	0
3	EDO	A	603	-	3,3,3	0.33	0	2,2,2	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	605	-	-	0/1/1/1	-
4	MPD	B	601	-	-	5/5/5/5	-
3	EDO	B	604	-	-	1/1/1/1	-
3	EDO	A	603	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	601	MPD	O2-C2-C3-C4
4	B	601	MPD	CM-C2-C3-C4
4	B	601	MPD	C2-C3-C4-O4
4	B	601	MPD	C2-C3-C4-C5
3	B	604	EDO	O1-C1-C2-O2
4	B	601	MPD	C1-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	491/541 (90%)	0.29	20 (4%)	41 41	14, 29, 48, 68	4 (0%)
1	B	488/541 (90%)	0.34	24 (4%)	35 33	13, 27, 42, 59	5 (1%)
All	All	979/1082 (90%)	0.32	44 (4%)	38 37	13, 28, 45, 68	9 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	THR	4.8
1	B	166	HIS	4.6
1	A	480	ILE	4.3
1	A	275	PRO	4.2
1	A	168	THR	4.1
1	A	56	THR	4.0
1	A	479	ILE	4.0
1	B	480	ILE	3.9
1	B	154	PHE	3.6
1	B	66	PHE	3.5
1	A	66	PHE	3.1
1	B	482	THR	3.0
1	A	287	GLY	3.0
1	A	481	GLN	2.8
1	B	295	ARG	2.8
1	A	67	THR	2.8
1	B	168	THR	2.8
1	B	478	PRO	2.7
1	A	274	LYS	2.7
1	A	429	ASN	2.6
1	B	477	ASN	2.6
1	A	476	ILE	2.6
1	A	450	LEU	2.5
1	B	56	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	333	TYR	2.5
1	B	342	GLN	2.5
1	A	124	SER	2.5
1	B	118	THR	2.5
1	B	36	THR	2.4
1	A	169	ASP	2.4
1	A	348[A]	LYS	2.4
1	B	165	VAL	2.3
1	B	479	ILE	2.3
1	B	67	THR	2.3
1	A	484	LEU	2.2
1	B	476	ILE	2.2
1	A	478	PRO	2.2
1	B	271	LYS	2.1
1	B	69	ALA	2.1
1	A	167	ASN	2.1
1	B	319	TYR	2.1
1	B	483	THR	2.0
1	B	169	ASP	2.0
1	B	254[A]	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	B	604	4/4	0.65	0.24	53,53,54,55	0
3	EDO	A	603	4/4	0.83	0.16	38,40,43,46	0
3	EDO	A	605	4/4	0.84	0.16	38,41,42,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MPD	B	601	8/8	0.89	0.14	38,41,42,42	0
2	CL	A	602	1/1	0.90	0.12	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.