



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:20 AM UTC

PDB ID : 3KKT / pdb_00003kkt
Title : Crystal structure of human PDE4b with 5-[3-[(1S,2S,4R)-Bicyclo[2.2.1]hept-2-yloxy]-4-methoxyp henyl]tetrahydro-2(1H)-pyrimidinone reveals ordering of the C-terminal helix residues 502-509.
Authors : Cheng, R.K.Y.; Crawley, L.; Barker, J.; Whittaker, M.
Deposited on : 2009-11-06
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	4% 74% 12% • 13%
1	B	398	3% 72% 11% • 15%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	S	4	0	0
			2809	1771	474	543	21			
1	B	338	Total	C	N	O	S	4	1	0
			2746	1734	464	527	21			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	MET	-	expression tag	UNP Q07343
A	132	GLY	-	expression tag	UNP Q07343
A	133	SER	-	expression tag	UNP Q07343
A	134	SER	-	expression tag	UNP Q07343
A	135	HIS	-	expression tag	UNP Q07343
A	136	HIS	-	expression tag	UNP Q07343
A	137	HIS	-	expression tag	UNP Q07343
A	138	HIS	-	expression tag	UNP Q07343
A	139	HIS	-	expression tag	UNP Q07343
A	140	HIS	-	expression tag	UNP Q07343
A	141	SER	-	expression tag	UNP Q07343
A	142	SER	-	expression tag	UNP Q07343
A	143	GLY	-	expression tag	UNP Q07343
A	144	LEU	-	expression tag	UNP Q07343
A	145	VAL	-	expression tag	UNP Q07343
A	146	PRO	-	expression tag	UNP Q07343
A	147	ARG	-	expression tag	UNP Q07343
A	148	GLY	-	expression tag	UNP Q07343
A	149	SER	-	expression tag	UNP Q07343
A	150	HIS	-	expression tag	UNP Q07343
A	151	MET	-	expression tag	UNP Q07343
B	131	MET	-	expression tag	UNP Q07343
B	132	GLY	-	expression tag	UNP Q07343
B	133	SER	-	expression tag	UNP Q07343
B	134	SER	-	expression tag	UNP Q07343

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	135	HIS	-	expression tag	UNP Q07343
B	136	HIS	-	expression tag	UNP Q07343
B	137	HIS	-	expression tag	UNP Q07343
B	138	HIS	-	expression tag	UNP Q07343
B	139	HIS	-	expression tag	UNP Q07343
B	140	HIS	-	expression tag	UNP Q07343
B	141	SER	-	expression tag	UNP Q07343
B	142	SER	-	expression tag	UNP Q07343
B	143	GLY	-	expression tag	UNP Q07343
B	144	LEU	-	expression tag	UNP Q07343
B	145	VAL	-	expression tag	UNP Q07343
B	146	PRO	-	expression tag	UNP Q07343
B	147	ARG	-	expression tag	UNP Q07343
B	148	GLY	-	expression tag	UNP Q07343
B	149	SER	-	expression tag	UNP Q07343
B	150	HIS	-	expression tag	UNP Q07343
B	151	MET	-	expression tag	UNP Q07343

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 5-{3-[(1S,2S,4R)-bicyclo[2.2.1]hept-2-yloxy]-4-methoxyphenyl}tetrahydropyrimidin-2(1H)-one (CCD ID: 0CP) (formula: C₁₈H₂₄N₂O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			23	18	2	3		
3	B	1	Total	C	N	O	0	0
			23	18	2	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: C₁₁H₂₆N₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			19	11	2	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	24	Total	O	0	0
			24	24		
6	B	36	Total	O	0	0
			36	36		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	89.74Å 89.74Å 337.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	70.59 – 2.48 70.59 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.8 (70.59-2.48) 99.8 (70.59-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.235 , 0.295 (Not available) , 0.277	Depositor DCC
R_{free} test set	1506 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5684	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B3P, ZN, OCP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2869	0.82	1/3890 (0.0%)
1	B	0.48	1/2803 (0.0%)	0.85	2/3793 (0.1%)
All	All	0.46	1/5672 (0.0%)	0.83	3/7683 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	328	LYS	CB-CG	10.04	1.82	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	328	LYS	CA-CB-CG	-12.24	89.62	114.10
1	B	328	LYS	CB-CG-CD	5.81	124.66	111.30
1	A	328	LYS	CA-CB-CG	-5.25	103.59	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2809	0	2720	29	0
1	B	2746	0	2662	30	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	23	0	24	1	0
3	B	23	0	24	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	19	0	26	0	0
6	A	24	0	0	0	0
6	B	36	0	0	0	0
All	All	5684	0	5456	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:MET:HA	1:A:221:MET:HE2	1.39	1.01
1:A:221:MET:HA	1:A:221:MET:CE	2.04	0.88
1:B:250:LEU:HD13	1:B:387:MET:HE1	1.60	0.81
1:A:193:THR:OG1	1:A:221:MET:HE3	1.87	0.74
1:A:190:ARG:CG	1:A:221:MET:HE1	2.20	0.71
1:B:159:ASN:H	1:B:159:ASN:HD22	1.41	0.66
1:A:327:THR:HG22	1:A:329:LYS:H	1.60	0.65
1:B:250:LEU:CD1	1:B:387:MET:HE1	2.27	0.63
1:A:246:THR:HG23	1:A:387:MET:HE3	1.81	0.63
1:A:180:ILE:HA	1:A:183:VAL:HG13	1.82	0.61
1:B:415:PHE:HA	1:B:431:MET:HE1	1.82	0.60
1:A:285:PHE:CE2	1:A:426:MET:HE1	2.36	0.60
1:A:507:GLN:O	1:A:508:PHE:HB2	2.00	0.60
1:A:348:SER:HA	1:A:509:GLU:HB3	1.85	0.58
1:A:172:ASP:HB2	1:A:178:LEU:HD12	1.89	0.55
1:B:313:LYS:O	1:B:316:GLN:HG2	2.07	0.54
1:B:418:GLY:HA3	1:B:431:MET:HE2	1.90	0.53
1:A:178:LEU:HD22	1:A:244:GLN:HG3	1.91	0.52
1:A:411:MET:HE3	1:A:439:VAL:HG13	1.90	0.52
1:B:221:MET:HA	1:B:221:MET:CE	2.39	0.52
1:A:395:ASN:HB2	1:A:396:PRO:HD3	1.93	0.51
1:B:410:ILE:O	1:B:414:PHE:HB2	2.11	0.51
1:A:382:GLN:HG3	1:A:385:ARG:HH21	1.75	0.51
1:B:191:PRO:HG3	1:B:237:LEU:HD12	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:ARG:HG3	1:A:221:MET:HE1	1.94	0.50
1:B:340:ASP:OD1	1:B:385:ARG:HD3	2.12	0.50
1:B:221:MET:HA	1:B:221:MET:HE2	1.94	0.50
1:A:333:THR:HG22	1:A:337:MET:HE2	1.95	0.49
1:B:382:GLN:HG3	1:B:385:ARG:NH2	2.27	0.48
1:B:159:ASN:HB2	1:B:162:ASN:ND2	2.29	0.48
1:A:190:ARG:HG2	1:A:221:MET:HE1	1.93	0.48
1:A:360:MET:HE1	1:A:383:VAL:HG22	1.96	0.47
1:B:156:PHE:HB2	1:B:158:VAL:HG23	1.97	0.47
1:B:397:THR:HB	1:B:469:ILE:HG23	1.98	0.46
1:A:421:GLU:HA	1:A:426:MET:HE3	1.98	0.46
1:B:159:ASN:HD22	1:B:159:ASN:N	2.10	0.45
1:A:507:GLN:HG2	1:A:508:PHE:H	1.82	0.45
1:B:395:ASN:HB2	1:B:396:PRO:HD3	1.99	0.45
1:A:258:VAL:HG11	1:A:374:LEU:HD12	1.99	0.45
1:B:426:MET:HE3	1:B:426:MET:HB2	1.89	0.44
1:B:333:THR:HG22	1:B:337:MET:HE2	2.00	0.44
1:B:440:GLU:H	1:B:440:GLU:CD	2.26	0.44
1:B:263:GLU:HG2	1:B:337:MET:SD	2.58	0.43
1:B:421:GLU:HA	1:B:426:MET:HE3	2.00	0.43
1:B:159:ASN:H	1:B:159:ASN:ND2	2.13	0.43
1:B:193:THR:OG1	1:B:221:MET:HE3	2.19	0.43
1:A:443:GLN:NE2	3:A:2:0CP:H9	2.34	0.42
1:A:463:GLN:HA	1:A:464:PRO:HA	1.66	0.42
1:A:180:ILE:HA	1:A:183:VAL:CG1	2.48	0.42
1:B:275:ASP:O	1:B:278:HIS:HB2	2.19	0.42
1:B:285:PHE:CE2	1:B:426:MET:HE1	2.54	0.42
1:B:159:ASN:N	1:B:159:ASN:ND2	2.68	0.41
1:B:154:SER:HB3	1:B:217:ILE:HG22	2.03	0.41
1:B:258:VAL:HG11	1:B:374:LEU:HD12	2.02	0.41
1:A:303:LEU:HD23	1:A:303:LEU:HA	1.98	0.41
1:A:374:LEU:HD22	1:A:379:ASP:HB3	2.03	0.41
1:A:242:VAL:HG12	1:A:271:ALA:HB1	2.03	0.41
1:A:275:ASP:O	1:A:278:HIS:HB2	2.20	0.40
1:B:463:GLN:HA	1:B:464:PRO:HA	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/398 (87%)	331 (96%)	12 (4%)	2 (1%)	21	35
1	B	333/398 (84%)	323 (97%)	9 (3%)	1 (0%)	36	53
All	All	678/796 (85%)	654 (96%)	21 (3%)	3 (0%)	30	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	508	PHE
1	B	321	ASP
1	A	488	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/360 (88%)	297 (94%)	19 (6%)	17	33
1	B	307/360 (85%)	287 (94%)	20 (6%)	15	30
All	All	623/720 (86%)	584 (94%)	39 (6%)	16	31

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	GLU
1	A	172	ASP
1	A	178	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	183	VAL
1	A	190	ARG
1	A	213	SER
1	A	221	MET
1	A	225	ASP
1	A	296	MET
1	A	313	LYS
1	A	328	LYS
1	A	334	LEU
1	A	356	ASP
1	A	373	LEU
1	A	422	ARG
1	A	432	CYS
1	A	436	THR
1	A	463	GLN
1	A	502	LEU
1	B	159	ASN
1	B	162	ASN
1	B	183	VAL
1	B	204	ASP
1	B	210	ARG
1	B	221	MET
1	B	225	ASP
1	B	237	LEU
1	B	250	LEU
1	B	313	LYS
1	B	316	GLN
1	B	319	HIS
1	B	334	LEU
1	B	366	VAL
1	B	371	VAL
1	B	372	LEU
1	B	440	GLU
1	B	499	CYS
1	B	502	LEU
1	B	507	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	332	GLN
1	A	435	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	443	GLN
1	A	452	HIS
1	A	463	GLN
1	B	159	ASN
1	B	162	ASN
1	B	288	ASN
1	B	316	GLN
1	B	382	GLN
1	B	481	GLN
1	B	500	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	B3P	B	529	-	18,18,18	0.72	0	23,23,23	0.91	1 (4%)
3	0CP	B	1	-	26,26,26	0.91	1 (3%)	29,37,37	0.99	3 (10%)
3	0CP	A	2	-	26,26,26	0.95	1 (3%)	29,37,37	1.05	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	B3P	B	529	-	-	19/28/28/28	-
3	0CP	B	1	-	-	2/10/37/37	0/5/4/4
3	0CP	A	2	-	-	2/10/37/37	0/5/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2	0CP	C3-C2	3.89	1.48	1.40
3	B	1	0CP	C3-C2	3.69	1.48	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2	0CP	C1-O1-C2	2.96	121.85	117.51
3	B	1	0CP	C1-O1-C2	2.59	121.32	117.51
5	B	529	B3P	C2-N2-C8	2.48	119.80	116.17
3	B	1	0CP	O1-C2-C3	2.47	118.75	115.40
3	A	2	0CP	C3-O2-C4	2.40	124.17	119.00
3	A	2	0CP	O1-C2-C3	2.37	118.62	115.40
3	B	1	0CP	C3-O2-C4	2.04	123.41	119.00

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	529	B3P	C5-C4-N1-C3
5	B	529	B3P	C7-C4-N1-C3
5	B	529	B3P	N1-C4-C6-O5
5	B	529	B3P	C5-C4-C6-O5
5	B	529	B3P	C7-C4-C6-O5
5	B	529	B3P	C9-C8-N2-C2
5	B	529	B3P	C10-C8-N2-C2
5	B	529	B3P	C11-C8-N2-C2
5	B	529	B3P	N2-C8-C9-O1
5	B	529	B3P	C10-C8-C9-O1
5	B	529	B3P	C11-C8-C9-O1
5	B	529	B3P	O2-C10-C8-N2

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
5	B	529	B3P	O2-C10-C8-C9
5	B	529	B3P	O2-C10-C8-C11
5	B	529	B3P	C3-C1-C2-N2
5	B	529	B3P	C1-C2-N2-C8
5	B	529	B3P	C2-C1-C3-N1
5	B	529	B3P	C1-C3-N1-C4
5	B	529	B3P	C6-C4-N1-C3
3	B	1	0CP	C13-C12-C15-C18
3	B	1	0CP	C11-C12-C15-C18
3	A	2	0CP	C11-C12-C15-C18
3	A	2	0CP	C13-C12-C15-C18


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	0CP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	347/398 (87%)	0.41	14 (4%)	42 38	23, 46, 65, 71	1 (0%)
1	B	338/398 (84%)	0.21	12 (3%)	46 42	21, 37, 63, 67	2 (0%)
All	All	685/796 (86%)	0.31	26 (3%)	44 40	21, 40, 63, 71	3 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	155[A]	ARG	4.7
1	B	434	LYS	3.6
1	B	154	SER	3.5
1	B	480	TYR	3.3
1	B	510	LEU	3.1
1	A	508	PHE	3.0
1	B	439	VAL	2.8
1	A	487	SER	2.7
1	A	429	SER	2.7
1	B	432	CYS	2.6
1	A	427	GLU	2.6
1	A	488	PRO	2.6
1	A	468	ASP	2.5
1	A	428	ILE	2.4
1	A	492	LEU	2.4
1	A	401	GLU	2.3
1	B	157	GLY	2.3
1	A	435	HIS	2.3
1	B	509	GLU	2.2
1	B	479	TRP	2.2
1	B	502	LEU	2.1
1	A	163	GLU	2.1
1	A	368	SER	2.1
1	A	430	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	162	ASN	2.1
1	A	422	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	A	529	1/1	0.82	0.10	70,70,70,70	0
3	0CP	B	1	23/23	0.93	0.10	37,38,39,39	0
5	B3P	B	529	19/19	0.93	0.10	28,29,30,30	19
3	0CP	A	2	23/23	0.95	0.08	34,35,36,36	0
4	MG	B	530	1/1	0.97	0.04	53,53,53,53	0
2	ZN	B	2	1/1	1.00	0.04	31,31,31,31	0
2	ZN	A	1	1/1	1.00	0.01	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 0CP B 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B3P B 529:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 0CP A 2:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.