



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:33 AM UTC

PDB ID : 7KKO / pdb_00007kko
Title : Structure of the catalytic domain of tankyrase 1 in complex with olaparib
Authors : Gajiwala, K.S.; Ryan, K.
Deposited on : 2020-10-27
Resolution : 1.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	2% 92% 6%
1	B	213	6% 92% 7%
1	C	213	21% 94%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly [ADP-ribose] polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	4	0
			1717	1079	313	312	13			
1	B	212	Total	C	N	O	S	0	6	0
			1746	1094	319	320	13			
1	C	210	Total	C	N	O	S	0	3	0
			1711	1076	312	311	12			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1102	GLY	-	expression tag	UNP Q59FX0
A	1103	SER	-	expression tag	UNP Q59FX0
B	1102	GLY	-	expression tag	UNP Q59FX0
B	1103	SER	-	expression tag	UNP Q59FX0
C	1102	GLY	-	expression tag	UNP Q59FX0
C	1103	SER	-	expression tag	UNP Q59FX0

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 4-(3-{[4-(cyclopropylcarbonyl)piperazin-1-yl]carbonyl}-4-fluorobenzyl)phthalazine-1(2H)-one (CCD ID: 09L) (formula: C₂₄H₂₃FN₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			32	24	1	4	3		
3	B	1	Total	C	F	N	O	0	0
			32	24	1	4	3		
3	C	1	Total	C	F	N	O	0	0
			32	24	1	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	198	Total	O	0	0
			198	198		
4	B	189	Total	O	0	0
			189	189		
4	C	142	Total	O	0	0
			142	142		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Poly [ADP-ribose] polymerase



- Molecule 1: Poly [ADP-ribose] polymerase



- Molecule 1: Poly [ADP-ribose] polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	79.29Å 90.41Å 199.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.12 – 1.56 57.12 – 1.56	Depositor EDS
% Data completeness (in resolution range)	80.8 (57.12-1.56) 80.8 (57.12-1.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.56Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.194 , 0.221 (Not available) , 0.236	Depositor DCC
R_{free} test set	4031 reflections (3.98%)	wwPDB-VP
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5802	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 09L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/1760	1.11	3/2366 (0.1%)
1	B	0.87	2/1789 (0.1%)	1.17	8/2405 (0.3%)
1	C	0.83	0/1754	1.18	3/2358 (0.1%)
All	All	0.85	2/5303 (0.0%)	1.15	14/7129 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1268	MET	SD-CE	5.96	1.94	1.79
1	B	1204	ILE	CA-C	5.72	1.60	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1115	ASP	CA-CB-CG	7.53	120.13	112.60
1	B	1204	ILE	CA-C-N	7.10	134.48	121.70
1	B	1204	ILE	C-N-CA	7.10	134.48	121.70
1	C	1115	ASP	CA-CB-CG	6.47	119.07	112.60
1	C	1208	PHE	CA-CB-CG	-6.35	107.45	113.80
1	B	1132	HIS	CA-C-N	6.34	129.64	120.38
1	B	1132	HIS	C-N-CA	6.34	129.64	120.38
1	B	1203	TYR	CA-C-N	5.88	132.55	121.97
1	B	1203	TYR	C-N-CA	5.88	132.55	121.97
1	B	1239	ASP	CA-CB-CG	5.82	118.42	112.60
1	A	1115	ASP	CA-CB-CG	5.67	118.28	112.60
1	C	1239	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	1239	ASP	CA-CB-CG	5.24	117.84	112.60
1	A	1268	MET	CB-CG-SD	-5.04	97.59	112.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1717	0	1648	4	0
1	B	1746	0	1670	6	0
1	C	1711	0	1644	4	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	32	0	22	0	0
3	B	32	0	22	5	0
3	C	32	0	22	0	0
4	A	198	0	0	1	0
4	B	189	0	0	0	0
4	C	142	0	0	0	0
All	All	5802	0	5028	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1228:ILE:HD11	3:B:1402:09L:H16	1.33	1.10
1:C:1207:MET:HG3	1:C:1265:THR:HG22	1.42	0.98
1:C:1207:MET:CG	1:C:1265:THR:HG22	2.17	0.74
1:B:1266:MET:HE2	1:B:1268:MET:SD	2.30	0.71
1:B:1228:ILE:CD1	3:B:1402:09L:H16	2.17	0.69
1:A:1261:LEU:HA	1:A:1279[B]:ILE:HG22	1.76	0.67
1:A:1158:LEU:HD11	1:A:1196:GLY:HA3	1.86	0.58
1:B:1261:LEU:HA	1:B:1279[B]:ILE:HG23	1.90	0.53
3:B:1402:09L:CAO	3:B:1402:09L:FAA	2.47	0.53
1:C:1128:THR:HB	1:C:1217:ASN:HA	1.93	0.51
1:C:1128:THR:HB	1:C:1217:ASN:HD22	1.79	0.48
3:B:1402:09L:CAU	3:B:1402:09L:H8	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1201:HIS:HE1	4:A:1660:HOH:O	1.99	0.46
1:A:1260:PHE:CG	1:A:1272:PRO:HG2	2.52	0.45
3:B:1402:09L:FAA	3:B:1402:09L:H8	2.09	0.41
1:B:1260:PHE:CG	1:B:1272:PRO:HG2	2.55	0.41
1:B:1279[B]:ILE:HD11	1:B:1281:ARG:CZ	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/213 (100%)	212 (100%)	0	0	100	100
1	B	216/213 (101%)	209 (97%)	5 (2%)	2 (1%)	14	3
1	C	211/213 (99%)	206 (98%)	5 (2%)	0	100	100
All	All	639/639 (100%)	627 (98%)	10 (2%)	2 (0%)	36	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1137	ASN
1	B	1204	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/180 (101%)	179 (98%)	3 (2%)	55	28
1	B	186/180 (103%)	183 (98%)	3 (2%)	55	28
1	C	181/180 (101%)	179 (99%)	2 (1%)	65	43
All	All	549/540 (102%)	541 (98%)	8 (2%)	57	31

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1110	ASP
1	A	1131	GLU
1	A	1314	GLU
1	B	1110	ASP
1	B	1144	ARG
1	B	1264	SER
1	C	1238	LYS
1	C	1285	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1143	ASN
1	A	1201	HIS
1	B	1223	GLN
1	B	1248	GLN
1	C	1143	ASN
1	C	1146	ASN
1	C	1177	HIS
1	C	1217	ASN
1	C	1237	HIS
1	C	1309	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	09L	B	1402	-	36,36,36	1.17	3 (8%)	49,52,52	1.56	10 (20%)
3	09L	C	1402	-	36,36,36	1.06	3 (8%)	49,52,52	1.47	12 (24%)
3	09L	A	1402	-	36,36,36	1.29	4 (11%)	49,52,52	1.28	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	09L	B	1402	-	-	10/20/32/32	0/5/5/5
3	09L	C	1402	-	-	4/20/32/32	0/5/5/5
3	09L	A	1402	-	-	5/20/32/32	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1402	09L	CAL-NAE	3.84	1.40	1.35
3	B	1402	09L	CAL-NAE	3.22	1.39	1.35
3	A	1402	09L	CBC-NAH	3.10	1.40	1.35
3	B	1402	09L	CAQ-NAF	2.35	1.39	1.34
3	C	1402	09L	CAL-NAE	2.33	1.38	1.35
3	A	1402	09L	CAQ-NAF	2.22	1.39	1.34
3	B	1402	09L	CAR-CAU	2.18	1.41	1.38
3	A	1402	09L	CAI-CAL	-2.08	1.48	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1402	09L	CAQ-NAF	2.08	1.39	1.34
3	C	1402	09L	CAI-CAL	-2.07	1.48	1.51

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1402	09L	CAK-CAI-CAL	4.76	123.01	116.91
3	B	1402	09L	CAJ-CAI-CAL	4.17	122.25	116.91
3	C	1402	09L	CAX-CAU-CAR	-3.39	119.42	123.07
3	B	1402	09L	CAI-CAL-NAE	-3.31	115.69	118.80
3	C	1402	09L	CAS-CAR-CAU	3.03	120.60	116.83
3	C	1402	09L	CAI-CAL-NAE	-2.87	116.10	118.80
3	B	1402	09L	CAS-CAR-CAU	2.78	120.28	116.83
3	C	1402	09L	CAZ-CAY-NAG	2.68	124.46	122.76
3	A	1402	09L	CAI-CAL-NAE	-2.66	116.30	118.80
3	A	1402	09L	CAJ-CAI-CAL	2.64	120.30	116.91
3	C	1402	09L	CAK-CAI-CAL	2.63	120.29	116.91
3	C	1402	09L	OAC-CAQ-CAR	-2.48	115.19	120.06
3	B	1402	09L	CAX-CAU-CAR	-2.44	120.44	123.07
3	C	1402	09L	CAP-NAF-CAQ	-2.42	115.30	122.79
3	A	1402	09L	OAB-CAL-NAE	2.37	124.44	121.61
3	B	1402	09L	CAV-CAT-CAS	2.37	123.98	120.38
3	C	1402	09L	CBB-CAZ-CAY	2.31	124.39	121.20
3	A	1402	09L	CAK-CAI-CAL	2.30	119.86	116.91
3	A	1402	09L	CAS-CAR-CAU	2.25	119.63	116.83
3	B	1402	09L	OAB-CAL-NAE	2.23	124.27	121.61
3	C	1402	09L	CBC-NAH-NAG	-2.21	125.51	127.98
3	B	1402	09L	CBB-CAZ-CAY	2.20	124.24	121.20
3	B	1402	09L	CAV-CAT-CAW	-2.20	117.66	120.89
3	A	1402	09L	CAX-CAU-CAR	-2.17	120.72	123.07
3	C	1402	09L	CAP-NAF-CAO	2.12	117.02	112.68
3	A	1402	09L	CBC-NAH-NAG	-2.05	125.69	127.98
3	B	1402	09L	OAD-CBC-NAH	2.04	123.49	120.88
3	C	1402	09L	CAJ-CAI-CAL	2.02	119.51	116.91
3	C	1402	09L	CAR-CAQ-NAF	2.00	121.35	118.24

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1402	09L	NAF-CAQ-CAR-CAU
3	B	1402	09L	OAC-CAQ-CAR-CAU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	1402	09L	CAT-CAV-CAY-CAZ
3	B	1402	09L	OAC-CAQ-CAR-CAS
3	B	1402	09L	NAF-CAQ-CAR-CAS
3	A	1402	09L	OAC-CAQ-CAR-CAS
3	B	1402	09L	OAC-CAQ-NAF-CAO
3	A	1402	09L	NAF-CAQ-CAR-CAS
3	C	1402	09L	NAF-CAQ-CAR-CAS
3	B	1402	09L	CAJ-CAI-CAL-OAB
3	B	1402	09L	CAJ-CAI-CAL-NAE
3	A	1402	09L	CAT-CAV-CAY-CAZ
3	C	1402	09L	CAT-CAV-CAY-CAZ
3	A	1402	09L	CAT-CAV-CAY-NAG
3	B	1402	09L	CAT-CAV-CAY-NAG
3	C	1402	09L	CAT-CAV-CAY-NAG
3	C	1402	09L	OAC-CAQ-CAR-CAS
3	A	1402	09L	OAC-CAQ-CAR-CAU
3	B	1402	09L	OAC-CAQ-NAF-CAP


There are no ring outliers.

1 monomer is involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1402	09L	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 09L B 1402



Ligand 09L C 1402





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/213 (98%)	0.37	5 (2%) 59 67	9, 24, 42, 103	4 (1%)
1	B	212/213 (99%)	0.47	13 (6%) 27 32	9, 24, 45, 95	6 (2%)
1	C	210/213 (98%)	0.97	45 (21%) 2 3	10, 29, 60, 115	3 (1%)
All	All	632/639 (98%)	0.60	63 (9%) 12 14	9, 26, 51, 115	13 (2%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1204	ILE	7.1
1	B	1203	TYR	6.2
1	C	1131	GLU	5.1
1	C	1132	HIS	4.9
1	B	1135	GLY	4.8
1	B	1133	ARG	4.2
1	C	1142	PHE	3.9
1	B	1136	GLY	3.8
1	C	1244	ILE	3.8
1	C	1228	ILE	3.7
1	C	1314	GLU	3.7
1	C	1139	GLY	3.7
1	C	1115	ASP	3.6
1	C	1243	TYR	3.5
1	C	1133	ARG	3.5
1	C	1137	ASN	3.4
1	C	1141	ILE	3.3
1	C	1312	LYS	3.3
1	C	1310	ILE	3.2
1	C	1136	GLY	3.2
1	B	1202	ALA	3.2
1	C	1112	ALA	3.0
1	C	1236	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1314	GLU	3.0
1	C	1233	GLY	3.0
1	C	1235	PRO	2.9
1	A	1105	GLY	2.9
1	C	1135	GLY	2.9
1	C	1234	CYS	2.9
1	C	1140	GLY	2.8
1	C	1129	ILE	2.8
1	C	1229	GLY	2.8
1	C	1237	HIS	2.8
1	C	1232	THR	2.7
1	C	1241	SER	2.7
1	A	1314	GLU	2.7
1	C	1148	ILE	2.7
1	C	1138	ALA	2.7
1	C	1313	PRO	2.7
1	B	1205	GLY	2.6
1	C	1284	VAL	2.6
1	C	1285	ASN	2.6
1	C	1242	CYS	2.6
1	B	1115	ASP	2.6
1	B	1137	ASN	2.5
1	C	1240	ARG	2.5
1	A	1157	LYS	2.4
1	B	1144	ARG	2.3
1	C	1143	ASN	2.3
1	A	1167	LYS	2.3
1	B	1189	ILE	2.3
1	C	1180	ARG	2.2
1	A	1174	HIS	2.2
1	C	1113	PRO	2.2
1	C	1230	GLY	2.2
1	C	1167	LYS	2.2
1	C	1238	LYS	2.2
1	C	1190	ASN	2.1
1	C	1134	ASP	2.1
1	C	1216	GLU	2.1
1	C	1265	THR	2.0
1	C	1226	TYR	2.0
1	B	1131	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	09L	B	1402	32/32	0.91	0.12	16,28,43,44	0
2	ZN	C	1401	1/1	0.94	0.07	47,47,47,47	0
3	09L	C	1402	32/32	0.95	0.06	17,21,24,26	0
3	09L	A	1402	32/32	0.97	0.05	15,18,22,24	0
2	ZN	A	1401	1/1	0.98	0.04	32,32,32,32	0
2	ZN	B	1401	1/1	0.98	0.04	32,32,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around 09L B 1402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 09L C 1402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.