



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:34 PM UTC

PDB ID : 7KKQ / pdb_00007kkq
Title : Structure of the catalytic domain of tankyrase 1 in complex with veliparib
Authors : Gajiwala, K.S.; Ryan, K.
Deposited on : 2020-10-27
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	10% 93% 5% .
1	B	213	15% 90% 5% 5%
1	C	213	10% 87% 9% .
1	D	213	10% 90% 5% . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly [ADP-ribose] polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	6	0
			1734	1089	321	310	14			
1	B	203	Total	C	N	O	S	0	3	0
			1664	1052	302	297	13			
1	C	206	Total	C	N	O	S	0	2	0
			1682	1062	307	300	13			
1	D	206	Total	C	N	O	S	0	4	0
			1697	1070	308	306	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1102	GLY	-	expression tag	UNP Q59FX0
A	1103	SER	-	expression tag	UNP Q59FX0
B	1102	GLY	-	expression tag	UNP Q59FX0
B	1103	SER	-	expression tag	UNP Q59FX0
C	1102	GLY	-	expression tag	UNP Q59FX0
C	1103	SER	-	expression tag	UNP Q59FX0
D	1102	GLY	-	expression tag	UNP Q59FX0
D	1103	SER	-	expression tag	UNP Q59FX0

- Molecule 2 is (2R)-2-(7-carbamoyl-1H-benzimidazol-2-yl)-2-methylpyrrolidinium (CCD ID: 78P) (formula: C₁₃H₁₆N₄O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			18	13	4	1		
2	B	1	Total	C	N	O	0	0
			18	13	4	1		
2	C	1	Total	C	N	O	0	0
			18	13	4	1		
2	D	1	Total	C	N	O	0	0
			18	13	4	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	218	Total	O	0	0
			218	218		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	200	Total 200	O 200	0	0
4	C	197	Total 197	O 197	0	0
4	D	218	Total 218	O 218	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Poly [ADP-ribose] polymerase




- Molecule 1: Poly [ADP-ribose] polymerase



- Molecule 1: Poly [ADP-ribose] polymerase



- Molecule 1: Poly [ADP-ribose] polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	76.19Å 159.10Å 84.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	79.55 – 1.59 79.55 – 1.59	Depositor EDS
% Data completeness (in resolution range)	79.5 (79.55-1.59) 79.9 (79.55-1.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.58Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.204 , 0.230 0.197 , 0.218	Depositor DCC
R_{free} test set	5532 reflections (3.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7686	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.3081e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 78P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/1777	1.08	1/2387 (0.0%)
1	B	0.83	1/1705 (0.1%)	1.10	2/2290 (0.1%)
1	C	0.82	0/1724	1.12	3/2316 (0.1%)
1	D	0.83	1/1739 (0.1%)	1.15	3/2336 (0.1%)
All	All	0.82	2/6945 (0.0%)	1.11	9/9329 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1228	ILE	CA-C	6.20	1.60	1.52
1	D	1284	VAL	CA-C	5.21	1.59	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1284	VAL	CA-C-N	12.46	144.13	121.70
1	D	1284	VAL	C-N-CA	12.46	144.13	121.70
1	C	1227	GLY	CA-C-N	5.84	128.76	120.53
1	C	1227	GLY	C-N-CA	5.84	128.76	120.53
1	D	1284	VAL	CA-C-O	-5.33	114.12	120.78
1	C	1115	ASP	CA-CB-CG	5.19	117.79	112.60
1	A	1134	ASP	CA-CB-CG	5.18	117.78	112.60
1	B	1285	ASN	CA-C-N	5.02	130.73	121.70
1	B	1285	ASN	C-N-CA	5.02	130.73	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1734	0	1671	5	0
1	B	1664	0	1613	4	0
1	C	1682	0	1627	8	0
1	D	1697	0	1634	10	0
2	A	18	0	16	1	0
2	B	18	0	16	1	0
2	C	18	0	16	3	0
2	D	18	0	16	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	218	0	0	0	0
4	B	200	0	0	0	0
4	C	197	0	0	1	0
4	D	218	0	0	0	0
All	All	7686	0	6609	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1266:MET:HE3	1:D:1268[A]:MET:HE1	1.38	1.03
1:D:1266:MET:CE	1:D:1268[A]:MET:HE1	2.13	0.78
1:D:1266:MET:HE3	1:D:1268[A]:MET:CE	2.17	0.73
1:D:1262:GLN:HG3	1:D:1266:MET:HE2	1.72	0.71
1:D:1284:VAL:HB	1:D:1285:ASN:HB2	1.79	0.65
1:C:1268[B]:MET:HG2	4:C:9266:HOH:O	2.01	0.60
1:B:1227:GLY:C	1:B:1229:GLY:HA3	2.27	0.59
1:A:1112:ALA:HB3	1:A:1115:ASP:OD2	2.04	0.57
1:D:1262:GLN:HG3	1:D:1266:MET:CE	2.33	0.56
2:D:9001:78P:H111	2:D:9001:78P:H9	1.54	0.56
2:B:9001:78P:H111	2:B:9001:78P:H9	1.56	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1228:ILE:HG23	2:C:9001:78P:H151	1.88	0.54
2:C:9001:78P:H111	2:C:9001:78P:H9	1.58	0.51
2:A:9001:78P:H111	2:A:9001:78P:H9	1.59	0.51
1:D:1284:VAL:HB	1:D:1285:ASN:CB	2.41	0.51
1:A:1268[A]:MET:HE1	1:A:1278:VAL:HG21	1.94	0.50
1:A:1128:THR:HB	1:A:1217:ASN:HA	1.95	0.49
1:A:1266:MET:HE2	1:A:1268[B]:MET:SD	2.55	0.45
1:D:1259:SER:HB3	1:D:1279[B]:ILE:HG22	1.99	0.45
1:B:1128:THR:HB	1:B:1217:ASN:HA	1.98	0.45
1:D:1207:MET:HG2	1:D:1208:PHE:CE2	2.52	0.44
1:C:1137:ASN:HB3	1:C:1138:ALA:H	1.68	0.44
1:A:1239:ASP:HB3	1:A:1242:CYS:HB2	2.01	0.43
1:C:1224:TYR:CE1	2:C:9001:78P:H183	2.55	0.42
1:C:1239:ASP:HB3	1:C:1242:CYS:HB2	2.01	0.41
1:B:1268[B]:MET:SD	1:B:1272:PRO:HG3	2.60	0.41
1:C:1197:PHE:HB3	1:C:1212:ILE:HD13	2.01	0.41
1:B:1165:ARG:HG2	1:B:1298:GLU:HG2	2.03	0.40
1:C:1124:GLU:HG2	1:C:1251:PHE:CZ	2.56	0.40
1:C:1262:GLN:HE21	1:C:1263:PHE:H	1.69	0.40
1:D:1165:ARG:HB2	1:D:1298[A]:GLU:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/213 (100%)	209 (98%)	4 (2%)	0	100	100
1	B	202/213 (95%)	196 (97%)	6 (3%)	0	100	100
1	C	204/213 (96%)	202 (99%)	1 (0%)	1 (0%)	24	10
1	D	206/213 (97%)	201 (98%)	3 (2%)	2 (1%)	12	3
All	All	825/852 (97%)	808 (98%)	14 (2%)	3 (0%)	30	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	1284	VAL
1	D	1285	ASN
1	C	1284	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/180 (102%)	182 (100%)	1 (0%)	81	70
1	B	176/180 (98%)	175 (99%)	1 (1%)	78	66
1	C	178/180 (99%)	174 (98%)	4 (2%)	45	22
1	D	180/180 (100%)	175 (97%)	5 (3%)	38	15
All	All	717/720 (100%)	706 (98%)	11 (2%)	55	35

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1110	ASP
1	B	1110	ASP
1	C	1110	ASP
1	C	1116	LYS
1	C	1143	ASN
1	C	1175	ASN
1	D	1110	ASP
1	D	1189	ILE
1	D	1207	MET
1	D	1241	SER
1	D	1284	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1151	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1177	HIS
1	A	1248	GLN
1	B	1151	GLN
1	B	1177	HIS
1	B	1190	ASN
1	B	1246	HIS
1	B	1248	GLN
1	B	1262	GLN
1	C	1146	ASN
1	C	1248	GLN
1	C	1262	GLN
1	C	1309	GLN
1	D	1146	ASN
1	D	1166	GLN
1	D	1248	GLN
1	D	1309	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	78P	C	9001	-	19,20,20	1.18	3 (15%)	20,30,30	2.33	5 (25%)
2	78P	D	9001	-	19,20,20	1.33	3 (15%)	20,30,30	2.75	5 (25%)
2	78P	A	9001	-	19,20,20	1.30	3 (15%)	20,30,30	2.17	3 (15%)
2	78P	B	9001	-	19,20,20	1.40	3 (15%)	20,30,30	2.27	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	78P	C	9001	-	-	0/4/19/19	0/3/3/3
2	78P	D	9001	-	-	0/4/19/19	0/3/3/3
2	78P	A	9001	-	-	0/4/19/19	0/3/3/3
2	78P	B	9001	-	-	0/4/19/19	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	9001	78P	C8-N9	-3.67	1.31	1.35
2	D	9001	78P	C8-N9	-3.33	1.31	1.35
2	A	9001	78P	C8-N9	-3.28	1.31	1.35
2	D	9001	78P	C8-N7	2.97	1.37	1.32
2	B	9001	78P	C8-N7	2.81	1.37	1.32
2	C	9001	78P	C8-N9	-2.71	1.32	1.35
2	C	9001	78P	C8-N7	2.69	1.37	1.32
2	B	9001	78P	C10-N11	2.67	1.37	1.33
2	D	9001	78P	C10-N11	2.65	1.37	1.33
2	A	9001	78P	C10-N11	2.60	1.37	1.33
2	A	9001	78P	C8-N7	2.52	1.37	1.32
2	C	9001	78P	C10-N11	2.10	1.36	1.33

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	9001	78P	C4-C3-N7	-8.16	104.33	109.91
2	B	9001	78P	C4-C3-N7	-7.04	105.10	109.91
2	C	9001	78P	C4-C3-N7	-6.84	105.23	109.91
2	A	9001	78P	C4-C3-N7	-6.70	105.33	109.91
2	D	9001	78P	C3-C4-N9	6.30	110.65	105.40
2	C	9001	78P	C3-C4-N9	5.36	109.86	105.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	9001	78P	C3-C4-N9	5.01	109.58	105.40
2	B	9001	78P	C3-C4-N9	4.83	109.42	105.40
2	D	9001	78P	C5-C4-C3	-3.78	117.85	122.52
2	D	9001	78P	C2-C3-C4	3.15	123.13	120.24
2	C	9001	78P	C2-C3-C4	2.98	122.97	120.24
2	D	9001	78P	C16-C17-C13	2.86	106.41	104.17
2	A	9001	78P	C2-C3-C4	2.85	122.86	120.24
2	B	9001	78P	C2-C3-C4	2.68	122.70	120.24
2	C	9001	78P	C5-C4-C3	-2.52	119.41	122.52
2	C	9001	78P	C16-C17-C13	2.48	106.12	104.17
2	B	9001	78P	C5-C4-C3	-2.28	119.70	122.52
2	B	9001	78P	N9-C8-N7	-2.05	111.73	114.05

There are no chirality outliers.



There are no torsion outliers.



There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	9001	78P	3	0
2	D	9001	78P	1	0
2	A	9001	78P	1	0
2	B	9001	78P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	209/213 (98%)	0.55	21 (10%)	12 12	9, 25, 43, 59	6 (2%)
1	B	203/213 (95%)	0.88	32 (15%)	5 4	11, 28, 50, 85	3 (1%)
1	C	206/213 (96%)	0.67	21 (10%)	12 12	11, 27, 48, 63	2 (0%)
1	D	206/213 (96%)	0.52	21 (10%)	12 12	10, 25, 46, 63	4 (1%)
All	All	824/852 (96%)	0.65	95 (11%)	9 9	9, 26, 48, 85	15 (1%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1228	ILE	11.1
1	D	1284	VAL	7.2
1	C	1284	VAL	6.3
1	B	1232	THR	5.6
1	C	1138	ALA	5.0
1	D	1228	ILE	4.7
1	B	1229	GLY	4.7
1	B	1230	GLY	4.7
1	D	1231	GLY	4.6
1	B	1131	GLU	4.5
1	B	1176	HIS	4.4
1	A	1137	ASN	4.4
1	A	1203	TYR	4.4
1	C	1228	ILE	4.4
1	B	1313	PRO	4.2
1	B	1107	ILE	4.1
1	C	1232	THR	4.0
1	D	1285	ASN	4.0
1	B	1284	VAL	4.0
1	D	1134	ASP	3.9
1	B	1289	TYR	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1137	ASN	3.8
1	A	1228	ILE	3.7
1	A	1132	HIS	3.6
1	B	1203	TYR	3.6
1	D	1232	THR	3.5
1	D	1138	ALA	3.5
1	D	1107	ILE	3.5
1	A	1107	ILE	3.4
1	C	1230	GLY	3.4
1	A	1133	ARG	3.4
1	B	1285	ASN	3.4
1	D	1230	GLY	3.3
1	B	1138	ALA	3.3
1	C	1240	ARG	3.3
1	C	1313	PRO	3.3
1	D	1203	TYR	3.2
1	A	1289	TYR	3.2
1	C	1231	GLY	3.1
1	A	1115	ASP	3.1
1	B	1105	GLY	3.0
1	D	1189	ILE	3.0
1	D	1313	PRO	3.0
1	B	1261	LEU	3.0
1	C	1287	LEU	2.9
1	A	1232	THR	2.9
1	B	1231	GLY	2.9
1	C	1105	GLY	2.9
1	A	1313	PRO	2.9
1	B	1115	ASP	2.9
1	B	1188	PHE	2.8
1	A	1284	VAL	2.8
1	B	1264	SER	2.7
1	A	1130	ARG	2.6
1	C	1238	LYS	2.6
1	C	1107	ILE	2.5
1	C	1188	PHE	2.5
1	C	1203	TYR	2.5
1	D	1175	ASN	2.5
1	B	1175	ASN	2.5
1	A	1285	ASN	2.5
1	B	1288	ALA	2.4
1	A	1230	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1163	CYS	2.4
1	D	1263	PHE	2.4
1	B	1235	PRO	2.3
1	D	1105	GLY	2.3
1	B	1263	PHE	2.3
1	C	1266	MET	2.3
1	A	1188	PHE	2.3
1	B	1286	GLY	2.3
1	D	1240	ARG	2.3
1	D	1204	ILE	2.3
1	A	1177	HIS	2.2
1	B	1227	GLY	2.2
1	A	1138	ALA	2.2
1	C	1283	SER	2.2
1	B	1177	HIS	2.2
1	B	1226	TYR	2.2
1	A	1205	GLY	2.2
1	C	1171	GLU	2.2
1	D	1235	PRO	2.1
1	A	1193	ILE	2.1
1	D	1229	GLY	2.1
1	C	1285	ASN	2.1
1	B	1180[A]	ARG	2.1
1	C	1286	GLY	2.1
1	B	1233	GLY	2.1
1	A	1134	ASP	2.1
1	A	1288	ALA	2.1
1	D	1236	THR	2.1
1	B	1204	ILE	2.1
1	C	1282	PRO	2.1
1	B	1130	ARG	2.0
1	D	1188	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	78P	D	9001	18/18	0.94	0.08	17,20,30,30	0
2	78P	C	9001	18/18	0.96	0.07	18,21,31,32	0
2	78P	B	9001	18/18	0.97	0.06	19,21,34,35	0
2	78P	A	9001	18/18	0.98	0.05	16,19,28,30	0
3	ZN	A	9002	1/1	0.98	0.04	30,30,30,30	0
3	ZN	B	9002	1/1	0.98	0.05	35,35,35,35	0
3	ZN	C	9002	1/1	0.99	0.03	23,23,23,23	0
3	ZN	D	9002	1/1	0.99	0.03	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around 78P C 9001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 78P B 9001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.