



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:31 PM UTC

PDB ID : 7KLG / pdb_00007klg
Title : SARS-CoV-2 RBD in complex with Fab 15033
Authors : Li, Z.; Rini, J.M.
Deposited on : 2020-10-30
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	225	8% 86% 13%
1	I	225	13% 88% 11%
2	L	214	90% 10%
2	M	214	3% 89% 11%
3	A	201	6% 88% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	201	8% 87% 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 15033 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	223	Total	C	N	O	S	0	0	0
			1635	1031	272	324	8			
1	I	223	Total	C	N	O	S	0	0	0
			1635	1031	272	324	8			

- Molecule 2 is a protein called Fab 15033 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	214	Total	C	N	O	S	0	0	0
			1648	1029	277	336	6			
2	M	214	Total	C	N	O	S	0	0	0
			1648	1029	277	336	6			

- Molecule 3 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	195	Total	C	N	O	S	0	0	0
			1543	989	257	289	8			
3	A	200	Total	C	N	O	S	0	0	0
			1580	1013	264	295	8			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab 15033 heavy chain




- Molecule 1: Fab 15033 heavy chain





- Molecule 2: Fab 15033 light chain





- Molecule 2: Fab 15033 light chain



- Molecule 3: Spike glycoprotein



• Molecule 3: Spike glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	197.10Å 197.10Å 211.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.99 – 3.20 47.99 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.99-3.20) 86.8 (47.99-3.20)	Depositor EDS
R_{merge}	0.29	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.52 (at 3.07Å)	Xtrriage
Refinement program	PHENIX 1.16rc1_3535	Depositor
R, R_{free}	0.271 , 0.290 0.277 , 0.298	Depositor DCC
R_{free} test set	2332 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	77.7	Xtrriage
Anisotropy	0.095	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9717	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.18 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5823e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	H	0.19	0/1675	0.40	0/2280
1	I	0.22	0/1675	0.42	0/2280
2	L	0.19	0/1683	0.42	0/2283
2	M	0.19	0/1683	0.42	0/2283
3	A	0.21	0/1625	0.39	0/2213
3	B	0.19	0/1587	0.39	0/2161
All	All	0.20	0/9928	0.41	0/13500

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1635	0	1586	18	0
1	I	1635	0	1586	15	0
2	L	1648	0	1602	18	0
2	M	1648	0	1602	14	0
3	A	1580	0	1498	14	0
3	B	1543	0	1459	12	0
4	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	14	0	13	0	0
All	All	9717	0	9359	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:156:LEU:HD21	2:L:195:LEU:HD22	1.72	0.70
1:I:52:TRP:CD1	2:M:116:ILE:HD11	2.28	0.69
3:B:383:SER:O	3:B:387:LEU:HD21	1.92	0.69
3:B:358:ILE:HG22	3:B:524:VAL:HG21	1.81	0.62
1:H:156:LEU:HD12	1:H:194:SER:HB3	1.83	0.60
3:A:338:PHE:HE1	3:A:358:ILE:HD13	1.66	0.59
1:I:167:VAL:HG23	1:I:213:VAL:HG22	1.85	0.58
2:L:135:VAL:HG11	2:L:225:VAL:HG11	1.86	0.57
2:L:195:LEU:HD23	2:L:196:SER:N	2.19	0.56
2:M:187:ASP:OD1	2:M:188:SER:N	2.39	0.56
3:B:359:SER:HA	3:B:524:VAL:HG23	1.88	0.55
1:I:127:SER:HB3	1:I:161:PHE:CZ	2.41	0.55
2:L:155:LEU:HD11	2:L:157:ASN:HB3	1.88	0.55
1:I:8:GLY:O	1:I:19:LEU:HD11	2.07	0.54
2:L:76:PHE:CD1	2:L:91:ILE:HD12	2.42	0.54
2:L:56:SER:O	2:L:57:ALA:HB3	2.08	0.54
1:H:2:VAL:HG11	1:H:117:TYR:CD1	2.43	0.54
1:H:127:SER:HB3	1:H:161:PHE:CZ	2.43	0.54
3:B:393:THR:O	3:B:523:THR:HG22	2.09	0.53
3:A:336:CYS:HB2	3:A:363:ALA:HB2	1.91	0.53
1:I:185:LEU:HD13	1:I:191:TYR:CZ	2.44	0.53
2:L:76:PHE:CE1	2:L:91:ILE:HD12	2.45	0.52
2:M:4:MET:HE3	2:M:23:CYS:SG	2.49	0.52
1:H:52:TRP:CG	2:L:116:ILE:HG12	2.44	0.52
3:B:433:VAL:HG22	3:B:512:VAL:HG22	1.91	0.52
2:L:36:SER:OG	2:L:37:SER:N	2.43	0.52
2:M:56:SER:O	2:M:57:ALA:HB3	2.10	0.51
1:H:106:ARG:O	1:H:115:MET:HA	2.10	0.51
1:I:12:LEU:C	1:I:12:LEU:HD13	2.36	0.51
1:H:52:TRP:CD1	2:L:116:ILE:HD11	2.46	0.50
2:M:195:LEU:HD23	2:M:196:SER:N	2.27	0.50
3:B:520:ALA:HB1	3:B:521:PRO:HD2	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:347:PHE:HB3	3:A:401:VAL:HG23	1.94	0.50
1:I:156:LEU:HD12	1:I:194:SER:HB3	1.94	0.50
2:M:36:SER:OG	2:M:37:SER:N	2.45	0.50
2:M:95:GLN:HG3	2:M:96:PRO:HD2	1.93	0.50
1:H:181:PHE:O	1:H:193:LEU:HD12	2.12	0.49
2:M:206:TYR:O	2:M:212:TYR:OH	2.30	0.48
2:M:155:LEU:C	2:M:156:LEU:HD12	2.38	0.47
1:H:229:LYS:HD3	2:L:234:CYS:HB2	1.97	0.47
1:I:12:LEU:HD13	1:I:13:VAL:N	2.29	0.47
2:L:133:PRO:HB2	2:L:156:LEU:HB2	1.96	0.47
1:I:2:VAL:HG11	1:I:117:TYR:CD1	2.50	0.47
1:I:13:VAL:HG11	1:I:94:LEU:HD13	1.97	0.46
1:H:91:MET:HE1	1:H:102:TYR:CZ	2.51	0.46
3:B:338:PHE:HE1	3:B:358:ILE:HD13	1.79	0.46
2:M:120:GLN:OE1	2:M:120:GLN:N	2.41	0.46
1:I:19:LEU:HD23	1:I:124:VAL:HG23	1.99	0.45
1:I:106:ARG:O	1:I:115:MET:HA	2.16	0.45
1:H:13:VAL:HG11	1:H:94:LEU:HD13	1.98	0.45
1:H:12:LEU:C	1:H:12:LEU:HD13	2.42	0.45
2:M:135:VAL:HG21	2:M:225:VAL:HG11	1.98	0.44
1:H:99:THR:HG23	1:H:99:THR:O	2.17	0.44
3:B:358:ILE:HB	3:B:395:VAL:HB	1.99	0.44
3:A:338:PHE:CE1	3:A:358:ILE:HD13	2.50	0.44
3:A:433:VAL:HG22	3:A:512:VAL:HG22	1.99	0.44
3:A:330:PRO:HD3	3:A:525:CYS:SG	2.58	0.43
1:H:216:LYS:N	1:H:217:PRO:CD	2.82	0.43
3:B:467:ASP:OD2	3:B:469:SER:HB3	2.18	0.43
1:H:166:THR:OG1	1:H:214:ASN:HB3	2.19	0.43
2:L:195:LEU:HD23	2:L:195:LEU:C	2.44	0.43
3:A:496:GLY:O	3:A:498:GLN:N	2.49	0.43
2:M:195:LEU:HD23	2:M:195:LEU:C	2.44	0.42
3:A:520:ALA:HB1	3:A:521:PRO:HD2	2.02	0.42
2:L:221:LEU:HD23	2:L:221:LEU:HA	1.93	0.42
3:B:388:ASN:O	3:B:526:GLY:HA3	2.20	0.42
2:M:128:ARG:NH2	2:M:129:THR:OG1	2.53	0.42
2:M:169:LYS:HG2	2:M:174:LEU:HD23	2.01	0.42
1:H:199:VAL:HB	1:H:200:PRO:HD2	2.01	0.41
1:H:8:GLY:O	1:H:19:LEU:HD11	2.20	0.41
2:L:187:ASP:OD2	2:L:188:SER:N	2.53	0.41
1:I:134:PRO:HB2	1:I:157:VAL:CG1	2.51	0.41
2:L:204:ALA:O	2:L:208:LYS:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:95:ARG:C	1:H:126:VAL:HG11	2.45	0.41
3:A:401:VAL:HG22	3:A:509:ARG:HG2	2.02	0.41
2:L:56:SER:HB3	3:A:417:LYS:HD3	2.02	0.41
2:L:109:ARG:NH2	3:A:477:SER:OG	2.54	0.41
1:I:91:MET:HE1	1:I:102:TYR:CZ	2.56	0.40
3:B:388:ASN:HB2	3:B:527:PRO:HD2	2.02	0.40
1:I:153:LEU:HD12	1:I:154:GLY:N	2.37	0.40
1:H:76:PHE:CE1	1:H:91:MET:HB3	2.56	0.40
3:A:437:ASN:OD1	3:A:438:SER:N	2.54	0.40
3:B:439:ASN:OD1	3:B:443:SER:HB3	2.22	0.40
3:A:348:ALA:HB2	3:A:354:ASN:ND2	2.37	0.40
3:A:468:ILE:C	3:A:468:ILE:HD12	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	221/225 (98%)	214 (97%)	7 (3%)	0	100	100
1	I	221/225 (98%)	214 (97%)	7 (3%)	0	100	100
2	L	212/214 (99%)	204 (96%)	8 (4%)	0	100	100
2	M	212/214 (99%)	205 (97%)	7 (3%)	0	100	100
3	A	198/201 (98%)	184 (93%)	13 (7%)	1 (0%)	24	59
3	B	193/201 (96%)	175 (91%)	17 (9%)	1 (0%)	24	59
All	All	1257/1280 (98%)	1196 (95%)	59 (5%)	2 (0%)	43	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	385	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	386	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	177/179 (99%)	177 (100%)	0	100	100
1	I	177/179 (99%)	177 (100%)	0	100	100
2	L	189/189 (100%)	189 (100%)	0	100	100
2	M	189/189 (100%)	189 (100%)	0	100	100
3	A	172/174 (99%)	172 (100%)	0	100	100
3	B	168/174 (97%)	168 (100%)	0	100	100
All	All	1072/1084 (99%)	1072 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	167	GLN
2	L	218	HIS
1	I	40	HIS
3	A	370	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	2001	3	14,14,15	0.28	0	17,19,21	0.54	0
4	NAG	B	2001	3	14,14,15	0.30	0	17,19,21	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	2001	3	-	2/6/23/26	0/1/1/1
4	NAG	B	2001	3	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	2001	NAG	C8-C7-N2-C2
4	B	2001	NAG	O7-C7-N2-C2
4	B	2001	NAG	O5-C5-C6-O6
4	A	2001	NAG	C4-C5-C6-O6
4	A	2001	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	223/225 (99%)	0.49	17 (7%) 20 13	70, 93, 142, 172	0
1	I	223/225 (99%)	0.78	29 (13%) 7 6	70, 94, 151, 172	0
2	L	214/214 (100%)	0.24	1 (0%) 87 76	74, 94, 110, 163	0
2	M	214/214 (100%)	0.62	6 (2%) 55 35	75, 99, 116, 158	0
3	A	200/201 (99%)	0.38	13 (6%) 25 16	67, 83, 122, 153	0
3	B	195/201 (97%)	0.64	16 (8%) 17 11	69, 90, 140, 165	0
All	All	1269/1280 (99%)	0.53	82 (6%) 25 16	67, 94, 137, 172	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	332	ILE	5.8
2	M	233	GLU	3.6
3	A	371	SER	3.5
3	A	330	PRO	3.5
1	H	131	THR	3.3
3	B	371	SER	3.3
1	I	97	GLU	3.2
3	A	370	ASN	3.1
3	A	329	PHE	3.0
1	I	179	HIS	3.0
1	I	127	SER	2.9
1	I	198	THR	2.9
3	B	498	GLN	2.9
1	I	231	CYS	2.9
2	M	97	GLU	2.9
1	I	151	ALA	2.8
3	B	369	TYR	2.8
1	H	222	VAL	2.8
1	I	196	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	I	161	PHE	2.8
3	B	518	LEU	2.7
3	A	387	LEU	2.7
2	M	188	SER	2.7
1	I	144	LYS	2.6
3	B	390	LEU	2.6
1	H	146	THR	2.6
1	H	161	PHE	2.6
1	I	142	SER	2.6
3	A	518	LEU	2.6
1	I	120	GLN	2.6
3	B	370	ASN	2.6
3	A	365	TYR	2.5
3	A	369	TYR	2.5
2	M	230	ASN	2.5
3	A	445	VAL	2.5
1	I	209	TYR	2.5
1	I	180	THR	2.5
1	H	151	ALA	2.4
1	I	128	SER	2.4
3	B	373	SER	2.4
1	I	226	VAL	2.4
1	I	183	ALA	2.4
3	A	519	HIS	2.4
3	A	373	SER	2.4
1	I	141	PRO	2.4
1	H	165	VAL	2.3
1	H	224	LYS	2.3
1	I	138	PRO	2.3
1	I	143	SER	2.3
1	I	157	VAL	2.3
1	I	217	PRO	2.2
1	I	228	PRO	2.2
1	H	147	SER	2.2
3	A	335	LEU	2.2
1	H	152	ALA	2.2
1	H	164	PRO	2.2
1	I	147	SER	2.2
3	B	392	PHE	2.2
1	H	209	TYR	2.2
1	H	226	VAL	2.2
2	M	174	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	365	TYR	2.1
1	H	156	LEU	2.1
1	I	195	SER	2.1
2	L	1	ASP	2.1
1	H	231	CYS	2.1
1	H	162	PRO	2.1
1	I	162	PRO	2.1
1	H	120	GLN	2.1
3	B	384	PRO	2.1
3	B	521	PRO	2.1
2	M	191	SER	2.1
3	B	358	ILE	2.1
1	I	223	ASP	2.1
1	I	131	THR	2.1
1	I	152	ALA	2.1
1	H	177	GLY	2.1
3	B	519	HIS	2.0
3	B	520	ALA	2.0
3	B	445	VAL	2.0
3	B	504	GLY	2.0
1	I	13	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	2001	14/15	0.75	0.16	107,108,111,113	0
4	NAG	A	2001	14/15	0.83	0.16	101,107,111,114	0

6.5 Other polymers [i](#)

There are no such residues in this entry.