



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 11:53 AM UTC

PDB ID : 1KMO / pdb_00001kmo
Title : Crystal structure of the Outer Membrane Transporter FecA
Authors : Ferguson, A.D.; Chakraborty, R.; Smith, B.S.; Esser, L.; van der Helm, D.;
Deisenhofer, J.
Deposited on : 2001-12-17
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	774	3% 63% 20% • 15%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iron(III) dicitrate transport protein fecA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	661	Total	C	N	O	S	0	0	0
			5184	3244	912	1014	14			

- Molecule 2 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: $C_{14}H_{31}NO$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		
2	A	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 3 is HEPTANE-1,2,3-TRIOL (CCD ID: HTO) (formula: $C_7H_{16}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	7	3		
3	A	1	Total	C	O	0	0
			10	7	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	386	Total	O	0	0
			386	386		

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	117.08Å 88.09Å 94.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.71 – 2.00 19.71 – 2.00	Depositor EDS
% Data completeness (in resolution range)	84.5 (19.71-2.00) 84.4 (19.71-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.94Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.207 , 0.245 0.212 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5862	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HTO, LDA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	1/5315 (0.0%)	1.03	28/7220 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	284	MET	SD-CE	-5.27	1.66	1.79

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	717	SER	N-CA-C	-7.43	97.10	109.07
1	A	427	GLY	N-CA-C	-6.95	96.71	113.18
1	A	535	GLU	N-CA-C	-6.48	99.33	109.25
1	A	269	GLY	N-CA-C	6.27	119.88	110.97
1	A	515	PHE	N-CA-C	6.23	118.77	108.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5184	0	4898	139	0
2	A	272	0	527	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	20	0	32	0	0
4	A	386	0	0	18	0
All	All	5862	0	5457	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:ARG:HG2	1:A:159:ARG:HH11	1.09	1.14
1:A:226:GLU:HG2	1:A:740:LYS:HG2	1.47	0.96
1:A:533:GLU:HG3	1:A:534:PRO:HD2	1.54	0.89
1:A:385:ARG:HH11	1:A:385:ARG:HB3	1.40	0.85
1:A:159:ARG:HG2	1:A:159:ARG:NH1	1.91	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	659/774 (85%)	638 (97%)	15 (2%)	6 (1%)	14 9

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	LEU
1	A	428	GLN
1	A	223	PHE
1	A	693	GLY
1	A	694	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	544/636 (86%)	515 (95%)	29 (5%)	20	18

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	507	LEU
1	A	701	LEU
1	A	575	VAL
1	A	695	GLN
1	A	551	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	570	GLN
1	A	603	ASN
1	A	695	GLN
1	A	275	HIS
1	A	351	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HTO	A	759	-	9,9,9	1.37	1 (11%)	10,10,10	0.38	0
2	LDA	A	756	-	13,15,15	1.88	1 (7%)	14,17,17	1.77	4 (28%)
2	LDA	A	754	-	13,15,15	1.91	1 (7%)	14,17,17	1.71	3 (21%)
2	LDA	A	744	-	13,15,15	1.96	1 (7%)	14,17,17	1.59	4 (28%)
3	HTO	A	760	-	9,9,9	1.58	2 (22%)	10,10,10	0.44	0
2	LDA	A	758	-	13,15,15	1.89	1 (7%)	14,17,17	1.75	4 (28%)
2	LDA	A	745	-	13,15,15	2.08	1 (7%)	14,17,17	1.58	4 (28%)
2	LDA	A	750	-	13,15,15	1.91	1 (7%)	14,17,17	1.75	5 (35%)
2	LDA	A	747	-	13,15,15	1.97	1 (7%)	14,17,17	1.65	4 (28%)
2	LDA	A	742	-	13,15,15	1.93	1 (7%)	14,17,17	1.76	4 (28%)
2	LDA	A	753	-	13,15,15	1.94	1 (7%)	14,17,17	1.69	4 (28%)
2	LDA	A	746	-	13,15,15	1.95	1 (7%)	14,17,17	1.69	4 (28%)
2	LDA	A	757	-	13,15,15	2.00	1 (7%)	14,17,17	1.67	4 (28%)
2	LDA	A	752	-	13,15,15	1.92	1 (7%)	14,17,17	1.71	4 (28%)
2	LDA	A	743	-	13,15,15	1.95	1 (7%)	14,17,17	1.73	4 (28%)
2	LDA	A	751	-	13,15,15	1.94	1 (7%)	14,17,17	1.76	5 (35%)
2	LDA	A	749	-	13,15,15	1.95	1 (7%)	14,17,17	1.66	4 (28%)
2	LDA	A	755	-	13,15,15	2.05	1 (7%)	14,17,17	1.66	4 (28%)
2	LDA	A	748	-	13,15,15	2.01	1 (7%)	14,17,17	1.63	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HTO	A	759	-	-	2/10/10/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LDA	A	756	-	-	8/13/13/13	-
2	LDA	A	754	-	-	11/13/13/13	-
2	LDA	A	744	-	-	8/13/13/13	-
3	HTO	A	760	-	-	2/10/10/10	-
2	LDA	A	758	-	-	8/13/13/13	-
2	LDA	A	745	-	-	11/13/13/13	-
2	LDA	A	750	-	-	8/13/13/13	-
2	LDA	A	747	-	-	9/13/13/13	-
2	LDA	A	742	-	-	8/13/13/13	-
2	LDA	A	753	-	-	11/13/13/13	-
2	LDA	A	746	-	-	11/13/13/13	-
2	LDA	A	757	-	-	10/13/13/13	-
2	LDA	A	752	-	-	9/13/13/13	-
2	LDA	A	743	-	-	8/13/13/13	-
2	LDA	A	751	-	-	11/13/13/13	-
2	LDA	A	749	-	-	8/13/13/13	-
2	LDA	A	755	-	-	7/13/13/13	-
2	LDA	A	748	-	-	10/13/13/13	-

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	745	LDA	O1-N1	-7.13	1.24	1.42
2	A	755	LDA	O1-N1	-6.94	1.25	1.42
2	A	757	LDA	O1-N1	-6.85	1.25	1.42
2	A	748	LDA	O1-N1	-6.81	1.25	1.42
2	A	744	LDA	O1-N1	-6.73	1.25	1.42

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	756	LDA	CM1-N1-C1	-3.75	102.35	110.23
2	A	742	LDA	CM1-N1-C1	-3.73	102.40	110.23
2	A	758	LDA	CM1-N1-C1	-3.69	102.47	110.23
2	A	751	LDA	CM1-N1-C1	-3.55	102.77	110.23
2	A	750	LDA	CM1-N1-C1	-3.55	102.77	110.23

There are no chirality outliers.

5 of 160 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	753	LDA	C2-C3-C4-C5
2	A	753	LDA	C11-C10-C9-C8
2	A	748	LDA	C2-C3-C4-C5
2	A	749	LDA	C7-C8-C9-C10
2	A	750	LDA	C7-C8-C9-C10

There are no ring outliers.

9 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	754	LDA	1	0
2	A	745	LDA	2	0
2	A	750	LDA	1	0
2	A	747	LDA	1	0
2	A	742	LDA	1	0
2	A	743	LDA	2	0
2	A	751	LDA	1	0
2	A	749	LDA	1	0
2	A	755	LDA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	661/774 (85%)	0.02	20 (3%)	52 51	13, 22, 39, 79	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	694	PRO	8.1
1	A	532	VAL	4.6
1	A	695	GLN	4.5
1	A	696	MET	4.2
1	A	693	GLY	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
2	LDA	A	753	16/16	0.51	0.33	81,82,85,85	0
3	HTO	A	759	10/10	0.52	0.32	60,62,63,65	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	LDA	A	755	16/16	0.62	0.31	64,71,74,74	0
2	LDA	A	746	16/16	0.62	0.29	71,72,75,75	0
2	LDA	A	747	16/16	0.64	0.31	80,81,84,84	0
2	LDA	A	758	16/16	0.65	0.31	76,77,80,80	0
2	LDA	A	748	16/16	0.65	0.29	64,65,67,67	0
3	HTO	A	760	10/10	0.65	0.24	67,67,69,71	0
2	LDA	A	757	16/16	0.68	0.27	74,75,79,79	0
2	LDA	A	750	16/16	0.69	0.25	57,62,71,72	0
2	LDA	A	744	16/16	0.72	0.30	66,69,73,73	0
2	LDA	A	752	16/16	0.72	0.28	85,85,85,86	0
2	LDA	A	756	16/16	0.73	0.27	59,62,67,67	0
2	LDA	A	743	16/16	0.74	0.28	60,62,67,67	0
2	LDA	A	749	16/16	0.75	0.22	58,62,67,67	0
2	LDA	A	754	16/16	0.76	0.24	66,67,68,69	0
2	LDA	A	751	16/16	0.78	0.26	56,64,75,75	0
2	LDA	A	745	16/16	0.78	0.25	66,70,72,72	0
2	LDA	A	742	16/16	0.82	0.20	53,64,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.