



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:46 PM UTC

PDB ID : 3KMU / pdb_00003kmu
Title : Crystal structure of the ILK/alpha-parvin core complex (apo)
Authors : Fukuda, K.; Qin, J.
Deposited on : 2009-11-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	5% 82% 16% ..
2	B	129	39% 84% 11% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin-linked kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	267	2136	1365	376	378	3	14	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	expression tag	UNP Q13418
A	346	SER	CYS	engineered mutation	UNP Q13418
A	422	SER	CYS	engineered mutation	UNP Q13418

- Molecule 2 is a protein called Alpha-parvin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	B	124	1011	662	160	186	1	2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP Q9NVD7
B	-3	SER	-	expression tag	UNP Q9NVD7
B	-2	HIS	-	expression tag	UNP Q9NVD7
B	-1	MSE	-	expression tag	UNP Q9NVD7


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	248	Total	O	0	0
			248	248		
3	B	50	Total	O	0	0
			50	50		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Integrin-linked kinase



• Molecule 2: Alpha-parvin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.38Å 117.72Å 47.34Å 90.00° 101.68° 90.00°	Depositor
Resolution (Å)	58.82 – 1.80 58.82 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (58.82-1.80) 97.5 (58.82-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.200 , 0.236 0.202 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3445	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	3/2181 (0.1%)	0.95	1/2936 (0.0%)
2	B	0.72	0/1032	0.94	1/1394 (0.1%)
All	All	0.93	3/3213 (0.1%)	0.94	2/4330 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	337	MSE	SE-CE	-13.39	1.55	1.95
1	A	398	MSE	SE-CE	-11.69	1.60	1.95
1	A	375	MSE	CG-SE	5.36	2.11	1.95

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	398	MSE	CG-SE-CE	5.51	111.05	98.92
2	B	276	LEU	N-CA-C	5.45	118.15	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2136	0	2139	44	0
2	B	1011	0	1019	7	0
3	A	248	0	0	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	50	0	0	1	0
All	All	3445	0	3158	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:398:MSE:HE3	1:A:402:MSE:HG2	1.52	0.90
1:A:398:MSE:HE1	3:B:134:HOH:O	1.73	0.89
1:A:247:HIS:HD2	1:A:249:ASN:H	1.22	0.87
1:A:340:VAL:HB	3:A:514:HOH:O	1.85	0.77
1:A:296:LYS:NZ	3:A:464:HOH:O	2.21	0.74
2:B:265:LYS:O	2:B:269:ILE:HG13	1.87	0.74
1:A:342:PHE:H	1:A:345:GLN:HE21	1.37	0.73
1:A:398:MSE:CE	1:A:402:MSE:SE	2.87	0.72
1:A:398:MSE:CE	1:A:402:MSE:HG2	2.20	0.72
1:A:282:HIS:HE1	1:A:384:GLU:OE2	1.72	0.71
1:A:247:HIS:CD2	1:A:249:ASN:H	2.07	0.71
1:A:423:LYS:O	1:A:427:ILE:HD13	1.92	0.70
1:A:402:MSE:HE2	1:A:406:LEU:HD12	1.75	0.68
1:A:402:MSE:HA	1:A:402:MSE:HE3	1.75	0.67
1:A:252:PRO:O	1:A:269:THR:HG23	1.96	0.65
1:A:282:HIS:HD2	3:A:21:HOH:O	1.81	0.63
1:A:402:MSE:HE2	1:A:406:LEU:CD1	2.30	0.61
2:B:365:PHE:O	2:B:369:ARG:HG3	2.02	0.59
1:A:398:MSE:HE3	1:A:402:MSE:CG	2.30	0.59
2:B:287:GLU:HG3	2:B:354:LEU:HD13	1.83	0.59
1:A:398:MSE:CE	1:A:402:MSE:CG	2.81	0.58
1:A:341:LYS:HA	1:A:345:GLN:HE21	1.68	0.58
1:A:398:MSE:HE2	1:A:402:MSE:SE	2.56	0.56
1:A:243:ARG:HD2	3:A:471:HOH:O	2.06	0.55
1:A:341:LYS:HA	1:A:345:GLN:NE2	2.27	0.50
1:A:342:PHE:N	1:A:345:GLN:HE21	2.06	0.50
2:B:257:ALA:HB1	2:B:260:LYS:HG2	1.92	0.50
1:A:450:GLN:O	1:A:451:ASP:CB	2.60	0.48
1:A:269:THR:HG21	3:A:61:HOH:O	2.14	0.48
1:A:311:LEU:HD12	1:A:315:ILE:HD13	1.96	0.47
1:A:345:GLN:C	3:A:526:HOH:O	2.58	0.47
1:A:402:MSE:CE	1:A:406:LEU:CD1	2.92	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:450:GLN:O	1:A:451:ASP:HB2	2.14	0.47
1:A:342:PHE:H	1:A:345:GLN:NE2	2.07	0.47
1:A:197:THR:OG1	3:A:482:HOH:O	2.21	0.46
2:B:257:ALA:HB1	2:B:260:LYS:CG	2.46	0.46
1:A:257:CYS:HB2	1:A:266:THR:HB	1.98	0.44
1:A:218:VAL:CG1	1:A:271:TRP:HE3	2.30	0.44
1:A:318:HIS:CD2	3:A:54:HOH:O	2.70	0.44
1:A:312:GLU:HA	1:A:313:PRO:HA	1.87	0.44
1:A:282:HIS:CD2	3:A:21:HOH:O	2.65	0.43
1:A:220:LYS:HE2	1:A:222:LEU:HD21	2.00	0.43
1:A:398:MSE:HE2	1:A:402:MSE:CG	2.49	0.42
1:A:192:GLN:NE2	3:A:37:HOH:O	2.52	0.42
2:B:295:TYR:O	2:B:299:LEU:HB2	2.20	0.41
1:A:269:THR:HG22	1:A:270:HIS:N	2.35	0.41
1:A:306:ALA:O	1:A:310:THR:HG23	2.21	0.41
1:A:211:ARG:NH1	1:A:216:ASP:OD1	2.54	0.41
1:A:199:LEU:HD11	1:A:271:TRP:HZ3	1.86	0.40
2:B:269:ILE:HG22	2:B:273:ASN:ND2	2.36	0.40
1:A:263:PRO:HB2	1:A:264:HIS:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/271 (98%)	264 (100%)	1 (0%)	0	100	100
2	B	122/129 (95%)	118 (97%)	4 (3%)	0	100	100
All	All	387/400 (97%)	382 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/226 (105%)	233 (98%)	4 (2%)	53	45
2	B	115/116 (99%)	111 (96%)	4 (4%)	32	19
All	All	352/342 (103%)	344 (98%)	8 (2%)	44	33

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	200	ASN
1	A	315	ILE
1	A	413	ILE
1	A	429	MSE
2	B	270	THR
2	B	282	GLU
2	B	299	LEU
2	B	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	GLN
1	A	194	ASN
1	A	213	GLN
1	A	247	HIS
1	A	264	HIS
1	A	282	HIS
1	A	286	ASN
1	A	293	GLN
1	A	345	GLN
1	A	450	GLN
2	B	262	ASN
2	B	275	HIS
2	B	363	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/271 (93%)	0.18	14 (5%) 30 29	9, 20, 37, 50	0
2	B	122/129 (94%)	1.89	50 (40%) 0 0	17, 40, 73, 74	1 (0%)
All	All	375/400 (93%)	0.73	64 (17%) 4 3	9, 25, 61, 74	1 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	249	ALA	6.4
2	B	263	VAL	6.2
2	B	264	VAL	5.8
2	B	253	LEU	5.5
2	B	261	LEU	5.1
2	B	259	ASP	5.0
1	A	185	HIS	4.4
2	B	256	HIS	4.2
2	B	254	PHE	4.1
2	B	269	ILE	4.1
2	B	250	PHE	4.1
2	B	371	VAL	3.9
1	A	412	THR	3.9
2	B	257	ALA	3.7
2	B	258	PRO	3.6
1	A	451	ASP	3.6
2	B	260	LYS	3.5
2	B	267	THR	3.3
2	B	354	LEU	3.2
2	B	366	THR	3.2
2	B	262	ASN	3.1
2	B	268	LEU	3.1
2	B	270	THR	3.1
2	B	266	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	413	ILE	3.0
2	B	271	PHE	2.9
1	A	346	SER	2.9
2	B	358	LEU	2.9
2	B	362	TYR	2.9
2	B	351	ASN	2.8
2	B	368	TYR	2.8
2	B	252	THR	2.8
1	A	287	PHE	2.8
1	A	196	LEU	2.7
2	B	344	PRO	2.7
2	B	265	LYS	2.7
2	B	272	VAL	2.6
2	B	339	LEU	2.6
1	A	191	LYS	2.6
2	B	283	VAL	2.6
1	A	286	ASN	2.6
1	A	416	GLY	2.5
2	B	372	GLU	2.4
2	B	369	ARG	2.3
2	B	364	LEU	2.3
2	B	365	PHE	2.3
1	A	427	ILE	2.3
1	A	284	GLY	2.3
2	B	350	VAL	2.3
2	B	361	LEU	2.3
1	A	285	THR	2.3
2	B	280	ASN	2.3
2	B	343	LYS	2.2
2	B	363	ASN	2.2
2	B	251	ASP	2.1
2	B	255	ASP	2.1
2	B	281	LEU	2.1
2	B	342	PRO	2.1
2	B	370	ASN	2.1
2	B	341	LYS	2.1
2	B	276	LEU	2.1
2	B	286	LEU	2.0
1	A	203	HIS	2.0
2	B	360	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.