



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 7, 2026 – 11:34 PM UTC

PDB ID : 2KN6 / pdb_00002kn6
Title : Structure of full-length human ASC (Apoptosis-associated speck-like protein containing a CARD)
Authors : De Alba, E.
Deposited on : 2009-08-16

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	215	

2 Ensemble composition and analysis

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:29, A:38-A:89 (79)	0.29	10
2	A:116-A:193 (78)	0.60	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 18, 19, 20
2	2, 16
Single-model clusters	17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3060 atoms, of which 1538 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Apoptosis-associated speck-like protein containing a CARD.

Mol	Chain	Residues	Atoms						Trace
1	A	195	Total	C	H	N	O	S	0
			3060	964	1538	266	286	6	

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q9ULZ3
A	-18	GLY	-	expression tag	UNP Q9ULZ3
A	-17	SER	-	expression tag	UNP Q9ULZ3
A	-16	SER	-	expression tag	UNP Q9ULZ3
A	-15	HIS	-	expression tag	UNP Q9ULZ3
A	-14	HIS	-	expression tag	UNP Q9ULZ3
A	-13	HIS	-	expression tag	UNP Q9ULZ3
A	-12	HIS	-	expression tag	UNP Q9ULZ3
A	-11	HIS	-	expression tag	UNP Q9ULZ3
A	-10	HIS	-	expression tag	UNP Q9ULZ3
A	-9	SER	-	expression tag	UNP Q9ULZ3
A	-8	SER	-	expression tag	UNP Q9ULZ3
A	-7	GLY	-	expression tag	UNP Q9ULZ3
A	-6	LEU	-	expression tag	UNP Q9ULZ3
A	-5	VAL	-	expression tag	UNP Q9ULZ3
A	-4	PRO	-	expression tag	UNP Q9ULZ3
A	-3	ARG	-	expression tag	UNP Q9ULZ3
A	-2	GLY	-	expression tag	UNP Q9ULZ3
A	-1	SER	-	expression tag	UNP Q9ULZ3
A	0	HIS	-	expression tag	UNP Q9ULZ3

5 Refinement protocol and experimental data overview

The models were refined using the following method: *DGSA-distance geometry simulated annealing*.

Of the 300 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	refinement	
TALOS	geometry optimization	

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.56±0.01	0±0/1273 (0.0± 0.0%)	1.10±0.01	0±0/1725 (0.0± 0.0%)
All	All	1.56	2/25460 (0.0%)	1.10	3/34500 (0.0%)

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	155	ASN	N-CA	5.26	1.50	1.46	17	2

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	153	PRO	N-CA-C	-6.59	106.29	114.20	17	2
1	A	78	LEU	N-CA-C	-5.08	107.58	112.97	17	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1253	1274	1274	28±4
All	All	25060	25480	25480	553

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

5 of 169 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:169:TRP:CG	1:A:173:CYS:HG	0.92	1.83	3	15
1:A:52:LEU:HD23	1:A:53:THR:N	0.84	1.87	8	2
1:A:171:TRP:CD1	1:A:172:THR:HG1	0.61	2.13	1	4
1:A:52:LEU:C	1:A:52:LEU:HD12	0.60	2.22	4	18
1:A:177:LEU:C	1:A:177:LEU:HD13	0.59	2.23	15	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/215 (73%)	149±1 (95±1%)	5±1 (3±1%)	2±1 (1±1%)	11	58
All	All	3140/4300 (73%)	2986 (95%)	109 (3%)	45 (1%)	11	58

5 of 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	127	THR	14
1	A	153	PRO	11
1	A	128	ASN	7
1	A	139	LYS	7
1	A	61	LEU	5

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/175 (76%)	119±2 (90±2%)	14±2 (10±2%)	9	53
All	All	2660/3500 (76%)	2388 (90%)	272 (10%)	9	53

5 of 42 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	52	LEU	20
1	A	154	THR	20
1	A	155	ASN	20
1	A	166	THR	19
1	A	8	ILE	16

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided