



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 08:07 PM UTC

PDB ID : 4KO6 / pdb_00004ko6
Title : Investigating the functional significance of the interlocked pair structural determinants in *Pseudomonas aeruginosa* azurin (V31I/V95K/Y108F)
Authors : Inampudi, K.K.; Meng, W.; Tobin, P.H.; Wilson, C.J.
Deposited on : 2013-05-11
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	 2% 80% 19% .
1	B	128	 2% 77% 17% . .
1	C	128	 11% 70% 27% .
1	D	128	 12% 81% 15% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Azurin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	128	Total	C	N	O	S	0	0	0
			975	609	165	192	9			
1	B	128	Total	C	N	O	S	0	0	0
			975	609	165	192	9			
1	C	128	Total	C	N	O	S	0	0	0
			975	609	165	192	9			
1	D	128	Total	C	N	O	S	0	0	0
			975	609	165	192	9			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	ILE	VAL	engineered mutation	UNP P00282
A	95	LYS	VAL	engineered mutation	UNP P00282
A	108	PHE	TYR	engineered mutation	UNP P00282
B	31	ILE	VAL	engineered mutation	UNP P00282
B	95	LYS	VAL	engineered mutation	UNP P00282
B	108	PHE	TYR	engineered mutation	UNP P00282
C	31	ILE	VAL	engineered mutation	UNP P00282
C	95	LYS	VAL	engineered mutation	UNP P00282
C	108	PHE	TYR	engineered mutation	UNP P00282
D	31	ILE	VAL	engineered mutation	UNP P00282
D	95	LYS	VAL	engineered mutation	UNP P00282
D	108	PHE	TYR	engineered mutation	UNP P00282

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	B	1	Total	Cu	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total 1	Cu 1	0	0
2	D	1	Total 1	Cu 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	122	Total 122	O 122	0	0
3	B	124	Total 124	O 124	0	0
3	C	96	Total 96	O 96	0	0
3	D	104	Total 104	O 104	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Azurin





- Molecule 1: Azurin



- Molecule 1: Azurin



- Molecule 1: Azurin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.67Å 50.17Å 70.81Å 90.00° 99.30° 90.00°	Depositor
Resolution (Å)	32.87 – 1.74 32.87 – 1.74	Depositor EDS
% Data completeness (in resolution range)	89.3 (32.87-1.74) 89.3 (32.87-1.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.14 (at 1.74Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.226 , 0.273 0.229 , 0.274	Depositor DCC
R_{free} test set	2243 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4350	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	4/992 (0.4%)	0.89	3/1335 (0.2%)
1	B	0.65	2/992 (0.2%)	0.96	5/1335 (0.4%)
1	C	0.46	0/992	0.84	2/1335 (0.1%)
1	D	0.53	0/992	0.94	3/1335 (0.2%)
All	All	0.61	6/3968 (0.2%)	0.91	13/5340 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	CYS	C-O	-8.75	1.13	1.24
1	B	2	GLU	CA-C	-7.03	1.44	1.52
1	A	4	SER	C-O	-6.97	1.15	1.23
1	B	95	LYS	C-O	-6.82	1.16	1.24
1	A	121	MET	C-O	-6.02	1.17	1.23
1	A	7	ILE	C-O	-5.02	1.19	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1	ALA	CA-C-N	-8.50	110.61	122.93
1	B	1	ALA	C-N-CA	-8.50	110.61	122.93
1	C	2	GLU	N-CA-C	-6.42	101.44	110.50
1	D	66	SER	CA-C-N	-6.41	115.93	122.27
1	D	66	SER	C-N-CA	-6.41	115.93	122.27
1	A	66	SER	CA-C-N	-6.39	115.95	122.27
1	A	66	SER	C-N-CA	-6.39	115.95	122.27
1	C	38	ASN	N-CA-C	6.06	122.05	113.02
1	B	66	SER	N-CA-C	-5.72	101.16	110.20
1	B	39	LEU	CA-C-N	5.33	125.80	120.52
1	B	39	LEU	C-N-CA	5.33	125.80	120.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	108	PHE	N-CA-C	-5.22	102.33	110.42
1	A	7	ILE	CB-CA-C	-5.00	102.79	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	975	0	958	23	0
1	B	975	0	958	28	0
1	C	975	0	958	33	0
1	D	975	0	958	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	122	0	0	10	0
3	B	124	0	0	7	0
3	C	96	0	0	17	0
3	D	104	0	0	8	1
All	All	4350	0	3832	97	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:24:LYS:NZ	3:D:402:HOH:O	1.84	1.07
1:D:13:MET:SD	3:D:349:HOH:O	2.16	1.03
1:C:107:GLN:O	3:C:380:HOH:O	1.85	0.94
1:D:64:MET:SD	3:D:307:HOH:O	2.38	0.81
1:D:106:GLU:O	3:D:342:HOH:O	1.99	0.81
1:A:113:THR:OG1	3:A:398:HOH:O	1.92	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1:ALA:O	3:C:366:HOH:O	2.00	0.79
1:C:28:GLN:OE1	3:C:361:HOH:O	2.04	0.75
1:B:8:GLN:HG3	3:B:347:HOH:O	1.87	0.73
1:A:128:LYS:NZ	1:B:2:GLU:CD	2.48	0.70
1:A:12:GLN:OE1	3:A:355:HOH:O	2.09	0.70
1:C:21:THR:HG23	1:C:128:LYS:HD3	1.74	0.68
1:A:107:GLN:H	1:B:21:THR:HG23	1.59	0.67
1:C:46:HIS:CE1	3:C:382:HOH:O	2.47	0.67
1:A:128:LYS:HZ3	1:B:2:GLU:CD	2.02	0.66
1:B:79:ARG:HA	3:B:415:HOH:O	1.94	0.66
1:C:44:MET:O	3:C:382:HOH:O	2.14	0.65
1:C:7:ILE:HD11	1:C:20:ILE:HD11	1.78	0.65
1:A:47:ASN:ND2	3:A:398:HOH:O	2.18	0.64
1:C:28:GLN:HB3	3:C:361:HOH:O	1.98	0.64
1:D:106:GLU:N	3:D:391:HOH:O	2.19	0.63
1:B:57:GLN:HG3	3:B:342:HOH:O	1.99	0.62
1:C:22:VAL:HB	1:C:127:LEU:HD22	1.82	0.62
1:C:44:MET:SD	3:C:382:HOH:O	2.56	0.60
1:D:82:ALA:HB1	1:D:95:LYS:HD3	1.84	0.59
1:A:73:LEU:HG	3:A:398:HOH:O	2.03	0.59
1:C:99:VAL:HA	1:C:102:LEU:HD12	1.86	0.58
1:C:46:HIS:CE1	1:C:121:MET:SD	2.97	0.57
1:D:103:LYS:O	3:D:391:HOH:O	2.17	0.57
1:B:13:MET:O	1:B:120:LEU:HD22	2.04	0.57
1:B:69:ASP:OD2	3:B:385:HOH:O	2.17	0.57
1:B:69:ASP:CG	3:B:385:HOH:O	2.48	0.57
1:C:95:LYS:HE3	1:C:97:PHE:HB3	1.86	0.56
1:D:102:LEU:HB3	1:D:108:PHE:CZ	2.41	0.56
1:B:103:LYS:NZ	1:D:69:ASP:OD2	2.34	0.56
1:A:128:LYS:NZ	1:B:2:GLU:HG2	2.21	0.56
1:C:28:GLN:CB	3:C:361:HOH:O	2.53	0.55
1:C:44:MET:C	3:C:382:HOH:O	2.47	0.55
1:A:48:TRP:HB2	3:A:406:HOH:O	2.07	0.55
1:C:24:LYS:N	1:C:128:LYS:O	2.37	0.55
1:C:28:GLN:HB3	3:C:378:HOH:O	2.06	0.54
1:C:2:GLU:HA	3:C:366:HOH:O	2.09	0.53
1:B:83:HIS:O	1:B:95:LYS:HG3	2.08	0.53
1:D:82:ALA:HB1	1:D:95:LYS:HE3	1.89	0.53
1:A:128:LYS:HZ3	1:B:2:GLU:CG	2.21	0.53
1:A:128:LYS:HZ3	1:B:2:GLU:HG2	1.75	0.52
1:A:124:THR:HG23	3:A:350:HOH:O	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:82:ALA:HB1	1:D:95:LYS:CE	2.41	0.51
1:C:126:THR:OG1	1:C:128:LYS:HD2	2.11	0.51
1:C:13:MET:HG2	1:C:117:HIS:CD2	2.45	0.50
1:A:128:LYS:NZ	1:B:2:GLU:OE2	2.40	0.50
1:A:128:LYS:NZ	1:B:2:GLU:CG	2.76	0.49
1:B:103:LYS:HD3	1:D:69:ASP:OD1	2.12	0.48
1:C:28:GLN:CB	3:C:378:HOH:O	2.60	0.48
1:C:37:GLY:O	1:C:89:SER:HB2	2.12	0.48
1:C:44:MET:CG	3:C:382:HOH:O	2.61	0.48
1:D:82:ALA:HB1	1:D:95:LYS:CD	2.44	0.48
1:B:69:ASP:HB2	3:B:385:HOH:O	2.14	0.47
1:D:7:ILE:HG13	1:D:31:ILE:HG23	1.96	0.47
1:A:10:ASN:HB2	3:A:335:HOH:O	2.15	0.47
1:B:10:ASN:HB2	3:B:326:HOH:O	2.13	0.47
1:D:10:ASN:OD1	1:D:14:GLN:HG3	2.15	0.47
1:C:18:ASN:ND2	3:C:348:HOH:O	2.48	0.46
1:D:103:LYS:C	1:D:127:LEU:HD23	2.40	0.46
1:D:34:SER:CB	1:D:92:LYS:HG2	2.45	0.46
1:D:92:LYS:HE3	3:D:382:HOH:O	2.16	0.46
1:A:106:GLU:HG3	1:A:107:GLN:N	2.30	0.45
1:D:46:HIS:HA	1:D:112:CYS:SG	2.56	0.45
1:A:46:HIS:CE1	1:A:121:MET:SD	3.10	0.45
1:D:34:SER:HB3	1:D:92:LYS:HG2	1.99	0.44
1:A:91:GLU:HG2	3:A:375:HOH:O	2.18	0.44
1:A:39:LEU:HD23	1:A:39:LEU:HA	1.81	0.44
1:B:13:MET:HE2	1:B:117:HIS:HE2	1.83	0.43
1:C:62:ASP:HB3	1:C:74:LYS:HD3	1.99	0.43
1:C:53:ALA:HB2	3:C:380:HOH:O	2.17	0.43
1:C:99:VAL:HG13	1:C:127:LEU:HD11	2.00	0.43
1:B:13:MET:HE3	1:B:44:MET:HE1	2.00	0.43
1:B:111:PHE:HB3	1:B:122:LYS:HB3	2.00	0.43
1:A:118:SER:O	1:A:122:LYS:HD3	2.20	0.42
1:B:84:THR:HG22	1:B:95:LYS:HB2	2.00	0.42
1:C:99:VAL:C	1:C:101:LYS:N	2.75	0.42
1:A:38:ASN:ND2	3:A:411:HOH:O	2.52	0.42
1:C:10:ASN:HB2	3:C:303:HOH:O	2.19	0.42
1:C:33:LEU:HG	1:C:87:ILE:HD11	2.02	0.42
1:D:82:ALA:CB	1:D:95:LYS:HE3	2.50	0.42
1:B:109:MET:HB3	1:B:109:MET:HE2	1.79	0.41
1:B:49:VAL:O	1:B:110:PHE:HA	2.19	0.41
1:C:114:PHE:CG	1:C:115:PRO:HD2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:71:ASP:OD1	3:C:337:HOH:O	2.22	0.41
1:D:102:LEU:HB2	1:D:127:LEU:HD22	2.03	0.41
1:A:47:ASN:C	3:A:406:HOH:O	2.63	0.41
1:B:121:MET:C	1:B:122:LYS:HG3	2.44	0.41
1:D:10:ASN:HB2	3:D:313:HOH:O	2.21	0.41
1:B:82:ALA:HB1	1:B:95:LYS:HD2	2.03	0.41
1:C:97:PHE:HE1	1:C:102:LEU:HD11	1.85	0.40
1:A:13:MET:HE3	1:A:117:HIS:CD2	2.56	0.40
1:B:38:ASN:O	1:B:38:ASN:ND2	2.49	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:314:HOH:O	3:D:367:HOH:O[2_555]	2.05	0.15

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/128 (98%)	122 (97%)	4 (3%)	0	100	100
1	B	126/128 (98%)	123 (98%)	2 (2%)	1 (1%)	16	4
1	C	126/128 (98%)	122 (97%)	3 (2%)	1 (1%)	16	4
1	D	126/128 (98%)	123 (98%)	3 (2%)	0	100	100
All	All	504/512 (98%)	490 (97%)	12 (2%)	2 (0%)	30	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	115	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	115	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/110 (100%)	108 (98%)	2 (2%)	51	30
1	B	110/110 (100%)	101 (92%)	9 (8%)	10	1
1	C	110/110 (100%)	106 (96%)	4 (4%)	31	9
1	D	110/110 (100%)	102 (93%)	8 (7%)	13	2
All	All	440/440 (100%)	417 (95%)	23 (5%)	21	4

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	69	ASP
1	B	2	GLU
1	B	21	THR
1	B	25	SER
1	B	38	ASN
1	B	64	MET
1	B	69	ASP
1	B	95	LYS
1	B	120	LEU
1	B	122	LYS
1	C	12	GLN
1	C	13	MET
1	C	31	ILE
1	C	103	LYS
1	D	2	GLU
1	D	7	ILE
1	D	14	GLN
1	D	39	LEU
1	D	64	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	69	ASP
1	D	115	PRO
1	D	128	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	38	ASN
1	B	107	GLN
1	D	8	GLN
1	D	107	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/128 (100%)	0.34	2 (1%) 70 77	8, 12, 17, 22	0
1	B	128/128 (100%)	0.33	3 (2%) 61 69	8, 12, 20, 26	0
1	C	128/128 (100%)	0.65	14 (10%) 10 14	8, 14, 28, 37	0
1	D	128/128 (100%)	0.73	16 (12%) 8 10	10, 15, 26, 38	0
All	All	512/512 (100%)	0.51	35 (6%) 23 29	8, 13, 25, 38	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	ALA	8.6
1	A	1	ALA	6.1
1	D	1	ALA	5.2
1	C	101	LYS	4.3
1	D	105	GLY	4.2
1	D	104	GLU	3.4
1	D	107	GLN	3.4
1	D	103	LYS	3.2
1	D	64	MET	3.1
1	D	106	GLU	3.1
1	A	128	LYS	3.1
1	C	38	ASN	3.1
1	D	67	GLY	3.0
1	C	100	SER	3.0
1	C	99	VAL	2.8
1	C	37	GLY	2.8
1	D	66	SER	2.7
1	C	102	LEU	2.6
1	B	66	SER	2.6
1	C	25	SER	2.5
1	D	65	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	98	ASP	2.5
1	C	31	ILE	2.5
1	B	2	GLU	2.5
1	C	27	LYS	2.4
1	D	127	LEU	2.4
1	D	39	LEU	2.3
1	D	120	LEU	2.3
1	C	97	PHE	2.3
1	C	28	GLN	2.2
1	D	57	GLN	2.2
1	B	67	GLY	2.2
1	C	2	GLU	2.1
1	D	24	LYS	2.1
1	D	2	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CU	D	201	1/1	0.93	0.13	30,30,30,30	0
2	CU	B	201	1/1	1.00	0.02	11,11,11,11	0
2	CU	C	201	1/1	1.00	0.05	10,10,10,10	0
2	CU	A	201	1/1	1.00	0.02	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.