



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 7, 2026 – 11:22 AM EDT

PDB ID : 5KO2 / pdb\_00005ko2  
Title : Mouse pgp 34 linker deleted mutant Hg derivative  
Authors : Xia, D.; Esser, L.; Zhou, F.  
Deposited on : 2016-06-29  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1248	
1	B	1248	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 37085 atoms, of which 18697 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug resistance protein 1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	1185	18539	5907	9350	1559	1685	38	0	0	0
1	B	1184	18532	5905	9347	1558	1684	38	0	0	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	552	GLN	GLU	engineered mutation	UNP P21447
A	?	-	MET	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	LYS	deletion	UNP P21447
A	?	-	ASP	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	GLY	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	LEU	deletion	UNP P21447
A	?	-	ILE	deletion	UNP P21447
A	?	-	ARG	deletion	UNP P21447
A	?	-	ARG	deletion	UNP P21447
A	?	-	ARG	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	THR	deletion	UNP P21447
A	?	-	ARG	deletion	UNP P21447
A	?	-	LYS	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	ILE	deletion	UNP P21447
A	?	-	CYS	deletion	UNP P21447
A	?	-	GLY	deletion	UNP P21447
A	?	-	PRO	deletion	UNP P21447
A	?	-	HIS	deletion	UNP P21447

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASP	deletion	UNP P21447
A	?	-	GLN	deletion	UNP P21447
A	?	-	ASP	deletion	UNP P21447
A	?	-	ARG	deletion	UNP P21447
A	?	-	LYS	deletion	UNP P21447
A	?	-	LEU	deletion	UNP P21447
A	?	-	SER	deletion	UNP P21447
A	?	-	THR	deletion	UNP P21447
A	?	-	LYS	deletion	UNP P21447
A	?	-	GLU	deletion	UNP P21447
A	1197	GLN	GLU	engineered mutation	UNP P21447
A	1277	HIS	-	expression tag	UNP P21447
A	1278	HIS	-	expression tag	UNP P21447
A	1279	HIS	-	expression tag	UNP P21447
A	1280	HIS	-	expression tag	UNP P21447
A	1281	HIS	-	expression tag	UNP P21447
A	1282	HIS	-	expression tag	UNP P21447
B	552	GLN	GLU	engineered mutation	UNP P21447
B	?	-	MET	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	LYS	deletion	UNP P21447
B	?	-	ASP	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	GLY	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	LEU	deletion	UNP P21447
B	?	-	ILE	deletion	UNP P21447
B	?	-	ARG	deletion	UNP P21447
B	?	-	ARG	deletion	UNP P21447
B	?	-	ARG	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	THR	deletion	UNP P21447
B	?	-	ARG	deletion	UNP P21447
B	?	-	LYS	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	ILE	deletion	UNP P21447
B	?	-	CYS	deletion	UNP P21447
B	?	-	GLY	deletion	UNP P21447
B	?	-	PRO	deletion	UNP P21447
B	?	-	HIS	deletion	UNP P21447

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ASP	deletion	UNP P21447
B	?	-	GLN	deletion	UNP P21447
B	?	-	ASP	deletion	UNP P21447
B	?	-	ARG	deletion	UNP P21447
B	?	-	LYS	deletion	UNP P21447
B	?	-	LEU	deletion	UNP P21447
B	?	-	SER	deletion	UNP P21447
B	?	-	THR	deletion	UNP P21447
B	?	-	LYS	deletion	UNP P21447
B	?	-	GLU	deletion	UNP P21447
B	1197	GLN	GLU	engineered mutation	UNP P21447
B	1277	HIS	-	expression tag	UNP P21447
B	1278	HIS	-	expression tag	UNP P21447
B	1279	HIS	-	expression tag	UNP P21447
B	1280	HIS	-	expression tag	UNP P21447
B	1281	HIS	-	expression tag	UNP P21447
B	1282	HIS	-	expression tag	UNP P21447


- Molecule 2 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	7	Total Hg 7 7	0	0
2	B	7	Total Hg 7 7	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Multidrug resistance protein 1A





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.19Å 114.74Å 375.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.54 – 3.30 33.54 – 3.30	Depositor EDS
% Data completeness (in resolution range)	92.2 (33.54-3.30) 90.9 (33.54-3.30)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 3.33Å)	Xtrriage
Refinement program	PHENIX dev_2443	Depositor
R, $R_{free}$	0.243 , 0.285 0.253 , 0.283	Depositor DCC
$R_{free}$ test set	2000 reflections (3.28%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	108.7	Xtrriage
Anisotropy	0.179	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	37085	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	145.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.20	0/9358	0.37	0/12650
1	B	0.20	0/9354	0.37	0/12645
All	All	0.20	0/18712	0.37	0/25295

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9189	9350	9367	17	0
1	B	9185	9347	9364	21	0
2	A	7	0	0	0	0
2	B	7	0	0	0	0
All	All	18388	18697	18731	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1134:ARG:NH1	1:B:1140:GLU:OE2	2.26	0.68
1:B:1189:GLN:OE1	1:B:1221:ARG:NH2	2.29	0.65
1:B:440:TYR:OH	1:B:901:ARG:NH2	2.30	0.65
1:A:440:TYR:OH	1:A:901:ARG:NH2	2.31	0.63
1:A:1207:GLU:OE2	1:A:1229:ARG:NH1	2.33	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1181/1248 (95%)	1139 (96%)	37 (3%)	5 (0%)	30	60
1	B	1180/1248 (95%)	1140 (97%)	35 (3%)	5 (0%)	30	60
All	All	2361/2496 (95%)	2279 (96%)	72 (3%)	10 (0%)	30	60

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1136	VAL
1	B	1160	LYS
1	A	1160	LYS
1	B	1136	VAL
1	A	690	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	977/1031 (95%)	951 (97%)	26 (3%)	39	63
1	B	977/1031 (95%)	957 (98%)	20 (2%)	48	68
All	All	1954/2062 (95%)	1908 (98%)	46 (2%)	43	65

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	121	VAL
1	B	520	VAL
1	B	166	GLU
1	B	306	TYR
1	B	549	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	608	HIS
1	B	910	GLN
1	B	1234	GLN
1	B	1132	ASN
1	B	1228	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1185/1248 (94%)	0.16	67 (5%) 29 20	70, 131, 234, 358	0
1	B	1184/1248 (94%)	0.12	55 (4%) 37 25	66, 140, 237, 383	0
All	All	2369/2496 (94%)	0.14	122 (5%) 33 22	66, 135, 236, 383	0

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	986	GLN	8.6
1	B	1135	VAL	8.3
1	A	1123	ILE	7.7
1	A	1231	SER	6.7
1	A	986	GLN	5.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HG	A	1301	1/1	0.75	0.13	274,274,274,274	0
2	HG	B	1306	1/1	0.79	0.07	414,414,414,414	0
2	HG	A	1306	1/1	0.83	0.12	368,368,368,368	0
2	HG	B	1307	1/1	0.84	0.08	536,536,536,536	0
2	HG	A	1307	1/1	0.87	0.07	415,415,415,415	0
2	HG	B	1304	1/1	0.91	0.15	385,385,385,385	0
2	HG	B	1302	1/1	0.92	0.05	321,321,321,321	0
2	HG	A	1305	1/1	0.95	0.04	303,303,303,303	0
2	HG	B	1305	1/1	0.95	0.04	347,347,347,347	0
2	HG	A	1302	1/1	0.96	0.08	286,286,286,286	0
2	HG	A	1304	1/1	0.96	0.06	228,228,228,228	0
2	HG	B	1301	1/1	0.97	0.14	182,182,182,182	0
2	HG	B	1303	1/1	0.98	0.03	142,142,142,142	0
2	HG	A	1303	1/1	0.99	0.05	127,127,127,127	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.