



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:41 PM UTC

PDB ID : 3KP6 / pdb_00003kp6
Title : Staphylococcus epidermidis TcaR in complex with salicylate
Authors : Chang, Y.M.; Chen, C.K.; Yeh, Y.J.; Wang, A.H.
Deposited on : 2009-11-15
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	
1	B	151	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SAL	B	3003	-	-	X	-
3	CAC	B	2001	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator TcaR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1119	709	196	211	3			
1	B	137	Total	C	N	O	S	0	0	0
			1099	697	188	211	3			

- Molecule 2 is 2-HYDROXYBENZOIC ACID (CCD ID: SAL) (formula: C₇H₆O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	7	3		
2	A	1	Total	C	O	0	0
			10	7	3		
2	B	1	Total	C	O	0	0
			10	7	3		
2	B	1	Total	C	O	0	0
			10	7	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			10	7	3		
2	B	1	Total	C	O	0	0
			10	7	3		
2	B	1	Total	C	O	0	0
			10	7	3		
2	B	1	Total	C	O	0	0
			10	7	3		

- Molecule 3 is CACODYLATE ION (CCD ID: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	As	C	O	0	0
			5	1	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	36	Total	O	0	0
			36	36		
4	B	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transcriptional regulator TcaR



• Molecule 1: Transcriptional regulator TcaR



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	105.38Å 105.38Å 51.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.45 30.00 – 2.45	Depositor EDS
% Data completeness (in resolution range)	93.7 (30.00-2.45) 93.7 (30.00-2.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.55 (at 2.45Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.234 , 0.271 0.236 , 0.271	Depositor DCC
R_{free} test set	597 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.523	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 63.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.055 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2383	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, SAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	7/1127 (0.6%)	1.18	10/1504 (0.7%)
1	B	0.63	0/1107	1.02	1/1478 (0.1%)
All	All	0.80	7/2234 (0.3%)	1.11	11/2982 (0.4%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	THR	CA-CB	8.84	1.69	1.54
1	A	51	ALA	CA-C	8.65	1.63	1.52
1	A	62	GLY	C-O	8.18	1.28	1.24
1	A	52	LEU	CA-C	8.09	1.63	1.53
1	A	54	VAL	N-CA	7.17	1.56	1.46
1	A	52	LEU	CA-CB	6.56	1.62	1.53
1	A	53	THR	N-CA	5.97	1.53	1.45

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	VAL	N-CA-C	-8.31	104.65	112.96
1	A	51	ALA	CA-C-O	-7.54	112.86	121.15
1	A	3	ARG	N-CA-C	7.52	119.78	109.11
1	A	72	VAL	N-CA-C	-7.38	103.66	110.82
1	A	149	SER	N-CA-C	-7.13	105.51	114.56
1	A	52	LEU	N-CA-C	7.05	117.58	108.34
1	A	144	ILE	N-CA-C	-6.29	104.62	110.53
1	B	130	ILE	N-CA-C	5.72	116.46	110.62
1	A	53	THR	CA-CB-CG2	5.66	120.12	110.50
1	A	123	SER	N-CA-C	5.63	119.25	112.38
1	A	69	SER	N-CA-C	-5.33	106.28	112.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1119	0	1193	58	0
1	B	1099	0	1164	36	0
2	A	20	0	10	3	0
2	B	60	0	35	7	0
3	B	5	0	0	1	0
4	A	36	0	0	4	0
4	B	44	0	0	1	0
All	All	2383	0	2402	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:GLN:H	1:A:56:GLN:NE2	1.59	1.00
1:A:56:GLN:HE21	1:A:56:GLN:N	1.61	0.97
1:A:108:LYS:NZ	4:A:1063:HOH:O	1.98	0.96
1:A:37:SER:H	1:A:40:GLN:NE2	1.70	0.88
1:A:37:SER:H	1:A:40:GLN:HE21	1.15	0.88
1:A:4:ARG:O	1:A:4:ARG:HG3	1.75	0.86
1:A:4:ARG:HH11	1:A:4:ARG:HG2	1.42	0.83
1:A:61:GLN:CG	1:A:64:ASN:ND2	2.43	0.82
1:A:126:ASP:OD2	1:A:128:LYS:HG2	1.80	0.81
1:B:12:LEU:HD13	2:B:3003:SAL:H5	1.61	0.80
1:A:61:GLN:HG3	1:A:64:ASN:ND2	2.02	0.75
1:A:61:GLN:HG2	1:A:64:ASN:ND2	2.03	0.72
1:A:145:GLN:C	1:A:147:TYR:H	1.99	0.71
1:A:16:ILE:HG12	1:B:16:ILE:HD13	1.73	0.70
1:A:42:HIS:HB3	4:A:1031:HOH:O	1.89	0.70
1:A:16:ILE:HG23	1:B:16:ILE:HG23	1.74	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:3005:SAL:H5	4:A:1073:HOH:O	1.93	0.68
1:A:114:MET:HE2	1:A:114:MET:HA	1.75	0.67
1:A:66:ALA:O	4:A:1060:HOH:O	2.12	0.67
1:B:74:LYS:NZ	4:B:1076:HOH:O	2.22	0.67
1:A:61:GLN:CG	1:A:64:ASN:HD22	2.09	0.65
1:B:70:ARG:NH1	3:B:2001:CAC:O1	2.27	0.63
1:A:39:GLU:H	1:A:39:GLU:CD	2.07	0.63
1:A:57:ILE:O	1:A:61:GLN:HB2	1.99	0.62
1:A:61:GLN:HG3	1:A:64:ASN:HD21	1.64	0.62
1:B:64:ASN:HD22	1:B:65:LYS:N	1.99	0.61
1:B:12:LEU:O	1:B:16:ILE:HG12	2.01	0.61
1:A:48:SER:HA	1:A:99:LEU:HD11	1.84	0.60
1:A:143:ARG:O	1:A:147:TYR:HB2	2.02	0.60
1:A:38:ALA:O	1:A:42:HIS:HD2	1.85	0.59
1:B:53:THR:OG1	1:B:56:GLN:HG3	2.02	0.58
1:B:46:MET:HE1	1:B:57:ILE:HA	1.85	0.58
1:A:50:GLU:O	1:A:50:GLU:HG3	2.05	0.57
1:A:47:LEU:HD22	1:A:97:ILE:HG22	1.87	0.56
1:A:69:SER:O	1:A:73:LYS:HG3	2.05	0.56
1:B:104:LYS:HB3	1:B:108:LYS:NZ	2.20	0.56
1:A:145:GLN:HA	1:A:148:THR:HG22	1.88	0.56
1:A:37:SER:OG	1:A:39:GLU:OE1	2.24	0.55
1:A:4:ARG:HG2	1:A:4:ARG:NH1	2.17	0.55
1:A:118:ALA:O	1:A:122:THR:HB	2.07	0.54
1:A:139:ILE:HG21	2:B:3004:SAL:H5	1.89	0.54
1:B:39:GLU:O	1:B:43:VAL:HG23	2.08	0.54
1:A:46:MET:HE1	1:A:57:ILE:HG23	1.90	0.53
1:A:145:GLN:C	1:A:147:TYR:N	2.66	0.53
1:B:5:ILE:HG13	1:B:6:GLU:H	1.73	0.53
1:A:136:VAL:O	1:A:140:ILE:HG13	2.09	0.52
1:A:96:ILE:O	1:A:97:ILE:HD12	2.09	0.52
1:B:114:MET:HA	1:B:114:MET:HE2	1.91	0.52
1:B:117:ILE:HG13	1:B:118:ALA:H	1.75	0.52
1:B:46:MET:HE3	1:B:60:LYS:CD	2.40	0.52
1:B:47:LEU:HD23	1:B:52:LEU:HD12	1.91	0.52
1:B:117:ILE:HG13	1:B:118:ALA:N	2.24	0.51
1:A:16:ILE:HG12	1:B:16:ILE:CD1	2.39	0.51
1:A:130:ILE:HG23	2:B:3003:SAL:O2	2.11	0.51
1:B:46:MET:HE1	1:B:57:ILE:HG23	1.92	0.50
1:A:64:ASN:HB3	1:A:68:VAL:HB	1.92	0.50
1:B:8:HIS:HD1	2:B:3003:SAL:C1'	2.24	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:LEU:HD23	1:B:102:LYS:HB3	1.94	0.50
1:A:14:LYS:NZ	1:B:145:GLN:OE1	2.44	0.50
1:B:43:VAL:CG2	2:B:3008:SAL:H4	2.41	0.50
1:A:81:VAL:HG13	1:A:97:ILE:HG23	1.95	0.49
1:B:5:ILE:HG13	1:B:6:GLU:N	2.28	0.49
1:B:46:MET:HE3	1:B:60:LYS:HD2	1.94	0.48
1:A:99:LEU:HB2	1:A:104:LYS:HG3	1.95	0.48
1:A:61:GLN:HG2	1:A:64:ASN:HD22	1.72	0.48
1:B:53:THR:H	1:B:56:GLN:HE21	1.62	0.47
1:B:58:THR:HG23	1:B:63:VAL:O	2.14	0.47
1:A:96:ILE:C	1:A:97:ILE:HD12	2.41	0.46
1:B:83:LEU:HD23	1:B:97:ILE:CD1	2.45	0.46
1:A:2:VAL:HB	1:A:3:ARG:HD3	1.98	0.46
1:A:82:LYS:HG3	1:A:100:SER:HB3	1.98	0.46
1:B:146:SER:O	1:B:149:SER:HB3	2.16	0.45
1:B:44:LEU:HD21	1:B:75:LEU:HD22	1.98	0.44
1:A:110:ARG:NH2	2:A:3005:SAL:O2'	2.45	0.44
1:B:46:MET:CE	1:B:57:ILE:HG23	2.48	0.44
1:A:6:GLU:HG3	1:A:7:ASP:N	2.32	0.44
1:B:43:VAL:HG22	2:B:3008:SAL:H4	2.00	0.44
1:A:131:GLU:O	1:A:135:GLN:HG3	2.18	0.44
1:A:4:ARG:HH11	1:A:4:ARG:CG	2.19	0.44
1:A:134:ARG:NH1	1:B:8:HIS:HA	2.33	0.44
2:A:3002:SAL:O2'	1:B:134:ARG:HB2	2.18	0.43
1:B:64:ASN:HD22	1:B:65:LYS:H	1.65	0.43
1:A:147:TYR:C	1:A:149:SER:H	2.28	0.42
1:A:50:GLU:HG3	1:A:52:LEU:CD2	2.50	0.42
1:A:134:ARG:HD3	2:B:3003:SAL:O2'	2.21	0.41
1:A:82:LYS:NZ	1:A:100:SER:HB2	2.35	0.41
1:A:50:GLU:HG3	1:A:52:LEU:HD21	2.02	0.41
1:A:145:GLN:O	1:A:147:TYR:N	2.53	0.40
1:B:46:MET:HE3	1:B:60:LYS:HD3	2.03	0.40
1:A:50:GLU:CG	1:A:52:LEU:HD21	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/151 (89%)	125 (93%)	9 (7%)	1 (1%)	18	24
1	B	133/151 (88%)	127 (96%)	6 (4%)	0	100	100
All	All	268/302 (89%)	252 (94%)	15 (6%)	1 (0%)	30	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/139 (91%)	113 (89%)	14 (11%)	6	5
1	B	125/139 (90%)	118 (94%)	7 (6%)	19	27
All	All	252/278 (91%)	231 (92%)	21 (8%)	10	12

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	VAL
1	A	4	ARG
1	A	53	THR
1	A	56	GLN
1	A	63	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	95	LYS
1	A	96	ILE
1	A	97	ILE
1	A	99	LEU
1	A	101	ASN
1	A	109	GLU
1	A	122	THR
1	A	145	GLN
1	A	151	LEU
1	B	9	ILE
1	B	39	GLU
1	B	50	GLU
1	B	81	VAL
1	B	84	GLU
1	B	109	GLU
1	B	151	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	GLN
1	A	42	HIS
1	A	45	ASN
1	A	56	GLN
1	A	64	ASN
1	A	77	ASN
1	A	101	ASN
1	A	145	GLN
1	B	20	ASN
1	B	56	GLN
1	B	64	ASN
1	B	135	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SAL	B	3007	-	10,10,10	2.43	5 (50%)	13,13,13	1.34	3 (23%)
2	SAL	B	3006	-	10,10,10	2.90	8 (80%)	13,13,13	0.97	0
2	SAL	B	3004	-	10,10,10	2.77	7 (70%)	13,13,13	1.21	2 (15%)
2	SAL	B	3008	-	10,10,10	2.37	5 (50%)	13,13,13	1.03	1 (7%)
3	CAC	B	2001	-	2,4,4	9.82	2 (100%)	4,6,6	2.59	2 (50%)
2	SAL	B	3001	-	10,10,10	2.06	5 (50%)	13,13,13	1.18	1 (7%)
2	SAL	A	3005	-	10,10,10	1.34	2 (20%)	13,13,13	0.68	0
2	SAL	B	3003	-	10,10,10	2.15	5 (50%)	13,13,13	1.31	2 (15%)
2	SAL	A	3002	-	10,10,10	1.35	1 (10%)	13,13,13	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAL	B	3007	-	-	0/4/4/4	0/1/1/1
2	SAL	B	3006	-	-	0/4/4/4	0/1/1/1
2	SAL	B	3004	-	-	0/4/4/4	0/1/1/1
2	SAL	B	3008	-	-	0/4/4/4	0/1/1/1
2	SAL	B	3001	-	-	0/4/4/4	0/1/1/1
2	SAL	A	3005	-	-	0/4/4/4	0/1/1/1
2	SAL	B	3003	-	-	0/4/4/4	0/1/1/1
2	SAL	A	3002	-	-	0/4/4/4	0/1/1/1

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2001	CAC	AS-C1	10.65	2.15	1.90
3	B	2001	CAC	AS-C2	8.91	2.11	1.90
2	B	3006	SAL	O2'-C1'	4.97	1.45	1.30
2	B	3008	SAL	O2'-C1'	4.09	1.42	1.30
2	B	3004	SAL	C6-C1	3.97	1.46	1.39
2	B	3004	SAL	C4-C3	3.95	1.45	1.38
2	B	3006	SAL	C4-C3	3.76	1.45	1.38
2	B	3006	SAL	C6-C1	3.75	1.45	1.39
2	B	3007	SAL	C4-C3	3.69	1.45	1.38
2	B	3004	SAL	C5-C4	3.66	1.46	1.38
2	B	3004	SAL	C3-C2	3.66	1.45	1.39
2	B	3007	SAL	O2'-C1'	3.65	1.41	1.30
2	B	3003	SAL	C3-C2	3.43	1.45	1.39
2	B	3007	SAL	C5-C4	3.30	1.45	1.38
2	B	3008	SAL	C6-C1	3.27	1.44	1.39
2	B	3001	SAL	C1-C2	2.96	1.45	1.40
2	A	3005	SAL	O2'-C1'	2.93	1.39	1.30
2	B	3001	SAL	O2'-C1'	2.93	1.39	1.30
2	B	3003	SAL	C5-C4	2.85	1.44	1.38
2	B	3008	SAL	O1'-C1'	2.82	1.30	1.22
2	B	3008	SAL	C5-C4	2.73	1.44	1.38
2	B	3006	SAL	C1-C2	2.65	1.45	1.40
2	B	3006	SAL	O2-C2	2.64	1.41	1.36
2	B	3003	SAL	C5-C6	2.62	1.43	1.38
2	B	3001	SAL	C4-C3	2.52	1.43	1.38
2	B	3003	SAL	O2'-C1'	2.45	1.37	1.30
2	B	3001	SAL	C5-C4	2.44	1.43	1.38
2	B	3004	SAL	O2-C2	2.34	1.41	1.36
2	B	3007	SAL	C6-C1	2.30	1.43	1.39
2	B	3006	SAL	C5-C6	2.20	1.42	1.38
2	B	3006	SAL	O1'-C1'	2.19	1.29	1.22
2	B	3004	SAL	C5-C6	2.16	1.42	1.38
2	B	3008	SAL	C5-C6	2.15	1.42	1.38
2	B	3007	SAL	C5-C6	2.15	1.42	1.38
2	B	3006	SAL	C5-C4	2.13	1.42	1.38
2	A	3002	SAL	C5-C4	2.12	1.42	1.38
2	B	3003	SAL	C4-C3	2.08	1.42	1.38
2	B	3001	SAL	C1-C1'	-2.08	1.45	1.49
2	B	3004	SAL	O2'-C1'	2.06	1.36	1.30
2	A	3005	SAL	C6-C1	2.04	1.43	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2001	CAC	O2-AS-C1	4.21	116.03	105.84
2	B	3003	SAL	O2-C2-C1	-3.17	115.83	121.71
2	B	3007	SAL	C6-C1-C2	2.64	121.47	118.15
2	B	3001	SAL	C6-C1-C2	2.45	121.22	118.15
2	B	3004	SAL	C6-C1-C2	2.22	120.94	118.15
2	B	3004	SAL	O2-C2-C1	-2.18	117.67	121.71
2	B	3007	SAL	O2'-C1'-C1	2.18	121.46	115.28
3	B	2001	CAC	O1-AS-C2	2.17	114.20	111.50
2	B	3003	SAL	C6-C1-C2	2.13	120.82	118.15
2	B	3007	SAL	O1'-C1'-C1	-2.11	116.93	121.97
2	B	3008	SAL	O2-C2-C1	-2.04	117.92	121.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3004	SAL	1	0
2	B	3008	SAL	2	0
3	B	2001	CAC	1	0
2	A	3005	SAL	2	0
2	B	3003	SAL	4	0
2	A	3002	SAL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	139/151 (92%)	0.71	21 (15%) 5 4	31, 60, 118, 123	0
1	B	137/151 (90%)	0.17	2 (1%) 72 74	25, 53, 73, 94	0
All	All	276/302 (91%)	0.44	23 (8%) 17 15	25, 55, 107, 123	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	63	VAL	7.4
1	A	57	ILE	5.7
1	A	64	ASN	4.6
1	A	66	ALA	3.6
1	A	60	LYS	3.5
1	A	83	LEU	3.3
1	A	68	VAL	3.2
1	A	2	VAL	3.0
1	A	67	ALA	3.0
1	B	5	ILE	2.9
1	A	96	ILE	2.6
1	A	61	GLN	2.6
1	A	58	THR	2.5
1	A	5	ILE	2.4
1	A	147	TYR	2.4
1	A	56	GLN	2.4
1	B	151	LEU	2.4
1	A	72	VAL	2.3
1	A	69	SER	2.2
1	A	81	VAL	2.2
1	A	103	GLY	2.2
1	A	70	ARG	2.1
1	A	80	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SAL	B	3008	10/10	0.72	0.16	53,54,57,58	0
2	SAL	B	3006	10/10	0.76	0.13	60,60,61,62	0
2	SAL	B	3007	10/10	0.86	0.11	51,52,52,56	0
2	SAL	B	3003	10/10	0.87	0.14	53,54,56,58	0
2	SAL	A	3002	10/10	0.89	0.17	37,38,40,45	0
2	SAL	B	3004	10/10	0.90	0.11	49,52,54,55	0
2	SAL	B	3001	10/10	0.90	0.09	38,40,43,44	0
2	SAL	A	3005	10/10	0.93	0.08	46,49,50,50	0
3	CAC	B	2001	5/5	0.93	0.18	89,89,90,91	0

6.5 Other polymers [i](#)

There are no such residues in this entry.