



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:37 AM UTC

PDB ID : 5KPI / pdb_00005kpi
Title : Mouse native PGP
Authors : Xia, D.; Esser, L.; Zhou, F.
Deposited on : 2016-07-04
Resolution : 4.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1095 (4.24-3.80)
Clashscore	190562	1142 (4.24-3.80)
Ramachandran outliers	187476	1074 (4.24-3.80)
Sidechain outliers	187428	1065 (4.24-3.80)
RSRZ outliers	180081	1095 (4.24-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1282	 88% 5% 8%
1	B	1282	 88% 5% 8%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 37077 atoms, of which 18703 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug resistance protein 1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	1185	Total	C	H	N	O	S	0	0	0
			18542	5907	9353	1557	1687	38			
1	B	1184	Total	C	H	N	O	S	0	0	0
			18535	5905	9350	1556	1686	38			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1277	HIS	-	expression tag	UNP P21447
A	1278	HIS	-	expression tag	UNP P21447
A	1279	HIS	-	expression tag	UNP P21447
A	1280	HIS	-	expression tag	UNP P21447
A	1281	HIS	-	expression tag	UNP P21447
A	1282	HIS	-	expression tag	UNP P21447
B	1277	HIS	-	expression tag	UNP P21447
B	1278	HIS	-	expression tag	UNP P21447
B	1279	HIS	-	expression tag	UNP P21447
B	1280	HIS	-	expression tag	UNP P21447
B	1281	HIS	-	expression tag	UNP P21447
B	1282	HIS	-	expression tag	UNP P21447

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	100.61Å 119.53Å 383.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.14 – 4.01 20.14 – 4.01	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.14-4.01) 94.0 (20.14-4.01)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 3.99Å)	Xtriage
Refinement program	PHENIX (dev_2443: ???)	Depositor
R, R_{free}	0.258 , 0.303 0.276 , 0.328	Depositor DCC
R_{free} test set	1870 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	184.9	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 158.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	37077	wwPDB-VP
Average B, all atoms (Å ²)	262.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/9358	0.63	2/12650 (0.0%)
1	B	0.30	0/9354	0.63	0/12645
All	All	0.30	0/18712	0.63	2/25295 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1159	ASP	CA-C-N	5.59	132.22	121.54
1	A	1159	ASP	C-N-CA	5.59	132.22	121.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1159	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9189	9353	9370	17	1
1	B	9185	9350	9367	19	1
All	All	18374	18703	18737	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:695:ARG:NH2	1:A:1009:GLU:OE2	2.26	0.68
1:A:471:GLN:NE2	1:A:552:GLU:OE1	2.36	0.59
1:B:514:HIS:O	1:B:518:THR:OG1	2.14	0.58
1:B:278:GLU:OE1	1:B:782:LYS:NZ	2.38	0.56
1:A:801:ASP:OD2	1:A:1083:TYR:OH	2.24	0.56

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:391:LYS:NZ	1:B:1023:LYS:O[1_565]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1181/1282 (92%)	1121 (95%)	55 (5%)	5 (0%)	30	65
1	B	1180/1282 (92%)	1122 (95%)	53 (4%)	5 (0%)	30	65
All	All	2361/2564 (92%)	2243 (95%)	108 (5%)	10 (0%)	30	65

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	515	GLN
1	A	1160	LYS
1	B	515	GLN
1	B	1160	LYS
1	A	1120	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	977/1063 (92%)	950 (97%)	27 (3%)	38	60
1	B	977/1063 (92%)	954 (98%)	23 (2%)	43	63
All	All	1954/2126 (92%)	1904 (97%)	50 (3%)	40	61

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	121	VAL
1	B	385	GLN
1	B	1200	SER
1	B	134	LEU
1	B	306	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1050	GLN
1	B	1176	GLN
1	A	1132	ASN
1	A	1149	ASN
1	B	471	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	1185/1282 (92%)	-0.44	5 (0%)	88 75	155, 250, 354, 546	0
1	B	1184/1282 (92%)	-0.46	2 (0%)	91 81	150, 257, 391, 647	0
All	All	2369/2564 (92%)	-0.45	7 (0%)	90 78	150, 253, 373, 647	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	553	ALA	3.7
1	B	1021	GLY	2.4
1	B	205	THR	2.4
1	A	312	TYR	2.3
1	A	69	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.