



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:17 AM UTC

PDB ID : 4KRP / pdb_00004krp
Title : Nanobody/VHH domain 9G8 in complex with the extracellular region of EGFR
Authors : Ferguson, K.M.; Schmitz, K.R.
Deposited on : 2013-05-16
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	625	20% 78% 8% 15%
2	B	136	22% 79% 6% 15%
3	C	211	89% 10%
4	D	220	2% 88% 9% .
5	E	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	2	 100%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14877 atoms, of which 7017 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	534	Total	C	H	N	O	S	0	0	0
			7001	2341	3252	651	712	45			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	614	THR	-	expression tag	UNP P00533
A	620	HIS	-	expression tag	UNP P00533
A	621	HIS	-	expression tag	UNP P00533
A	622	HIS	-	expression tag	UNP P00533
A	623	HIS	-	expression tag	UNP P00533
A	624	HIS	-	expression tag	UNP P00533
A	625	HIS	-	expression tag	UNP P00533

- Molecule 2 is a protein called Nanobody/VHH domain 9G8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	116	Total	C	H	N	O	S	0	0	0
			1473	516	655	135	163	4			

- Molecule 3 is a protein called Cetuximab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	211	Total	C	H	N	O	S	0	0	0
			3099	992	1506	270	327	4			

- Molecule 4 is a protein called Cetuximab heavy chain.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	213	Total	C	H	N	O	S	0	0	0
			3079	1007	1494	259	314	5			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	2	Total	C	H	N	O	0	0	0
			55	16	27	2	10			
5	F	2	Total	C	H	N	O	0	0	0
			55	16	27	2	10			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
6	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total 1	O 1	0	0
7	C	1	Total 1	O 1	0	0
7	D	1	Total 1	O 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



- Molecule 2: Nanobody/VHH domain 9G8




- Molecule 3: Cetuximab light chain

Chain C:  89% 10%



- Molecule 4: Cetuximab heavy chain

Chain D:  2% 88% 9%




- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.37Å 95.84Å 129.48Å 90.00° 99.95° 90.00°	Depositor
Resolution (Å)	44.46 – 2.82 44.46 – 2.82	Depositor EDS
% Data completeness (in resolution range)	98.0 (44.46-2.82) 98.0 (44.46-2.82)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.27 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.215 , 0.263 0.221 , 0.266	Depositor DCC
R_{free} test set	1889 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	51.9	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	14877	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/3825	0.69	1/5230 (0.0%)
2	B	0.28	0/838	0.61	0/1146
3	C	0.41	0/1627	0.68	0/2218
4	D	0.42	0/1627	0.72	0/2233
All	All	0.37	0/7917	0.69	1/10827 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	417	VAL	N-CA-C	5.13	115.36	107.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3749	3252	3319	23	0
2	B	818	655	655	4	0
3	C	1593	1506	1503	14	0
4	D	1585	1494	1492	9	0
5	E	28	27	25	0	0
5	F	28	27	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	42	42	39	1	0
6	D	14	14	13	0	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
All	All	7860	7017	7071	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:337:ASN:HD22	6:A:703:NAG:H83	1.60	0.67
1:A:376:GLU:OE1	1:A:403:ARG:NH1	2.28	0.66
1:A:455:LYS:O	2:B:27:ARG:NH2	2.31	0.64
1:A:403:ARG:NH2	2:B:100:TYR:O	2.31	0.64
2:B:54:SER:O	2:B:104:SER:OG	2.18	0.61
3:C:80:SER:HA	3:C:106:LEU:HD22	1.83	0.61
4:D:32:TYR:CZ	4:D:101:TYR:CE1	2.90	0.59
1:A:14:LEU:HG	1:A:89:TYR:OH	2.02	0.58
1:A:245:LEU:HD13	1:A:577:GLY:O	2.05	0.56
3:C:164:THR:HG22	3:C:174:SER:H	1.72	0.55
1:A:276:VAL:N	1:A:284:VAL:O	2.40	0.53
1:A:28:GLN:HG3	1:A:29:ARG:N	2.23	0.53
2:B:101:GLN:OE1	2:B:107:TYR:OH	2.26	0.51
3:C:13:VAL:HG13	3:C:17:GLU:HB2	1.94	0.50
3:C:158:ASN:N	3:C:158:ASN:OD1	2.46	0.49
4:D:6:GLN:O	4:D:111:GLN:NE2	2.46	0.49
1:A:588:ASP:O	1:A:591:HIS:N	2.48	0.47
1:A:407:LYS:NZ	1:A:434:ASP:OD2	2.39	0.47
1:A:447:TYR:O	1:A:449:ASN:N	2.43	0.47
1:A:365:PRO:HB3	1:A:387:PRO:HG3	1.97	0.46
3:C:140:TYR:CG	3:C:141:PRO:HA	2.50	0.46
3:C:182:SER:OG	3:C:185:ASP:OD2	2.32	0.46
1:A:41:LEU:HB3	1:A:65:VAL:HG22	1.99	0.45
1:A:272:PRO:HG2	1:A:275:TYR:CG	2.51	0.45
3:C:124:GLN:O	3:C:127:SER:OG	2.32	0.45
4:D:47:TRP:HZ2	4:D:50:VAL:HG23	1.82	0.44
3:C:44:PRO:HG2	4:D:109:TRP:CE2	2.53	0.44
1:A:584:TRP:NE1	1:A:601:THR:O	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:72:ASP:OD1	4:D:74:SER:OG	2.28	0.44
3:C:19:VAL:HG23	3:C:78:VAL:CG2	2.48	0.44
1:A:443:LYS:HE3	3:C:94:TRP:CZ2	2.53	0.43
1:A:268:VAL:HG12	1:A:270:LYS:H	1.83	0.43
1:A:208:CYS:N	1:A:216:CYS:SG	2.91	0.43
4:D:207:LYS:N	4:D:208:PRO:HD2	2.34	0.42
3:C:37:GLN:O	3:C:45:ARG:N	2.51	0.42
4:D:4:LEU:HD12	4:D:22:CYS:SG	2.59	0.41
1:A:326:SER:OG	1:A:327:ILE:N	2.52	0.41
3:C:108:ARG:HG3	3:C:109:THR:O	2.21	0.41
4:D:105:GLU:O	4:D:105:GLU:HG2	2.21	0.41
5:F:1:NAG:O3	5:F:2:NAG:O5	2.30	0.41
1:A:246:TYR:OH	1:A:563:ASP:OD2	2.27	0.41
3:C:13:VAL:HG11	3:C:78:VAL:HG21	2.02	0.41
4:D:105:GLU:OE1	4:D:105:GLU:N	2.43	0.41
1:A:427:ARG:HA	1:A:492:TRP:CD1	2.56	0.41
1:A:272:PRO:HG2	1:A:275:TYR:CD2	2.56	0.40
1:A:97:VAL:HB	1:A:126:PHE:CE2	2.57	0.40
3:C:19:VAL:CG2	3:C:78:VAL:HG21	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/625 (85%)	478 (90%)	51 (10%)	1 (0%)	43	71
2	B	112/136 (82%)	96 (86%)	16 (14%)	0	100	100
3	C	209/211 (99%)	204 (98%)	4 (2%)	1 (0%)	24	53
4	D	209/220 (95%)	198 (95%)	11 (5%)	0	100	100
All	All	1060/1192 (89%)	976 (92%)	82 (8%)	2 (0%)	43	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	214	ALA
3	C	138	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/546 (67%)	362 (98%)	6 (2%)	55	82
2	B	66/110 (60%)	64 (97%)	2 (3%)	36	69
3	C	179/188 (95%)	177 (99%)	2 (1%)	65	86
4	D	176/190 (93%)	169 (96%)	7 (4%)	28	61
All	All	789/1034 (76%)	772 (98%)	17 (2%)	45	77

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	THR
1	A	278	THR
1	A	391	THR
1	A	505	VAL
1	A	568	VAL
1	A	604	CYS
2	B	49	VAL
2	B	53	TRP
3	C	106	LEU
3	C	181	LEU
4	D	24	VAL
4	D	25	SER
4	D	30	THR
4	D	50	VAL
4	D	51	ILE
4	D	81	LYS
4	D	134	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	334	HIS
1	A	480	GLN
1	A	554	ASN
2	B	77	ASN
3	C	137	ASN
3	C	138	ASN
4	D	170	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	E	1	1,5	14,14,15	0.68	1 (7%)	17,19,21	0.65	0
5	NAG	E	2	5	14,14,15	0.54	0	17,19,21	1.15	0
5	NAG	F	1	1,5	14,14,15	0.52	0	17,19,21	1.15	0
5	NAG	F	2	5	14,14,15	0.55	0	17,19,21	1.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	1/6/23/26	0/1/1/1
5	NAG	F	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	F	2	5	-	3/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	1	NAG	O5-C1	-2.25	1.39	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	F	2	NAG	C8-C7-N2-C2
5	F	2	NAG	O7-C7-N2-C2
5	E	1	NAG	C4-C5-C6-O6
5	E	1	NAG	O5-C5-C6-O6
5	F	2	NAG	O5-C5-C6-O6
5	F	1	NAG	O5-C5-C6-O6
5	E	2	NAG	C1-C2-N2-C7


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	2	NAG	1	0
5	F	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	A	704	1	14,14,15	0.53	0	17,19,21	1.13	0
6	NAG	A	703	1	14,14,15	0.31	0	17,19,21	0.57	0
6	NAG	D	301	4	14,14,15	0.55	0	17,19,21	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	A	707	1	14,14,15	0.28	0	17,19,21	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	704	1	-	3/6/23/26	0/1/1/1
6	NAG	A	703	1	-	3/6/23/26	0/1/1/1
6	NAG	D	301	4	-	2/6/23/26	0/1/1/1
6	NAG	A	707	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	703	NAG	C8-C7-N2-C2
6	A	703	NAG	O7-C7-N2-C2
6	A	704	NAG	C8-C7-N2-C2
6	A	704	NAG	O7-C7-N2-C2
6	A	707	NAG	C8-C7-N2-C2
6	A	707	NAG	O7-C7-N2-C2
6	D	301	NAG	C8-C7-N2-C2
6	D	301	NAG	O7-C7-N2-C2
6	A	703	NAG	O5-C5-C6-O6
6	A	704	NAG	O5-C5-C6-O6
6	A	707	NAG	C4-C5-C6-O6
6	A	707	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	703	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/625 (85%)	0.99	126 (23%) 2 1	27, 92, 160, 179	0
2	B	116/136 (85%)	1.39	30 (25%) 1 1	62, 102, 132, 151	0
3	C	211/211 (100%)	-0.33	0 100 100	19, 38, 60, 79	0
4	D	213/220 (96%)	-0.24	5 (2%) 61 51	19, 37, 56, 92	0
All	All	1074/1192 (90%)	0.53	161 (14%) 5 4	19, 56, 153, 179	0

All (161) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	THR	7.1
1	A	81	GLN	6.5
1	A	104	ASN	6.4
1	A	82	ILE	5.7
1	A	211	GLN	4.9
1	A	103	ALA	4.5
1	A	126	PHE	4.5
1	A	107	GLY	4.5
1	A	587	ALA	4.5
1	A	38	LEU	4.3
1	A	110	GLU	4.3
1	A	101	TYR	4.3
2	B	82	GLN	4.2
2	B	11	LEU	4.1
1	A	74	ARG	4.1
1	A	116	LEU	4.0
1	A	274	ASN	4.0
1	A	47	GLN	4.0
1	A	24	PHE	4.0
1	A	238	ASP	4.0
1	A	20	PHE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	51	ASP	3.9
1	A	290	ASP	3.9
1	A	50	TYR	3.9
1	A	115	ASN	3.9
1	A	105	LYS	3.8
1	A	78	GLU	3.7
1	A	214	ALA	3.7
1	A	54	PHE	3.7
1	A	59	GLN	3.7
2	B	84	ASN	3.7
1	A	49	ASN	3.7
1	A	210	ASN	3.7
1	A	296	GLU	3.6
1	A	125	ARG	3.6
1	A	36	VAL	3.6
2	B	83	MET	3.5
1	A	219	PRO	3.5
2	B	90	ASP	3.5
1	A	234	ALA	3.5
1	A	279	ASP	3.4
2	B	67	ARG	3.3
1	A	213	ALA	3.3
1	A	225	LEU	3.2
1	A	132	LEU	3.2
1	A	586	TYR	3.1
1	A	579	ASN	3.1
1	A	605	THR	3.1
1	A	215	GLY	3.1
1	A	58	ILE	3.1
1	A	123	ALA	3.0
4	D	218	GLU	3.0
1	A	91	ASN	3.0
1	A	90	GLU	2.9
1	A	607	PRO	2.9
1	A	228	ARG	2.9
1	A	37	VAL	2.9
1	A	612	CYS	2.9
2	B	96	CYS	2.9
1	A	80	LEU	2.9
1	A	317	GLY	2.9
1	A	300	ARG	2.9
1	A	48	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	232	ASP	2.8
1	A	108	LEU	2.8
1	A	474	SER	2.8
2	B	69	THR	2.8
1	A	578	GLU	2.8
2	B	91	THR	2.8
2	B	35	GLY	2.8
2	B	47	PHE	2.8
2	B	97	ALA	2.7
1	A	252	GLN	2.7
1	A	359	HIS	2.7
1	A	84	ARG	2.7
2	B	55	SER	2.7
1	A	21	GLU	2.7
1	A	79	ASN	2.7
2	B	73	ASP	2.7
1	A	592	VAL	2.7
2	B	124	THR	2.7
4	D	41	PRO	2.6
2	B	109	PHE	2.6
1	A	217	THR	2.6
1	A	86	ASN	2.6
2	B	89	GLU	2.6
1	A	35	GLU	2.6
1	A	129	ASN	2.6
2	B	68	PHE	2.6
1	A	297	ASP	2.6
1	A	291	SER	2.5
1	A	294	MET	2.5
1	A	71	THR	2.5
1	A	208	CYS	2.5
1	A	55	LEU	2.5
1	A	111	LEU	2.5
1	A	226	VAL	2.5
1	A	299	VAL	2.5
1	A	561	TYR	2.5
2	B	74	ASN	2.5
1	A	19	THR	2.4
1	A	100	ASN	2.4
1	A	96	ALA	2.4
2	B	94	TYR	2.4
1	A	588	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	14	LEU	2.4
1	A	83	ILE	2.4
1	A	119	ILE	2.4
1	A	233	GLU	2.4
1	A	113	MET	2.4
1	A	574	GLY	2.3
1	A	117	GLN	2.3
2	B	54	SER	2.3
1	A	602	TYR	2.3
1	A	109	LYS	2.3
1	A	124	VAL	2.3
1	A	128	ASN	2.3
2	B	42	GLY	2.3
1	A	102	ASP	2.3
1	A	67	ILE	2.3
1	A	114	ARG	2.3
1	A	598	PRO	2.3
1	A	220	ARG	2.3
1	A	32	ASN	2.3
1	A	66	LEU	2.3
2	B	20	LEU	2.3
4	D	43	LYS	2.3
1	A	250	THR	2.3
1	A	130	PRO	2.3
2	B	80	TYR	2.3
1	A	603	GLY	2.3
1	A	589	ALA	2.2
1	A	131	ALA	2.2
1	A	56	LYS	2.2
1	A	611	GLY	2.2
1	A	127	SER	2.2
1	A	8	GLN	2.1
2	B	22	CYS	2.1
2	B	45	ARG	2.1
1	A	562	ILE	2.1
1	A	512	VAL	2.1
1	A	264	GLY	2.1
2	B	66	GLY	2.1
2	B	30	SER	2.1
1	A	254	ASP	2.1
1	A	285	ARG	2.1
1	A	43	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	306	GLU	2.1
1	A	295	GLU	2.1
1	A	53	SER	2.0
2	B	3	GLN	2.0
4	D	135	LYS	2.0
1	A	61	VAL	2.0
1	A	112	PRO	2.0
1	A	590	GLY	2.0
1	A	44	THR	2.0
1	A	269	LYS	2.0
2	B	44	GLU	2.0
1	A	93	TYR	2.0
4	D	32	TYR	2.0
1	A	97	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	E	2	14/15	0.67	0.16	66,92,111,114	0
5	NAG	F	2	14/15	0.76	0.13	69,89,105,118	0
5	NAG	F	1	14/15	0.87	0.10	43,58,70,77	0
5	NAG	E	1	14/15	0.89	0.10	53,79,96,104	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	A	704	14/15	0.41	0.20	71,97,116,121	0
6	NAG	D	301	14/15	0.69	0.16	64,83,97,112	0
6	NAG	A	707	14/15	0.76	0.12	60,85,102,106	0
6	NAG	A	703	14/15	0.76	0.11	59,95,116,120	0

6.5 Other polymers [i](#)

There are no such residues in this entry.