



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:03 PM UTC

PDB ID : 4KRX / pdb_00004krx
Title : Structure of Aes from E. coli
Authors : Schiefner, A.; Gerber, K.; Brosig, A.; Boos, W.
Deposited on : 2013-05-17
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	333	3% 86% 9% 5%
1	B	333	85% 10% 5%
1	C	333	3% 86% 8% 6%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl esterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	5	0
			2546	1622	429	476	19			
1	B	316	Total	C	N	O	S	0	4	0
			2542	1620	430	473	19			
1	C	313	Total	C	N	O	S	0	7	0
			2532	1615	427	471	19			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	expression tag	UNP P23872
A	-12	ARG	-	expression tag	UNP P23872
A	-11	GLY	-	expression tag	UNP P23872
A	-10	SER	-	expression tag	UNP P23872
A	-9	HIS	-	expression tag	UNP P23872
A	-8	HIS	-	expression tag	UNP P23872
A	-7	HIS	-	expression tag	UNP P23872
A	-6	HIS	-	expression tag	UNP P23872
A	-5	HIS	-	expression tag	UNP P23872
A	-4	HIS	-	expression tag	UNP P23872
A	-3	THR	-	expression tag	UNP P23872
A	-2	ASP	-	expression tag	UNP P23872
A	-1	PRO	-	expression tag	UNP P23872
A	0	ILE	-	expression tag	UNP P23872
B	-13	MET	-	expression tag	UNP P23872
B	-12	ARG	-	expression tag	UNP P23872
B	-11	GLY	-	expression tag	UNP P23872
B	-10	SER	-	expression tag	UNP P23872
B	-9	HIS	-	expression tag	UNP P23872
B	-8	HIS	-	expression tag	UNP P23872
B	-7	HIS	-	expression tag	UNP P23872
B	-6	HIS	-	expression tag	UNP P23872
B	-5	HIS	-	expression tag	UNP P23872

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	HIS	-	expression tag	UNP P23872
B	-3	THR	-	expression tag	UNP P23872
B	-2	ASP	-	expression tag	UNP P23872
B	-1	PRO	-	expression tag	UNP P23872
B	0	ILE	-	expression tag	UNP P23872
C	-13	MET	-	expression tag	UNP P23872
C	-12	ARG	-	expression tag	UNP P23872
C	-11	GLY	-	expression tag	UNP P23872
C	-10	SER	-	expression tag	UNP P23872
C	-9	HIS	-	expression tag	UNP P23872
C	-8	HIS	-	expression tag	UNP P23872
C	-7	HIS	-	expression tag	UNP P23872
C	-6	HIS	-	expression tag	UNP P23872
C	-5	HIS	-	expression tag	UNP P23872
C	-4	HIS	-	expression tag	UNP P23872
C	-3	THR	-	expression tag	UNP P23872
C	-2	ASP	-	expression tag	UNP P23872
C	-1	PRO	-	expression tag	UNP P23872
C	0	ILE	-	expression tag	UNP P23872

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	8	5		
2	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	250	Total 250	O 250	0	0
3	B	222	Total 222	O 222	0	0
3	C	186	Total 186	O 186	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl esterase



- Molecule 1: Acetyl esterase



- Molecule 1: Acetyl esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	113.71Å 113.71Å 151.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.80 30.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-1.80) 99.3 (30.00-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 1.80Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.161 , 0.190 0.173 , 0.199	Depositor DCC
R_{free} test set	2083 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8304	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2502e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.33	11/2614 (0.4%)	1.14	1/3555 (0.0%)
1	B	1.31	10/2610 (0.4%)	1.08	1/3549 (0.0%)
1	C	1.23	4/2606 (0.2%)	1.12	6/3545 (0.2%)
All	All	1.29	25/7830 (0.3%)	1.11	8/10649 (0.1%)

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	274	THR	C-O	8.55	1.33	1.24
1	A	168	ALA	C-O	7.66	1.33	1.24
1	A	270	LEU	C-O	7.63	1.33	1.24
1	B	270	LEU	C-O	7.50	1.32	1.24
1	B	282	CYS	N-CA	7.28	1.54	1.46
1	B	271	LEU	N-CA	7.06	1.54	1.46
1	A	306	GLU	C-O	6.53	1.31	1.24
1	B	111	ALA	N-CA	6.33	1.54	1.46
1	A	274	THR	C-O	6.28	1.31	1.24
1	A	240	TYR	N-CA	6.27	1.53	1.46
1	A	275	LEU	C-O	6.14	1.31	1.24
1	C	262	ASP	CA-C	6.07	1.57	1.52
1	C	253	PRO	N-CA	-5.94	1.40	1.47
1	B	116	CYS	CA-C	-5.77	1.46	1.52
1	A	22	ASN	N-CA	5.46	1.53	1.46
1	B	306	GLU	C-O	5.42	1.30	1.24
1	A	136	GLU	N-CA	5.40	1.52	1.46
1	A	261	PHE	CA-CB	5.35	1.60	1.53
1	C	110	LEU	C-O	5.35	1.30	1.24
1	B	269[A]	ARG	C-O	5.26	1.30	1.24
1	B	269[B]	ARG	C-O	5.26	1.30	1.24
1	C	286	LEU	CA-C	-5.18	1.46	1.52
1	B	261	PHE	CA-CB	5.13	1.60	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	102	THR	C-O	-5.09	1.17	1.24
1	A	290	THR	CA-CB	5.04	1.61	1.53

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	252	PRO	O-C-N	7.53	124.77	121.31
1	A	20	VAL	N-CA-C	6.57	116.73	110.42
1	C	300	MET	CG-SD-CE	-5.74	88.27	100.90
1	B	298	SER	N-CA-C	5.70	118.27	111.71
1	C	20	VAL	N-CA-C	5.56	115.76	110.42
1	C	217	THR	N-CA-C	-5.43	103.73	110.41
1	C	252	PRO	CA-C-N	5.26	125.52	119.83
1	C	252	PRO	C-N-CA	5.26	125.52	119.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2546	0	2458	16	0
1	B	2542	0	2459	11	0
1	C	2532	0	2453	11	0
2	A	13	0	18	2	0
2	B	13	0	18	1	0
3	A	250	0	0	4	0
3	B	222	0	0	2	0
3	C	186	0	0	1	0
All	All	8304	0	7406	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:61:ARG:HD2	1:B:63:TYR:OH	1.92	0.69
1:B:19:THR:HG22	1:B:300:MET:HE1	1.74	0.69
1:A:302:LYS:NZ	3:A:675:HOH:O	2.26	0.67
1:A:16:GLU:HG2	1:A:301:MET:HE3	1.79	0.65
1:A:4:GLU:N	1:A:4:GLU:OE1	2.30	0.65
1:A:16:GLU:HB3	3:A:539:HOH:O	1.98	0.62
1:B:177:TRP:CE2	1:B:181:LYS:HE3	2.35	0.61
1:A:76:LEU:HG	1:A:119:ILE:HG12	1.84	0.60
1:B:4:GLU:HG2	1:B:5:ASN:H	1.67	0.60
1:A:16:GLU:HG3	1:A:301:MET:HG2	1.88	0.55
1:A:157[B]:SER:O	1:A:158:ARG:HG3	2.07	0.55
1:C:16:GLU:HG2	3:C:432:HOH:O	2.07	0.54
1:C:144:TYR:HH	1:C:152:TYR:HH	1.54	0.54
1:C:76:LEU:HG	1:C:119:ILE:HG12	1.92	0.52
1:B:61:ARG:NH2	3:B:679:HOH:O	2.42	0.51
1:A:166:ALA:HB2	2:A:401:PG4:H21	1.92	0.50
1:A:300:MET:HB2	3:A:551:HOH:O	2.13	0.47
1:C:195:TYR:CZ	1:C:293:ALA:HA	2.49	0.47
1:A:76:LEU:N	1:A:76:LEU:HD12	2.29	0.47
1:C:197:LEU:H	1:C:197:LEU:HD23	1.80	0.47
1:A:96:ILE:HG13	1:A:97:LEU:HG	1.96	0.47
1:A:187:LYS:NZ	3:A:746:HOH:O	2.43	0.47
1:B:4:GLU:HG2	3:B:664:HOH:O	2.14	0.47
1:C:264:LEU:HD12	1:C:292:HIS:CE1	2.51	0.45
1:C:7:LEU:HD12	1:C:209:LEU:HD21	1.98	0.45
1:B:76:LEU:HG	1:B:119:ILE:HG12	1.99	0.44
1:B:195:TYR:CZ	1:B:293:ALA:HA	2.53	0.44
1:C:195:TYR:CE1	1:C:293:ALA:HA	2.52	0.44
1:A:110:LEU:HD23	1:A:110:LEU:HA	1.87	0.43
1:A:86:THR:O	1:A:159:ILE:HA	2.18	0.43
1:C:58:MET:HE3	1:C:58:MET:HB2	1.96	0.43
1:B:199:GLY:HA3	1:B:241:CYS:HA	2.02	0.42
1:B:13:ILE:HG13	1:B:18:LYS:HG3	2.02	0.41
1:A:94:GLY:HA2	2:A:401:PG4:H21	2.01	0.41
1:A:206:ARG:HH11	1:A:206:ARG:HD3	1.76	0.41
1:C:126:SER:HB3	1:C:127:PRO:HA	2.02	0.40
1:B:216:LEU:HD21	2:B:401:PG4:H41	2.02	0.40
1:C:14:SER:HB3	1:C:289:GLY:HA3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/333 (96%)	311 (98%)	8 (2%)	0	100	100
1	B	318/333 (96%)	307 (96%)	11 (4%)	0	100	100
1	C	318/333 (96%)	308 (97%)	10 (3%)	0	100	100
All	All	955/999 (96%)	926 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/279 (96%)	265 (99%)	3 (1%)	65	60
1	B	267/279 (96%)	262 (98%)	5 (2%)	50	41
1	C	267/279 (96%)	264 (99%)	3 (1%)	65	60
All	All	802/837 (96%)	791 (99%)	11 (1%)	61	52

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	16	GLU
1	A	34	THR
1	B	16	GLU
1	B	90	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	158	ARG
1	B	219	GLN
1	B	222	GLN
1	C	61[A]	ARG
1	C	61[B]	ARG
1	C	65	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	148	GLN
1	A	218	GLN
1	A	313	GLN
1	C	40	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PG4	A	401	-	12,12,12	0.68	0	11,11,11	0.64	0
2	PG4	B	401	-	12,12,12	0.66	0	11,11,11	0.80	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	A	401	-	-	5/10/10/10	-
2	PG4	B	401	-	-	5/10/10/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	PG4	O2-C2-C1	2.06	119.21	110.11

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	PG4	O3-C5-C6-O4
2	A	401	PG4	O3-C5-C6-O4
2	A	401	PG4	O4-C7-C8-O5
2	B	401	PG4	O2-C3-C4-O3
2	B	401	PG4	O4-C7-C8-O5
2	A	401	PG4	O2-C3-C4-O3
2	A	401	PG4	C4-C3-O2-C2
2	A	401	PG4	C6-C5-O3-C4
2	B	401	PG4	C6-C5-O3-C4
2	B	401	PG4	C4-C3-O2-C2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	PG4	2	0
2	B	401	PG4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/333 (94%)	-0.27	4 (1%) 75 75	11, 23, 46, 77	5 (1%)
1	B	316/333 (94%)	-0.20	3 (0%) 81 81	11, 24, 48, 79	4 (1%)
1	C	313/333 (93%)	-0.06	9 (2%) 53 53	10, 26, 53, 90	7 (2%)
All	All	945/999 (94%)	-0.18	16 (1%) 69 69	10, 24, 49, 90	16 (1%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	7	LEU	5.1
1	C	8	PRO	4.0
1	A	31	TRP	2.9
1	C	61[A]	ARG	2.7
1	B	30	PRO	2.7
1	A	30	PRO	2.6
1	B	31	TRP	2.5
1	C	213	TRP	2.5
1	C	31	TRP	2.4
1	C	10	LEU	2.4
1	A	24	LEU	2.4
1	B	4	GLU	2.3
1	C	30	PRO	2.2
1	C	212	VAL	2.1
1	C	70	GLY	2.1
1	A	29	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PG4	A	401	13/13	0.83	0.16	34,47,63,64	0
2	PG4	B	401	13/13	0.83	0.17	39,49,68,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.