



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:03 PM UTC

PDB ID : 3KSL / pdb_00003ksl
Title : Structure of FPT bound to DATFP-DH-GPP
Authors : Hovlid, M.L.; Edelstein, R.L.; Henry, O.; Ochocki, J.; DeGraw, A.; Lenevich, S.; Talbot, T.; Young, V.; Hruza, A.W.; Lopez-Gallego, F.; Labello, N.P.; Strickland, C.L.; Schmidt-Dannert, C.; Distefano, M.D.
Deposited on : 2009-11-23
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	
2	B	437	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SZH	B	438[A]	X	-	-	-
5	SZH	B	438[B]	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Farnesyltransferase, CAAX box, alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2680	1706	470	499	5			

- Molecule 2 is a protein called Protein farnesyltransferase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	403	Total	C	N	O	S	0	0	0
			3169	2026	545	575	23			


- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is (2S,6E)-8-{[(R)-hydroxy(phosphonooxy)phosphoryl]oxy}-2,6-dimethyl oct-6-en-1-yl (2S)-3,3,3-trifluoro-2-hydrazinopropanoate (CCD ID: SZH) (formula: C₁₃H₂₅F₃N₂O₉P₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	B	1	Total	C	F	N	O	P	0	1
			58	26	6	4	18	4		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	212	Total O 212 212	0	0
6	B	215	Total O 215 215	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	171.19Å 171.19Å 69.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.45 – 2.05 29.45 – 2.05	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.45-2.05) 97.8 (29.45-2.05)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.04Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, R_{free}	0.180 , 0.214 0.178 , 0.212	Depositor DCC
R_{free} test set	3569 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6336	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SZH, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/2746 (0.0%)	1.23	7/3728 (0.2%)
2	B	0.87	0/3255	1.21	2/4421 (0.0%)
All	All	0.85	1/6001 (0.0%)	1.22	9/8149 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	152	MET	SD-CE	-5.19	1.66	1.79

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	217	ASP	N-CA-C	7.49	120.39	111.33
1	A	185	GLU	N-CA-C	7.43	119.07	110.97
2	B	352	ASP	CA-CB-CG	7.15	119.75	112.60
1	A	183	SER	N-CA-C	5.67	118.23	111.71
1	A	348	LYS	N-CA-C	5.63	119.99	113.18
1	A	219	GLU	N-CA-C	5.25	117.09	111.36
1	A	311	LEU	CA-C-N	5.02	126.88	120.56
1	A	311	LEU	C-N-CA	5.02	126.88	120.56
2	B	301	SER	N-CA-C	-5.00	105.96	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2680	0	2600	8	0
2	B	3169	0	3098	16	0
3	B	1	0	0	0	0
4	B	1	0	0	0	0
5	B	58	0	40	2	0
6	A	212	0	0	0	0
6	B	215	0	0	0	0
All	All	6336	0	5738	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:21:GLU:HB2	2:B:22:PRO:CD	2.22	0.69
2:B:21:GLU:HB2	2:B:22:PRO:HD2	1.78	0.64
1:A:67:ARG:HH22	1:A:94:GLU:CD	2.15	0.55
2:B:245:MET:HA	2:B:245:MET:HE3	1.89	0.53
1:A:342:GLU:HG3	1:A:346:LYS:HD2	1.91	0.53
2:B:239:ILE:HB	2:B:252:THR:HA	1.92	0.51
2:B:192:LEU:HD23	2:B:199:VAL:HG23	1.92	0.51
2:B:250:GLY:HA3	5:B:438[A]:SZH:H10A	1.93	0.51
1:A:77:PRO:HB2	1:A:101:ASP:HB3	1.93	0.50
2:B:250:GLY:HA3	5:B:438[B]:SZH:H12	1.94	0.50
2:B:23:LEU:HD12	2:B:26:LEU:HD22	1.95	0.48
2:B:301:SER:O	2:B:305:ALA:HB3	2.15	0.47
1:A:300:LEU:HD12	1:A:303:GLN:HG3	1.97	0.47
1:A:303:GLN:O	1:A:307:SER:HB3	2.14	0.47
2:B:193:MET:SD	2:B:203:SER:HB3	2.56	0.45
1:A:112:ARG:HA	1:A:140:LEU:CD2	2.47	0.44
2:B:253:PHE:HA	2:B:307:LEU:HD21	2.00	0.43
2:B:268:LEU:HD13	2:B:273:LEU:HD12	2.01	0.43
2:B:306:GLY:HA2	2:B:372:ILE:HD13	2.02	0.42
2:B:297:ASP:HB3	2:B:300:TYR:CD1	2.55	0.42
1:A:204:GLN:HA	2:B:245:MET:SD	2.61	0.41
2:B:308:LEU:HD13	2:B:308:LEU:HA	1.97	0.40
1:A:287:ARG:HB3	1:A:291:ARG:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/377 (83%)	300 (96%)	12 (4%)	1 (0%)	36	30
2	B	401/437 (92%)	396 (99%)	5 (1%)	0	100	100
All	All	714/814 (88%)	696 (98%)	17 (2%)	1 (0%)	48	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/338 (87%)	284 (97%)	9 (3%)	35	30
2	B	340/371 (92%)	333 (98%)	7 (2%)	47	46
All	All	633/709 (89%)	617 (98%)	16 (2%)	42	39

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	ASP
1	A	85	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	GLU
1	A	286	ASP
1	A	300	LEU
1	A	323	LEU
1	A	331	GLU
1	A	363	LEU
1	A	368	SER
2	B	21	GLU
2	B	108	LEU
2	B	245	MET
2	B	266	ARG
2	B	351	LEU
2	B	382	LEU
2	B	418	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	GLN
1	A	149	GLN
1	A	184	GLN
2	B	80	HIS
2	B	128	GLN
2	B	327	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SZH	B	438[A]	-	26,28,28	1.80	1 (3%)	33,40,40	1.44	5 (15%)
5	SZH	B	438[B]	-	26,28,28	1.90	2 (7%)	33,40,40	1.12	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SZH	B	438[A]	-	1/1/7/9	10/34/36/36	-
5	SZH	B	438[B]	-	2/2/7/9	12/34/36/36	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	438[A]	SZH	C3-N15	-7.97	1.31	1.47
5	B	438[B]	SZH	C3-N15	-7.64	1.32	1.47
5	B	438[B]	SZH	P29-O24	4.29	1.64	1.59

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	438[B]	SZH	F25-C13-C3	3.65	118.45	111.97
5	B	438[A]	SZH	F25-C13-C3	3.62	118.41	111.97
5	B	438[A]	SZH	C7-C1-C2	3.33	131.66	126.20
5	B	438[A]	SZH	F26-C13-F25	-2.64	101.13	106.87
5	B	438[B]	SZH	F26-C13-F25	-2.15	102.19	106.87
5	B	438[A]	SZH	C9-C8-C2	2.09	118.56	113.47
5	B	438[A]	SZH	O22-C4-O19	2.03	127.86	124.14

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	438[A]	SZH	C3
5	B	438[B]	SZH	C12
5	B	438[B]	SZH	C3

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	438[A]	SZH	F25-C13-C3-C4
5	B	438[A]	SZH	F26-C13-C3-C4
5	B	438[A]	SZH	F27-C13-C3-C4
5	B	438[B]	SZH	F25-C13-C3-C4
5	B	438[B]	SZH	F26-C13-C3-C4
5	B	438[B]	SZH	F27-C13-C3-C4
5	B	438[B]	SZH	C9-C10-C12-C11
5	B	438[B]	SZH	C7-O23-P29-O21
5	B	438[B]	SZH	C7-O23-P29-O24
5	B	438[B]	SZH	C2-C8-C9-C10
5	B	438[A]	SZH	C3-C4-O22-C11
5	B	438[A]	SZH	C2-C8-C9-C10
5	B	438[A]	SZH	O19-C4-O22-C11
5	B	438[A]	SZH	P28-O24-P29-O18
5	B	438[B]	SZH	O22-C11-C12-C10
5	B	438[B]	SZH	C7-O23-P29-O18
5	B	438[B]	SZH	C9-C10-C12-C6
5	B	438[B]	SZH	O22-C11-C12-C6
5	B	438[A]	SZH	C13-C3-C4-O19
5	B	438[A]	SZH	C13-C3-C4-O22
5	B	438[B]	SZH	C5-C2-C8-C9
5	B	438[A]	SZH	P28-O24-P29-O21


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	438[A]	SZH	1	0
5	B	438[B]	SZH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	315/377 (83%)	0.05	14 (4%) 39 39	20, 34, 64, 90	0
2	B	403/437 (92%)	-0.13	8 (1%) 65 67	20, 32, 54, 94	0
All	All	718/814 (88%)	-0.05	22 (3%) 51 52	20, 33, 59, 94	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	423	PHE	8.1
1	A	368	SER	5.3
1	A	55	PHE	5.3
2	B	422	GLY	4.6
2	B	64	PHE	4.6
2	B	21	GLU	3.3
1	A	306	HIS	3.0
1	A	302	LEU	3.0
1	A	331	GLU	3.0
1	A	54	GLY	2.9
2	B	22	PRO	2.8
1	A	304	PRO	2.8
1	A	303	GLN	2.7
1	A	330	LYS	2.6
2	B	70	ARG	2.6
1	A	305	SER	2.4
1	A	300	LEU	2.3
1	A	166	TYR	2.3
1	A	307	SER	2.2
2	B	62	TYR	2.1
2	B	382	LEU	2.1
1	A	56	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SZH	B	438[A]	29/29	0.95	0.14	20,47,70,71	29
5	SZH	B	438[B]	29/29	0.95	0.14	75,77,79,79	29
3	MG	B	1001	1/1	0.97	0.13	32,32,32,32	0
4	ZN	B	2001	1/1	1.00	0.03	36,36,36,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SZH B 438 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SZH B 438 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.