



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:29 AM UTC

PDB ID : 4KTK / pdb_00004ktk
Title : Crystal structure of Mycobacterium tuberculosis CYP121 in complex with 4-(3-amino-4-(4-hydroxyphenyl)-1H-pyrazol-5-yl)benzene-1,3-diol
Authors : Hudson, S.A.
Deposited on : 2013-05-20
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	2% 84% 14%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 121.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	21	1
			3053	1950	532	561	10			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 4-[3-amino-4-(4-hydroxyphenyl)-1H-pyrazol-5-yl]benzene-1,3-diol (CCD ID: KTK) (formula: C₁₅H₁₃N₃O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	O	0
			21	15	3	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			21	15	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	468	Total	O	0	8
			476	476		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome P450 121



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.77Å 77.77Å 263.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.47 – 1.40 38.47 – 1.40	Depositor EDS
% Data completeness (in resolution range)	81.2 (38.47-1.40) 81.2 (38.47-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.155 , 0.183 0.155 , 0.184	Depositor DCC
R_{free} test set	3792 reflections (4.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.4	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3624	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, KTK, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.81	38/3211 (1.2%)	1.55	33/4373 (0.8%)

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	290	ALA	CA-C	-7.99	1.43	1.52
1	A	57	ASP	C-O	7.15	1.32	1.23
1	A	388	ILE	CA-CB	-6.66	1.45	1.54
1	A	292	ILE	C-O	6.64	1.30	1.24
1	A	184	ARG	C-O	6.62	1.31	1.24
1	A	186	ILE	C-O	6.54	1.31	1.24
1	A	309	LEU	C-O	6.46	1.31	1.24
1	A	32	ARG	CZ-NH2	-6.31	1.25	1.33
1	A	144	ALA	C-O	6.23	1.31	1.24
1	A	32	ARG	CZ-NH1	6.22	1.41	1.32
1	A	138	ALA	C-O	6.10	1.31	1.24
1	A	292	ILE	CA-C	-5.86	1.45	1.52
1	A	67	ALA	CA-CB	-5.82	1.43	1.53
1	A	25	LEU	CA-CB	5.81	1.62	1.53
1	A	252	ARG	N-CA	5.80	1.53	1.46
1	A	220	ASP	CA-C	5.80	1.60	1.52
1	A	25	LEU	N-CA	-5.76	1.39	1.46
1	A	55	LEU	CA-C	5.64	1.60	1.52
1	A	215	TYR	N-CA	5.57	1.52	1.46
1	A	63	LYS	CA-CB	5.56	1.62	1.53
1	A	277	ASN	C-O	-5.51	1.17	1.23
1	A	288	ALA	CA-CB	-5.51	1.44	1.53
1	A	54	VAL	CA-CB	5.42	1.60	1.54
1	A	253	PRO	C-O	-5.38	1.17	1.24
1	A	251[A]	GLN	CA-CB	-5.37	1.44	1.53
1	A	251[B]	GLN	CA-CB	-5.37	1.44	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	329	ARG	CA-C	5.29	1.59	1.53
1	A	86	MET	N-CA	5.24	1.52	1.46
1	A	208	ARG	CD-NE	-5.22	1.39	1.46
1	A	117	ALA	CA-CB	-5.12	1.45	1.53
1	A	20	ASP	CA-C	-5.10	1.45	1.52
1	A	14	ARG	N-CA	5.09	1.52	1.46
1	A	199	THR	CA-CB	5.08	1.62	1.54
1	A	293	GLN	N-CA	-5.07	1.40	1.46
1	A	140	PRO	C-N	-5.06	1.27	1.33
1	A	373	VAL	N-CA	5.05	1.51	1.46
1	A	151	GLY	CA-C	-5.04	1.44	1.51
1	A	187	GLU	C-O	5.01	1.29	1.24

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	252	ARG	O-C-N	-8.14	114.09	121.34
1	A	32	ARG	NE-CZ-NH2	-7.86	112.13	119.20
1	A	32	ARG	NE-CZ-NH1	7.59	129.09	121.50
1	A	221	GLU	CB-CG-CD	7.30	125.01	112.60
1	A	329	ARG	O-C-N	6.56	128.74	121.53
1	A	195	ASN	CA-C-O	6.22	123.33	119.29
1	A	164	LEU	N-CA-C	6.08	118.87	111.82
1	A	383[A]	ARG	NE-CZ-NH2	5.93	124.54	119.20
1	A	383[B]	ARG	NE-CZ-NH2	5.93	124.54	119.20
1	A	304	LEU	CA-C-O	5.88	127.19	120.54
1	A	191	GLY	O-C-N	-5.64	116.72	122.19
1	A	53	GLN	CA-C-O	5.63	126.92	120.90
1	A	388	ILE	N-CA-CB	5.57	119.01	111.21
1	A	50	LEU	CA-C-O	5.56	126.32	120.42
1	A	225	THR	O-C-N	-5.41	115.99	122.15
1	A	321	ASN	CA-C-O	5.37	124.56	120.26
1	A	184	ARG	NE-CZ-NH2	-5.35	114.39	119.20
1	A	8	GLU	CB-CG-CD	-5.28	103.63	112.60
1	A	290	ALA	O-C-N	-5.27	117.73	123.26
1	A	270	VAL	O-C-N	-5.26	116.77	121.87
1	A	321	ASN	O-C-N	5.25	123.94	121.53
1	A	250	ILE	CA-C-O	-5.24	115.61	121.05
1	A	277	ASN	CA-C-O	5.22	127.00	121.05
1	A	208	ARG	NE-CZ-NH1	-5.21	116.29	121.50
1	A	55	LEU	N-CA-C	5.18	117.83	111.82
1	A	353	ARG	CG-CD-NE	-5.13	100.71	112.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	366	MET	O-C-N	-5.07	116.94	121.20
1	A	72	ARG	CA-C-N	5.05	127.31	120.44
1	A	72	ARG	C-N-CA	5.05	127.31	120.44
1	A	360	GLU	CA-C-N	5.04	127.00	120.44
1	A	360	GLU	C-N-CA	5.04	127.00	120.44
1	A	267	PRO	O-C-N	-5.03	116.19	122.23
1	A	191	GLY	N-CA-C	-5.02	106.34	112.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3053	0	3071	14	1
2	A	43	0	30	2	0
3	A	10	0	0	1	0
4	A	42	0	23	0	0
5	A	476	0	0	5	1
All	All	3624	0	3124	17	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:GLU:CD	5:A:807:HOH:O	1.92	1.10
3:A:402:SO4:O4	5:A:524:HOH:O	1.95	0.84
1:A:8:GLU:CG	5:A:807:HOH:O	2.23	0.77
1:A:181:ASN:OD1	1:A:184:ARG:NH2	2.17	0.74
1:A:353:ARG:HH11	1:A:356:GLN:HE21	1.37	0.72
1:A:74:ASN:C	1:A:74:ASN:HD22	2.11	0.57
1:A:104:PRO:HA	1:A:109[B]:LEU:HD23	1.91	0.52
1:A:331:ASN:HD22	1:A:331:ASN:H	1.57	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:ARG:HH11	1:A:356:GLN:NE2	2.05	0.51
1:A:8:GLU:HG2	5:A:807:HOH:O	2.01	0.48
1:A:352:ARG:NH1	5:A:878:HOH:O	2.49	0.44
1:A:144:ALA:HA	1:A:154:GLN:HE22	1.82	0.43
1:A:56:GLU:OE2	1:A:340[B]:ARG:HD2	2.21	0.41
1:A:331:ASN:HD22	1:A:331:ASN:N	2.18	0.40
2:A:401:HEM:HBB2	2:A:401:HEM:CMB	2.51	0.40
2:A:401:HEM:HMC1	2:A:401:HEM:HBC2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:ASN:OD1	5:A:936:HOH:O[8_565]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	414/395 (105%)	410 (99%)	3 (1%)	1 (0%)	43 19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/325 (101%)	324 (99%)	4 (1%)	63	34

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ARG
1	A	74	ASN
1	A	277	ASN
1	A	331	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	154	GLN
1	A	331	ASN
1	A	342	GLN
1	A	356	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	402	-	4,4,4	0.60	0	6,6,6	0.27	0
4	KTK	A	405	-	23,23,23	2.34	10 (43%)	27,33,33	2.63	11 (40%)
3	SO4	A	403	-	4,4,4	1.04	0	6,6,6	1.46	2 (33%)
4	KTK	A	404	-	23,23,23	2.38	11 (47%)	27,33,33	1.94	10 (37%)
2	HEM	A	401	5,1	50,50,50	1.63	12 (24%)	67,82,82	1.72	12 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	KTK	A	405	-	-	1/8/8/8	0/3/3/3
4	KTK	A	404	-	-	0/8/8/8	0/3/3/3
2	HEM	A	401	5,1	-	1/14/54/54	-

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	404	KTK	CAN-CAJ	-5.72	1.38	1.49
4	A	405	KTK	CAN-CAJ	-5.12	1.40	1.49
4	A	404	KTK	CAC-CAB	-4.85	1.40	1.48
2	A	401	HEM	FE-NB	4.80	2.09	1.94
4	A	405	KTK	CAC-CAB	-4.53	1.41	1.48
4	A	404	KTK	CAH-CAC	4.08	1.46	1.39
4	A	405	KTK	CAC-CAD	3.37	1.46	1.40
4	A	405	KTK	CAS-CAN	3.32	1.44	1.39
4	A	405	KTK	CAE-CAF	3.19	1.43	1.39
2	A	401	HEM	FE-NA	3.05	2.05	1.95
2	A	401	HEM	FE-NC	3.00	2.05	1.95
4	A	404	KTK	OAU-CAD	3.00	1.42	1.36
4	A	404	KTK	CAB-NAA	2.98	1.40	1.35
2	A	401	HEM	O2D-CGD	-2.93	1.21	1.30
2	A	401	HEM	C1B-NB	-2.86	1.35	1.40
4	A	404	KTK	CAO-CAN	-2.85	1.35	1.39
4	A	405	KTK	CAO-CAN	2.82	1.43	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	405	KTK	CAE-CAD	-2.57	1.35	1.38
2	A	401	HEM	CBD-CGD	2.56	1.56	1.50
2	A	401	HEM	CMA-C3A	-2.54	1.45	1.50
4	A	404	KTK	CAE-CAD	-2.53	1.35	1.38
2	A	401	HEM	CHB-C1B	2.40	1.43	1.38
4	A	404	KTK	CAK-NAL	2.30	1.35	1.33
4	A	404	KTK	CAP-CAQ	-2.27	1.34	1.39
4	A	404	KTK	CAJ-CAB	-2.24	1.35	1.39
2	A	401	HEM	CMB-C2B	-2.22	1.46	1.50
2	A	401	HEM	FE-ND	2.21	2.01	1.94
4	A	405	KTK	CAS-CAR	2.21	1.42	1.38
4	A	404	KTK	CAS-CAR	2.19	1.42	1.38
2	A	401	HEM	C2A-C3A	-2.13	1.33	1.38
4	A	405	KTK	OAI-CAF	-2.09	1.32	1.37
2	A	401	HEM	C4A-C3A	2.05	1.47	1.43
4	A	405	KTK	CAG-CAH	2.03	1.42	1.38

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	C4C-C3C-C2C	5.86	111.90	106.81
4	A	405	KTK	CAP-CAO-CAN	5.06	126.21	120.80
4	A	405	KTK	CAJ-CAK-NAM	4.83	132.55	126.13
2	A	401	HEM	C1B-NB-C4B	4.78	110.86	105.21
2	A	401	HEM	C3B-C4B-NB	-4.51	106.23	109.47
4	A	405	KTK	NAM-CAK-NAL	-4.39	116.20	122.05
4	A	405	KTK	CAD-CAE-CAF	4.38	123.71	119.67
4	A	405	KTK	CAS-CAR-CAQ	4.10	124.21	119.88
4	A	405	KTK	CAG-CAF-CAE	-3.92	115.88	120.19
4	A	405	KTK	CAO-CAP-CAQ	-3.83	115.83	119.88
2	A	401	HEM	C3C-C2C-C1C	-3.79	103.46	107.05
4	A	405	KTK	CAC-CAB-CAJ	-3.73	127.52	131.65
4	A	404	KTK	CAC-CAB-NAA	-3.49	116.75	121.25
2	A	401	HEM	CHC-C1C-NC	-3.45	120.69	124.45
4	A	405	KTK	CAC-CAB-NAA	3.43	125.67	121.25
2	A	401	HEM	CMA-C3A-C2A	3.37	132.76	125.62
4	A	404	KTK	CAS-CAR-CAQ	-3.30	116.39	119.88
4	A	405	KTK	CAR-CAS-CAN	-3.19	117.39	120.80
4	A	404	KTK	NAM-CAK-NAL	-3.17	117.82	122.05
4	A	404	KTK	CAH-CAC-CAD	-3.14	114.20	118.15
4	A	404	KTK	CAC-CAB-CAJ	3.00	134.98	131.65
4	A	404	KTK	CAN-CAJ-CAB	2.64	132.08	127.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	404	KTK	CAR-CAS-CAN	2.63	123.61	120.80
2	A	401	HEM	CMA-C3A-C4A	-2.54	121.55	125.42
3	A	403	SO4	O4-S-O3	2.52	122.46	108.54
4	A	404	KTK	CAS-CAN-CAO	-2.47	115.43	118.57
2	A	401	HEM	O1D-CGD-CBD	-2.34	115.67	123.09
4	A	404	KTK	CAJ-CAK-NAM	2.27	129.14	126.13
2	A	401	HEM	C1A-C2A-C3A	2.26	110.36	106.87
4	A	404	KTK	OAU-CAD-CAE	-2.23	113.44	119.45
2	A	401	HEM	CMC-C2C-C3C	2.19	133.73	128.43
2	A	401	HEM	C3B-C2B-C1B	2.19	108.06	106.41
4	A	405	KTK	CAS-CAN-CAO	-2.13	115.86	118.57
2	A	401	HEM	O2A-CGA-O1A	-2.02	118.14	123.33
3	A	403	SO4	O4-S-O1	-2.01	99.04	109.56

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	405	KTK	NAA-CAB-CAC-CAD
2	A	401	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	SO4	1	0
2	A	401	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/395 (100%)	-0.06	7 (1%) 67 71	7, 14, 29, 48	23 (5%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	THR	10.4
1	A	3	ALA	3.9
1	A	4	THR	3.1
1	A	217	HIS	2.7
1	A	108	GLY	2.5
1	A	106	ALA	2.2
1	A	107	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	402	5/5	0.92	0.12	54,67,71,79	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	403	5/5	0.95	0.09	21,21,31,36	0
4	KTK	A	405	21/21	0.95	0.07	14,17,20,22	0
4	KTK	A	404	21/21	0.96	0.07	13,17,20,21	0
2	HEM	A	401	43/43	0.98	0.06	8,11,17,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KTK A 405:


2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KTK A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.