



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:58 AM UTC

PDB ID : 5KTE / pdb_00005kte
Title : Crystal structure of Deinococcus radiodurans MntH, an Nramp-family transition metal transporter
Authors : Bane, L.B.; Gaudet, R.; Weihofen, W.A.; Singharoy, A.
Deposited on : 2016-07-11
Resolution : 3.94 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1022 (4.12-3.76)
Clashscore	190562	1000 (4.10-3.78)
Ramachandran outliers	187476	1009 (4.12-3.76)
Sidechain outliers	187428	1002 (4.12-3.76)
RSRZ outliers	180081	1022 (4.12-3.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	
2	H	213	
3	L	213	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Divalent metal cation transporter MntH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	338	2407	1586	385	420	16	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	MET	-	initiating methionine	UNP Q9RTP8
A	18	HIS	-	expression tag	UNP Q9RTP8
A	19	HIS	-	expression tag	UNP Q9RTP8
A	20	HIS	-	expression tag	UNP Q9RTP8
A	21	HIS	-	expression tag	UNP Q9RTP8
A	22	HIS	-	expression tag	UNP Q9RTP8
A	23	HIS	-	expression tag	UNP Q9RTP8
A	24	HIS	-	expression tag	UNP Q9RTP8
A	25	HIS	-	expression tag	UNP Q9RTP8
A	168	HIS	GLN	engineered mutation	UNP Q9RTP8
A	169	HIS	LYS	engineered mutation	UNP Q9RTP8
A	251	TYR	GLU	engineered mutation	UNP Q9RTP8
A	252	TYR	GLU	engineered mutation	UNP Q9RTP8
A	253	TYR	LYS	engineered mutation	UNP Q9RTP8
A	398	HIS	ARG	engineered mutation	UNP Q9RTP8
A	399	HIS	ARG	engineered mutation	UNP Q9RTP8

- Molecule 2 is a protein called Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	211	1567	992	251	316	8	0	0	0

- Molecule 3 is a protein called Fab Light Chain.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1648	1019	282	341	6	0	0	0

- Molecule 4 is OSMIUM ION (CCD ID: OS) (formula: Os).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Os	0	0
			2	2		
4	H	1	Total	Os	0	0
			1	1		



● Molecule 3: Fab Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	113.13Å 132.08Å 221.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.47 – 3.94 46.47 – 3.94	Depositor EDS
% Data completeness (in resolution range)	76.0 (46.47-3.94) 75.1 (46.47-3.94)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.43 (at 4.00Å)	Xtrriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.268 , 0.313 0.265 , 0.306	Depositor DCC
R_{free} test set	1179 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	-4.7	Xtrriage
Anisotropy	5.659	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.74	EDS
Total number of atoms	5625	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.15	0/2455	0.39	0/3363
2	H	0.19	0/1609	0.46	0/2205
3	L	0.18	0/1685	0.44	0/2286
All	All	0.17	0/5749	0.42	0/7854

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	195	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2407	0	2426	66	0
2	H	1567	0	1532	53	0
3	L	1648	0	1565	54	0
4	A	2	0	0	0	0
4	H	1	0	0	0	0
All	All	5625	0	5523	166	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 166 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:189:HIS:O	3:L:211:ARG:NH1	2.07	0.86
2:H:49:VAL:HG12	2:H:59:TYR:HA	1.68	0.75
3:L:147:LYS:NZ	3:L:156:GLN:HG2	2.00	0.75
2:H:53:SER:HA	2:H:71:LYS:HE3	1.67	0.74
3:L:149:LYS:HB2	3:L:193:THR:HB	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/420 (78%)	301 (92%)	26 (8%)	1 (0%)	36	70
2	H	207/213 (97%)	184 (89%)	21 (10%)	2 (1%)	12	45
3	L	211/213 (99%)	191 (90%)	18 (8%)	2 (1%)	14	48
All	All	746/846 (88%)	676 (91%)	65 (9%)	5 (1%)	18	53

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	291	VAL
3	L	51	VAL
2	H	160	LEU
3	L	77	SER
2	H	42	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/326 (72%)	231 (99%)	2 (1%)	70	76
2	H	185/187 (99%)	185 (100%)	0	100	100
3	L	192/193 (100%)	190 (99%)	2 (1%)	68	76
All	All	610/706 (86%)	606 (99%)	4 (1%)	76	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	259	LEU
1	A	291	VAL
3	L	76	ASN
3	L	77	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	L	31	ASN
3	L	53	GLN
3	L	90	GLN
1	A	399	HIS
1	A	59	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/420 (80%)	2.37	162 (47%) 0 1	61, 116, 169, 230	0
2	H	211/213 (99%)	0.79	28 (13%) 7 10	17, 56, 98, 157	0
3	L	213/213 (100%)	1.41	49 (23%) 2 4	44, 73, 116, 175	0
All	All	762/846 (90%)	1.66	239 (31%) 1 3	17, 86, 154, 230	0

The worst 5 of 239 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	108	ARG	16.1
1	A	109	GLU	15.0
1	A	199	PRO	11.7
3	L	17	ASP	11.2
1	A	232	HIS	9.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	OS	H	301	1/1	0.89	0.16	173,173,173,173	0
4	OS	A	502	1/1	0.90	0.14	192,192,192,192	0
4	OS	A	501	1/1	0.94	0.16	205,205,205,205	0

6.5 Other polymers [i](#)

There are no such residues in this entry.