



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 04:15 PM UTC

PDB ID : 6KU0 / pdb_00006ku0
Title : Crystal structure of MyoVa-GTD in complex with MICAL1-GTBM
Authors : Niu, F.; Wei, Z.
Deposited on : 2019-08-29
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	 3% 88% 6% 5%
1	C	389	 4% 85% 7% 8%
2	B	27	 7% 74% 7% 15%
2	D	27	 4% 78% 15%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-Va.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	13	0
			2989	1903	512	549	25			
1	C	359	Total	C	N	O	S	0	14	0
			2944	1874	504	542	24			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1465	GLY	-	expression tag	UNP Q99104
A	1466	PRO	-	expression tag	UNP Q99104
A	1467	GLY	-	expression tag	UNP Q99104
A	1468	SER	-	expression tag	UNP Q99104
C	1465	GLY	-	expression tag	UNP Q99104
C	1466	PRO	-	expression tag	UNP Q99104
C	1467	GLY	-	expression tag	UNP Q99104
C	1468	SER	-	expression tag	UNP Q99104

- Molecule 2 is a protein called Peptide from [F-actin]-monooxygenase MICAL1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	23	Total	C	N	O	0	1	0
			195	116	42	37			
2	D	23	Total	C	N	O	0	1	0
			195	116	42	37			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	796	GLY	-	expression tag	UNP Q8TDZ2
B	797	PRO	-	expression tag	UNP Q8TDZ2
B	798	GLY	-	expression tag	UNP Q8TDZ2
D	796	GLY	-	expression tag	UNP Q8TDZ2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	797	PRO	-	expression tag	UNP Q8TDZ2
D	798	GLY	-	expression tag	UNP Q8TDZ2

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	325	Total O 325 325	0	0
4	B	31	Total O 31 31	0	0
4	C	320	Total O 320 320	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	24	Total	O	0	0
			24	24		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Unconventional myosin-Va




- Molecule 1: Unconventional myosin-Va



- Molecule 2: Peptide from [F-actin]-monooxygenase MICAL1



- Molecule 2: Peptide from [F-actin]-monooxygenase MICAL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.29Å 63.22Å 74.42Å 65.78° 86.71° 85.81°	Depositor
Resolution (Å)	50.00 – 1.60 50.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	91.8 (50.00-1.60) 88.4 (50.00-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.161 , 0.199 0.164 , 0.199	Depositor DCC
R_{free} test set	2014 reflections (1.66%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.003 for -h,-k,-k+l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7043	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	2/3076 (0.1%)	1.02	2/4149 (0.0%)
1	C	1.03	3/3031 (0.1%)	1.01	2/4088 (0.0%)
2	B	1.34	1/199 (0.5%)	1.29	3/265 (1.1%)
2	D	1.08	1/199 (0.5%)	1.13	0/265
All	All	1.07	7/6505 (0.1%)	1.03	7/8767 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	1624	VAL	C-O	-6.01	1.17	1.24
1	C	1512	PHE	C-O	-5.40	1.17	1.24
1	A	1623	ILE	C-O	-5.37	1.18	1.24
2	D	803	ARG	CA-C	-5.36	1.46	1.52
1	A	1512	PHE	C-O	-5.26	1.17	1.24
1	C	1622	MET	C-O	-5.24	1.17	1.24
2	B	802	THR	C-O	-5.00	1.18	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1623	ILE	N-CA-C	9.01	119.04	110.30
1	A	1623	ILE	N-CA-C	7.69	117.76	110.30
1	A	1810	LYS	N-CA-C	-6.58	103.76	112.72
1	C	1547	ASP	N-CA-C	-5.87	106.16	113.38
2	B	802	THR	N-CA-C	-5.80	100.53	109.76
2	B	800	GLN	CA-C-N	-5.74	114.70	120.21
2	B	800	GLN	C-N-CA	-5.74	114.70	120.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2989	0	3074	17	0
1	C	2944	0	3020	16	0
2	B	195	0	208	3	0
2	D	195	0	208	1	0
3	A	12	0	18	1	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
4	A	325	0	0	3	0
4	B	31	0	0	0	0
4	C	320	0	0	2	0
4	D	24	0	0	0	0
All	All	7043	0	6540	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:800:GLN:HG3	2:B:803:ARG:HH12	1.53	0.71
1:A:1726:GLU:OE1	1:A:1729:ARG:NH2	2.27	0.65
1:A:1573:GLU:HA	1:A:1576:MET:HE2	1.80	0.63
1:C:1701:ASN:OD1	1:C:1705:ARG:HD2	1.99	0.63
1:C:1485:LEU:HD13	1:C:1850[B]:ILE:HD11	1.82	0.62
1:C:1660:ASP:HB3	1:C:1664:ARG:NH1	2.19	0.58
1:C:1701:ASN:O	1:C:1705:ARG:HG3	2.08	0.54
1:A:1499:ALA:HB1	1:A:1842:PRO:HG3	1.89	0.53
1:A:1729:ARG:NH1	4:A:2003:HOH:O	2.44	0.50
1:C:1631:GLU:HA	1:C:1716:GLN:HE22	1.77	0.49
1:A:1846:GLY:HA2	3:A:1901:EDO:H11	1.94	0.49
1:C:1847:LEU:O	1:C:1850[A]:ILE:HG12	2.14	0.48
1:C:1479:ARG:HD3	1:C:1526:LYS:HG3	1.97	0.47
1:A:1729:ARG:HD3	4:A:2267:HOH:O	2.15	0.47
1:A:1542:LEU:O	1:A:1546:GLY:HA3	2.16	0.46
1:C:1546:GLY:HA3	1:C:1610:GLN:NE2	2.31	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1718:ARG:HD3	1:C:1753:GLN:OE1	2.15	0.46
1:A:1546:GLY:HA3	1:A:1610:GLN:NE2	2.30	0.46
2:B:800:GLN:OE1	2:B:803:ARG:NH2	2.42	0.46
1:A:1620:GLN:HG3	4:A:2168:HOH:O	2.15	0.45
1:C:1702:LEU:HD23	1:C:1702:LEU:C	2.41	0.45
1:A:1499:ALA:CB	1:A:1842:PRO:HG3	2.48	0.44
1:A:1571:GLY:O	2:B:800:GLN:HG2	2.18	0.44
1:C:1739:LYS:HE3	4:C:2031:HOH:O	2.18	0.43
1:C:1526:LYS:NZ	4:C:2016:HOH:O	2.51	0.43
2:D:800:GLN:HG3	2:D:803:ARG:HH12	1.83	0.42
1:C:1574:GLY:HA2	1:C:1577:LYS:HE3	2.00	0.42
1:A:1572:GLU:O	1:A:1576:MET:HG3	2.20	0.42
1:C:1705:ARG:HD2	1:C:1705:ARG:HH11	1.72	0.42
1:A:1805[B]:ARG:HA	1:A:1805[B]:ARG:HD3	1.86	0.41
1:A:1704:LEU:HD23	1:A:1704:LEU:HA	1.91	0.41
1:C:1496:ARG:HE	1:C:1496:ARG:HB2	1.65	0.41
1:A:1801:THR:O	1:A:1805[B]:ARG:HG2	2.21	0.41
1:A:1613:ARG:HD3	1:A:1613:ARG:HA	1.81	0.40
1:C:1725[A]:GLU:HG2	1:C:1739:LYS:HG3	2.03	0.40
1:A:1512:PHE:CZ	1:A:1569:TYR:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/389 (97%)	373 (99%)	4 (1%)	0	100	100
1	C	369/389 (95%)	364 (99%)	5 (1%)	0	100	100
2	B	22/27 (82%)	22 (100%)	0	0	100	100
2	D	22/27 (82%)	22 (100%)	0	0	100	100
All	All	790/832 (95%)	781 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/352 (97%)	338 (98%)	5 (2%)	57	35
1	C	338/352 (96%)	336 (99%)	2 (1%)	78	66
2	B	24/25 (96%)	23 (96%)	1 (4%)	26	7
2	D	24/25 (96%)	24 (100%)	0	100	100
All	All	729/754 (97%)	721 (99%)	8 (1%)	68	46

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1493	LEU
1	A	1495	PRO
1	A	1498	VAL
1	A	1712[A]	SER
1	A	1712[B]	SER
2	B	800	GLN
1	C	1739	LYS
1	C	1820	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1634	GLN
1	A	1670	HIS
1	A	1732	ASN
1	A	1821	HIS
1	A	1830	ASN
1	C	1830	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	1901	-	3,3,3	0.46	0	2,2,2	0.41	0
3	EDO	D	901	-	3,3,3	0.49	0	2,2,2	0.56	0
3	EDO	C	1901	-	3,3,3	0.42	0	2,2,2	0.51	0
3	EDO	A	1903	-	3,3,3	0.67	0	2,2,2	0.36	0
3	EDO	A	1902	-	3,3,3	0.64	0	2,2,2	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1901	-	-	1/1/1/1	-
3	EDO	D	901	-	-	0/1/1/1	-
3	EDO	C	1901	-	-	0/1/1/1	-
3	EDO	A	1903	-	-	0/1/1/1	-
3	EDO	A	1902	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1901	EDO	O1-C1-C2-O2
3	A	1902	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1901	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/389 (94%)	0.01	10 (2%) 56 59	11, 25, 49, 69	15 (4%)
1	C	359/389 (92%)	0.00	15 (4%) 40 42	11, 25, 48, 77	16 (4%)
2	B	23/27 (85%)	0.09	2 (8%) 16 16	14, 25, 45, 61	1 (4%)
2	D	23/27 (85%)	0.05	1 (4%) 40 41	14, 23, 33, 56	3 (13%)
All	All	773/832 (92%)	0.01	28 (3%) 46 48	11, 25, 49, 77	35 (4%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	822	THR	4.5
1	C	1790	PHE	4.1
1	A	1790	PHE	4.0
1	C	1788	ASN	3.8
1	A	1639	VAL	3.7
1	C	1498	VAL	3.5
1	C	1471	PHE	3.2
1	C	1789	GLU	3.1
1	A	1640	LYS	3.0
1	C	1787	VAL	3.0
1	C	1813	PRO	3.0
1	A	1813	PRO	2.9
2	B	822	THR	2.7
1	C	1656	THR	2.7
1	A	1471	PHE	2.6
1	A	1814	GLN	2.6
1	A	1637	SER	2.5
1	C	1502	LEU	2.5
1	A	1500	VAL	2.5
1	C	1630	HIS	2.5
1	C	1757	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1502	LEU	2.3
1	C	1729	ARG	2.2
1	A	1787	VAL	2.1
1	C	1499	ALA	2.0
2	B	800	GLN	2.0
1	C	1500	VAL	2.0
1	C	1851	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	D	901	4/4	0.78	0.21	55,57,58,60	0
3	EDO	A	1903	4/4	0.80	0.16	48,49,52,54	0
3	EDO	A	1902	4/4	0.83	0.13	40,42,46,46	0
3	EDO	C	1901	4/4	0.86	0.14	46,49,49,51	0
3	EDO	A	1901	4/4	0.87	0.13	49,55,59,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.