



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2026 – 01:09 PM UTC

PDB ID : 6KUZ / pdb_00006kuz
Title : E.coli beta-galactosidase (E537Q) in complex with fluorescent probe KSL01
Authors : Chen, X.; Hu, Y.L.; Li, X.K.; Guo, Y.; Li, J.
Deposited on : 2019-09-03
Resolution : 2.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.



Whole archive
(#Entries)

The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Quality of chain

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 33232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1021	Total	C	N	O	S	0	3	0
			8164	5167	1446	1513	38			
1	B	1021	Total	C	N	O	S	0	3	0
			8156	5161	1450	1508	37			
1	C	1021	Total	C	N	O	S	0	0	0
			8066	5113	1429	1487	37			
1	D	1020	Total	C	N	O	S	0	0	0
			8071	5115	1432	1487	37			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	537	GLN	GLU	engineered mutation	UNP P00722
B	537	GLN	GLU	engineered mutation	UNP P00722
C	537	GLN	GLU	engineered mutation	UNP P00722
D	537	GLN	GLU	engineered mutation	UNP P00722

- Molecule 2 is 3-(1,3-benzothiazol-2-yl)-2-[[4-[(2 {S},3 {R},4 {S},5 {R},6 {R})-6-(hydroxymethyl)-3,4,5-tris(oxidanyl)oxan-2-yl]oxyphenyl]methoxy]-5-methyl-benzaldehyde (CCD ID: DVL) (formula: C₂₈H₂₇NO₈S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			38	28	1	8	1		
2	B	1	Total	C	N	O	S	0	0
			38	28	1	8	1		
2	C	1	Total	C	N	O	S	0	0
			38	28	1	8	1		
2	D	1	Total	C	N	O	S	0	0
			38	28	1	8	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Na	0	0
			3	3		
3	B	3	Total	Na	0	0
			3	3		
3	C	3	Total	Na	0	0
			3	3		
3	D	3	Total	Na	0	0
			3	3		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	B	1	Total 4	C 2	O 1	S 1	0	0
4	C	1	Total 4	C 2	O 1	S 1	0	0
4	C	1	Total 4	C 2	O 1	S 1	0	0
4	C	1	Total 4	C 2	O 1	S 1	0	0
4	C	1	Total 4	C 2	O 1	S 1	0	0
4	C	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0
4	D	1	Total 4	C 2	O 1	S 1	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total 2	Mg 2	0	0
5	B	2	Total 2	Mg 2	0	0
5	C	2	Total 2	Mg 2	0	0
5	D	2	Total 2	Mg 2	0	0

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total 6	C 3	O 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	132	Total 132	O 132	0	0
7	B	121	Total 121	O 121	0	0
7	C	91	Total 91	O 91	0	0
7	D	113	Total 113	O 113	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-galactosidase




• Molecule 1: Beta-galactosidase




- Molecule 1: Beta-galactosidase

Chain C:  89% 11%



- Molecule 1: Beta-galactosidase

Chain D:  86% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	248.17Å 85.51Å 243.05Å 90.00° 94.09° 90.00°	Depositor
Resolution (Å)	29.50 – 2.83 29.50 – 2.83	Depositor EDS
% Data completeness (in resolution range)	96.9 (29.50-2.83) 96.9 (29.50-2.83)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.85Å)	Xtriage
Refinement program	REFMAC 7.0.076	Depositor
R, R_{free}	0.182 , 0.224 0.185 , 0.224	Depositor DCC
R_{free} test set	5871 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	41.3	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 31.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	33232	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, DMS, MG, GOL, DVL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	0/8406	1.06	0/11475
1	B	0.91	0/8396	1.05	2/11459 (0.0%)
1	C	0.93	0/8307	1.05	0/11349
1	D	0.92	0/8312	1.06	2/11357 (0.0%)
All	All	0.92	0/33421	1.06	4/45640 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	752	GLY	CA-C-N	5.30	127.64	120.38
1	D	752	GLY	C-N-CA	5.30	127.64	120.38
1	B	859	ASP	CA-CB-CG	5.29	117.89	112.60
1	B	323	ILE	N-CA-C	-5.05	106.88	111.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8164	0	7732	71	0
1	B	8156	0	7731	80	0
1	C	8066	0	7609	69	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	8071	0	7619	83	0
2	A	38	0	0	2	0
2	B	38	0	0	4	0
2	C	38	0	0	2	0
2	D	38	0	0	2	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
4	A	52	0	78	3	0
4	B	32	0	48	0	0
4	C	20	0	30	1	0
4	D	36	0	54	5	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	D	2	0	0	0	0
6	D	6	0	8	0	0
7	A	132	0	0	3	0
7	B	121	0	0	3	0
7	C	91	0	0	2	0
7	D	113	0	0	1	0
All	All	33232	0	30909	302	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 302 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:773:LYS:HE2	1:B:775:GLN:HE22	1.44	0.82
1:D:290:THR:H	4:D:1108:DMS:H21	1.46	0.80
1:B:305:ILE:HD11	1:B:645:ARG:HB3	1.66	0.77
1:B:545:SER:HB2	1:B:791:ASN:HD21	1.53	0.73
1:A:427:THR:HA	1:A:436:MET:HE1	1.70	0.73

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1022/1024 (100%)	983 (96%)	39 (4%)	0	100	100
1	B	1022/1024 (100%)	983 (96%)	39 (4%)	0	100	100
1	C	1019/1024 (100%)	974 (96%)	44 (4%)	1 (0%)	48	69
1	D	1018/1024 (99%)	978 (96%)	40 (4%)	0	100	100
All	All	4081/4096 (100%)	3918 (96%)	162 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	684	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	863/876 (98%)	849 (98%)	14 (2%)	55	76
1	B	859/876 (98%)	841 (98%)	18 (2%)	47	70
1	C	842/876 (96%)	830 (99%)	12 (1%)	59	78
1	D	844/876 (96%)	829 (98%)	15 (2%)	51	74
All	All	3408/3504 (97%)	3349 (98%)	59 (2%)	53	75

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	690	SER
1	D	710	GLU
1	C	546	LEU
1	D	690	SER
1	D	519	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	363	HIS
1	D	887	GLN
1	D	424	ASN
1	D	757	GLN
1	D	965	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 60 ligands modelled in this entry, 20 are monoatomic - leaving 40 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DMS	D	1108	-	3,3,3	0.33	0	3,3,3	0.21	0
4	DMS	B	1109	-	3,3,3	0.37	0	3,3,3	0.05	0
4	DMS	D	1107	-	3,3,3	0.34	0	3,3,3	0.06	0
4	DMS	C	1108	-	3,3,3	0.34	0	3,3,3	0.08	0
4	DMS	A	1109	-	3,3,3	0.40	0	3,3,3	0.25	0
4	DMS	A	1113	-	3,3,3	0.37	0	3,3,3	0.05	0
4	DMS	B	1107	-	3,3,3	0.38	0	3,3,3	0.20	0
4	DMS	C	1105	-	3,3,3	0.42	0	3,3,3	0.05	0
4	DMS	A	1106	-	3,3,3	0.38	0	3,3,3	0.17	0
4	DMS	C	1106	-	3,3,3	0.46	0	3,3,3	0.15	0
4	DMS	A	1116	-	3,3,3	0.36	0	3,3,3	0.23	0
4	DMS	D	1106	-	3,3,3	0.37	0	3,3,3	0.24	0
4	DMS	A	1115	-	3,3,3	0.35	0	3,3,3	0.38	0
4	DMS	B	1108	-	3,3,3	0.46	0	3,3,3	0.41	0
4	DMS	B	1106	-	3,3,3	0.54	0	3,3,3	0.29	0
4	DMS	A	1112	-	3,3,3	0.29	0	3,3,3	0.17	0
4	DMS	B	1112	-	3,3,3	0.40	0	3,3,3	0.08	0
4	DMS	D	1113	-	3,3,3	0.18	0	3,3,3	0.18	0
4	DMS	D	1110	-	3,3,3	0.38	0	3,3,3	0.13	0
4	DMS	A	1108	-	3,3,3	0.36	0	3,3,3	0.14	0
2	DVL	C	1101	5,3	42,42,42	1.62	3 (7%)	60,60,60	2.23	19 (31%)
4	DMS	B	1105	-	3,3,3	0.41	0	3,3,3	0.12	0
4	DMS	D	1112	-	3,3,3	0.38	0	3,3,3	0.15	0
4	DMS	D	1109	-	3,3,3	0.45	0	3,3,3	0.11	0
4	DMS	D	1111	-	3,3,3	0.36	0	3,3,3	0.19	0
2	DVL	D	1101	3	42,42,42	1.53	2 (4%)	60,60,60	1.86	12 (20%)
2	DVL	A	1101	3	42,42,42	1.55	2 (4%)	60,60,60	1.92	14 (23%)
4	DMS	C	1109	-	3,3,3	0.34	0	3,3,3	0.13	0
4	DMS	A	1105	3	3,3,3	0.41	0	3,3,3	0.16	0
4	DMS	B	1110	-	3,3,3	0.38	0	3,3,3	0.11	0
4	DMS	A	1117	-	3,3,3	0.38	0	3,3,3	0.11	0
4	DMS	B	1111	-	3,3,3	0.40	0	3,3,3	0.07	0
4	DMS	D	1105	-	3,3,3	0.42	0	3,3,3	0.16	0
6	GOL	D	1114	-	5,5,5	0.14	0	5,5,5	0.34	0
2	DVL	B	1101	5	42,42,42	1.59	3 (7%)	60,60,60	1.80	16 (26%)
4	DMS	C	1107	-	3,3,3	0.28	0	3,3,3	0.24	0
4	DMS	A	1110	-	3,3,3	0.25	0	3,3,3	0.30	0
4	DMS	A	1114	-	3,3,3	0.30	0	3,3,3	0.24	0
4	DMS	A	1111	-	3,3,3	0.40	0	3,3,3	0.24	0
4	DMS	A	1107	-	3,3,3	0.43	0	3,3,3	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DVL	C	1101	5,3	-	12/17/37/37	0/5/5/5
6	GOL	D	1114	-	-	0/4/4/4	-
2	DVL	B	1101	5	-	10/17/37/37	0/5/5/5
2	DVL	D	1101	3	-	12/17/37/37	0/5/5/5
2	DVL	A	1101	3	-	10/17/37/37	0/5/5/5

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	DVL	C10-C08	-6.92	1.36	1.47
2	D	1101	DVL	C10-C08	-6.68	1.37	1.47
2	C	1101	DVL	C10-C08	-6.63	1.37	1.47
2	B	1101	DVL	C10-C08	-6.55	1.37	1.47
2	B	1101	DVL	C14-C37	-4.69	1.36	1.47

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1101	DVL	O16-C29-C28	6.25	116.26	107.16
2	A	1101	DVL	C11-C10-C08	-6.22	110.03	120.08
2	A	1101	DVL	C15-C10-C08	5.95	130.11	122.06
2	C	1101	DVL	C15-C10-C08	5.90	130.04	122.06
2	C	1101	DVL	C11-C10-C08	-5.43	111.31	120.08

There are no chirality outliers.

5 of 44 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	DVL	S07-C08-C10-C15
2	A	1101	DVL	C15-C14-C37-O38
2	A	1101	DVL	C28-C29-O16-C23
2	B	1101	DVL	N09-C08-C10-C15
2	B	1101	DVL	S07-C08-C10-C15


There are no ring outliers.


12 monomers are involved in 19 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1108	DMS	2	0
4	A	1115	DMS	1	0
4	A	1112	DMS	1	0
4	D	1113	DMS	1	0
4	D	1110	DMS	1	0
2	C	1101	DVL	2	0
4	D	1112	DMS	1	0
2	D	1101	DVL	2	0
2	A	1101	DVL	2	0
4	C	1109	DMS	1	0
4	A	1105	DMS	1	0
2	B	1101	DVL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1021/1024 (99%)	-0.43	4 (0%) 88 87	15, 35, 56, 84	3 (0%)
1	B	1021/1024 (99%)	-0.40	3 (0%) 90 89	14, 36, 58, 106	3 (0%)
1	C	1021/1024 (99%)	-0.31	4 (0%) 88 87	25, 40, 65, 115	0
1	D	1020/1024 (99%)	-0.36	7 (0%) 84 80	24, 37, 61, 112	0
All	All	4083/4096 (99%)	-0.37	18 (0%) 88 87	14, 37, 61, 115	6 (0%)

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	3	ILE	4.0
1	D	3	ILE	3.1
1	D	733	ALA	2.7
1	A	653[A]	HIS	2.6
1	A	1023	LYS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	DMS	B	1108	4/4	0.68	0.23	62,63,67,76	0
4	DMS	B	1107	4/4	0.70	0.27	79,81,85,89	0
2	DVL	A	1101	38/38	0.76	0.26	56,95,149,156	0
2	DVL	C	1101	38/38	0.77	0.28	67,124,184,192	0
4	DMS	B	1111	4/4	0.77	0.23	79,91,92,95	0
2	DVL	B	1101	38/38	0.78	0.25	69,117,181,184	0
4	DMS	D	1111	4/4	0.78	0.20	58,66,69,70	0
4	DMS	A	1111	4/4	0.79	0.21	82,84,97,101	0
4	DMS	A	1117	4/4	0.79	0.24	84,85,92,95	0
2	DVL	D	1101	38/38	0.79	0.25	82,122,165,168	0
4	DMS	B	1110	4/4	0.82	0.22	82,91,94,96	0
4	DMS	D	1109	4/4	0.83	0.22	61,63,67,73	0
4	DMS	A	1107	4/4	0.84	0.24	69,78,81,82	0
4	DMS	A	1109	4/4	0.85	0.20	60,61,62,65	0
4	DMS	D	1108	4/4	0.85	0.21	61,65,67,72	0
3	NA	A	1104	1/1	0.85	0.09	60,60,60,60	0
4	DMS	A	1108	4/4	0.85	0.20	73,77,81,85	0
4	DMS	D	1112	4/4	0.85	0.18	57,64,68,72	0
4	DMS	C	1106	4/4	0.86	0.20	66,78,80,85	0
4	DMS	A	1110	4/4	0.87	0.16	82,84,88,93	0
4	DMS	A	1114	4/4	0.87	0.16	63,65,70,71	0
4	DMS	A	1112	4/4	0.88	0.18	66,69,70,73	0
4	DMS	B	1106	4/4	0.90	0.17	58,58,59,66	0
4	DMS	B	1112	4/4	0.91	0.16	65,65,71,73	0
4	DMS	D	1110	4/4	0.91	0.19	65,71,77,78	0
4	DMS	C	1108	4/4	0.91	0.15	64,64,71,72	0
4	DMS	C	1105	4/4	0.91	0.18	61,69,71,71	0
4	DMS	D	1105	4/4	0.92	0.17	67,78,79,81	0
4	DMS	A	1105	4/4	0.92	0.17	62,68,73,74	0
4	DMS	A	1115	4/4	0.92	0.17	47,51,52,56	0
4	DMS	A	1106	4/4	0.92	0.15	63,65,70,70	0
4	DMS	C	1107	4/4	0.92	0.14	67,70,71,74	0
4	DMS	A	1113	4/4	0.92	0.12	55,57,67,70	0
4	DMS	D	1106	4/4	0.93	0.14	53,59,59,63	0
4	DMS	D	1113	4/4	0.93	0.15	66,73,79,82	0
4	DMS	C	1109	4/4	0.94	0.12	50,56,57,60	0
4	DMS	A	1116	4/4	0.94	0.13	51,56,57,57	0
4	DMS	B	1109	4/4	0.94	0.16	63,71,73,76	0
4	DMS	D	1107	4/4	0.94	0.15	67,77,80,83	0
4	DMS	B	1105	4/4	0.94	0.16	55,60,60,63	0
3	NA	D	1104	1/1	0.95	0.06	50,50,50,50	0
3	NA	B	1104	1/1	0.95	0.06	46,46,46,46	0
6	GOL	D	1114	6/6	0.95	0.07	54,59,62,62	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	D	1116	1/1	0.96	0.13	42,42,42,42	0
3	NA	C	1102	1/1	0.96	0.07	37,37,37,37	0
3	NA	B	1103	1/1	0.97	0.05	30,30,30,30	0
3	NA	D	1102	1/1	0.97	0.04	34,34,34,34	0
3	NA	D	1103	1/1	0.98	0.06	34,34,34,34	0
5	MG	A	1119	1/1	0.98	0.07	31,31,31,31	0
3	NA	C	1104	1/1	0.98	0.05	41,41,41,41	0
3	NA	C	1103	1/1	0.98	0.08	28,28,28,28	0
5	MG	A	1118	1/1	0.99	0.03	36,36,36,36	0
3	NA	A	1102	1/1	0.99	0.05	44,44,44,44	0
5	MG	B	1113	1/1	0.99	0.03	37,37,37,37	0
5	MG	B	1114	1/1	0.99	0.04	40,40,40,40	0
5	MG	C	1110	1/1	0.99	0.03	40,40,40,40	0
5	MG	C	1111	1/1	0.99	0.06	42,42,42,42	0
5	MG	D	1115	1/1	0.99	0.03	43,43,43,43	0
3	NA	B	1102	1/1	0.99	0.07	44,44,44,44	0
3	NA	A	1103	1/1	0.99	0.05	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around DVL C 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DVL B 1101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around DVL D 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.