



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:42 AM UTC

PDB ID : 3KV4 / pdb_00003kv4
Title : Structure of PHF8 in complex with histone H3
Authors : Horton, J.R.; Upadhyay, A.K.; Qi, H.H.; Zhang, X.; Shi, Y.; Cheng, X.
Deposited on : 2009-11-29
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	
2	B	24	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHD finger protein 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	432	Total	C	N	O	S	0	2	0
			3475	2224	591	637	23			

- Molecule 2 is a protein called Histone H3-like.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	14	Total	C	N	O	0	2	0
			125	74	29	22			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe	0	0
			1	1		

- Molecule 5 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ni	0	0
			1	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is N-OXALYLGLYCINE (CCD ID: OGA) (formula: C₄H₅NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			10	4	1	5		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	193	Total	O	0	0
			193	193		
8	B	12	Total	O	0	0
			12	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHD finger protein 8



• Molecule 2: Histone H3-like



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.50Å 73.50Å 210.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.72 – 2.19 34.72 – 2.19	Depositor EDS
% Data completeness (in resolution range)	93.0 (34.72-2.19) 93.1 (34.72-2.19)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.217 , 0.255 0.213 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3871	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI, EDO, MLY, ZN, FE2, OGA, M3L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3561	0.88	6/4832 (0.1%)
2	B	0.37	0/101	0.99	1/132 (0.8%)
All	All	0.39	0/3662	0.89	7/4964 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	12	GLY	N-CA-C	-6.58	105.40	114.67
1	A	255	VAL	N-CA-C	6.11	116.92	108.12
1	A	128	LEU	N-CA-C	5.74	117.54	111.28
1	A	107	THR	N-CA-C	-5.58	102.50	110.59
1	A	27	GLN	N-CA-C	-5.42	105.43	113.40
1	A	58	GLY	CA-C-N	5.35	126.53	119.84
1	A	58	GLY	C-N-CA	5.35	126.53	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3475	0	3365	98	0
2	B	125	0	142	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	52	0	78	7	0
7	A	10	0	3	0	0
8	A	193	0	0	0	0
8	B	12	0	0	0	0
All	All	3871	0	3588	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:LYS:HG2	1:A:79:PRO:HD2	1.43	0.99
1:A:21:ILE:HG22	2:B:3:THR:HG22	1.53	0.89
1:A:336:HIS:H	1:A:343:GLN:HE22	1.32	0.77
2:B:1:ALA:O	2:B:3:THR:HG23	1.86	0.75
1:A:208:LYS:HE2	1:A:212:LYS:HE3	1.70	0.73
1:A:257:TYR:OH	1:A:264:LYS:HE3	1.88	0.72
1:A:291:MET:HE3	1:A:296:GLN:NE2	2.05	0.72
1:A:382:LEU:HD12	1:A:389:PRO:HG3	1.73	0.70
1:A:215:TRP:CD2	1:A:371:VAL:HG21	2.26	0.70
1:A:78:LYS:CG	1:A:79:PRO:HD2	2.20	0.67
1:A:107:THR:HG22	1:A:110:GLN:HG3	1.80	0.62
1:A:35:VAL:HB	6:A:461:EDO:H22	1.82	0.62
1:A:406:ALA:HA	1:A:409:ARG:NH1	2.16	0.61
1:A:433:LEU:O	1:A:437:ILE:HD13	2.01	0.61
1:A:98:SER:O	1:A:101:GLU:HG2	2.03	0.59
1:A:383:ARG:HG2	1:A:383:ARG:HH11	1.68	0.59
1:A:114:GLU:CD	1:A:114:GLU:H	2.11	0.59
1:A:26:CYS:O	1:A:27:GLN:HB2	2.02	0.58
1:A:107:THR:HG23	1:A:109:ASN:H	1.69	0.57
1:A:145:ARG:HD2	1:A:180:TYR:CE2	2.40	0.57
1:A:29:TRP:CZ3	2:B:4:M3L:HG2	2.41	0.56
1:A:229:PRO:HB3	1:A:362:PRO:HG2	1.87	0.55
1:A:333:ASN:OD1	2:B:9:MLY:HH22	2.05	0.55
1:A:106:PRO:HD2	1:A:125:ILE:HG23	1.89	0.55
1:A:351:LYS:HG3	1:A:360:ARG:HH22	1.71	0.54
1:A:24:ASP:CG	1:A:47:LEU:HD11	2.32	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:379:PHE:CE1	1:A:389:PRO:HG2	2.43	0.53
1:A:425:ARG:HG3	1:A:428:GLN:H	1.73	0.52
1:A:374:HIS:CE1	1:A:378:ILE:HD11	2.44	0.52
1:A:54:GLU:HG3	1:A:59:PRO:HA	1.92	0.51
1:A:221:PRO:HG2	1:A:224:CYS:HB3	1.92	0.51
1:A:365:GLU:HG2	1:A:407:TRP:CH2	2.46	0.50
1:A:116:LEU:CD1	1:A:213:LEU:HD11	2.41	0.50
1:A:120:SER:O	1:A:121:PHE:C	2.54	0.50
1:A:385:ASN:O	1:A:387:ARG:HG3	2.11	0.50
1:A:121:PHE:CE2	1:A:314:PRO:HG3	2.47	0.49
1:A:380:ARG:HG2	1:A:380:ARG:HH11	1.77	0.49
1:A:52:ASN:O	1:A:55:VAL:HG22	2.13	0.49
1:A:336:HIS:N	1:A:343:GLN:HE22	2.07	0.49
1:A:236:LEU:N	1:A:236:LEU:HD22	2.28	0.49
1:A:437:ILE:O	1:A:441:GLU:HG3	2.13	0.49
1:A:40:GLU:OE1	1:A:40:GLU:N	2.36	0.48
1:A:144:VAL:HG22	1:A:180:TYR:CZ	2.47	0.48
1:A:62:MET:HG3	6:A:460:EDO:O2	2.12	0.48
1:A:389:PRO:HB3	1:A:393:LEU:HD23	1.95	0.48
1:A:48:TYR:OH	6:A:461:EDO:H22	2.14	0.48
1:A:336:HIS:H	1:A:343:GLN:NE2	2.07	0.47
1:A:252:GLY:O	1:A:315:THR:HG23	2.14	0.47
1:A:325:VAL:O	1:A:326:ASP:C	2.56	0.47
1:A:45:ILE:HB	2:B:1:ALA:HB2	1.98	0.46
1:A:379:PHE:CD1	1:A:389:PRO:HG2	2.51	0.46
1:A:25:MET:HB2	1:A:53:CYS:SG	2.56	0.46
1:A:347:TYR:CE1	1:A:360:ARG:NH2	2.84	0.46
1:A:178:TYR:O	1:A:181:SER:HB3	2.16	0.45
1:A:313:ILE:HA	1:A:314:PRO:HD3	1.82	0.45
1:A:134:GLY:HA2	6:A:458:EDO:H21	1.99	0.45
1:A:282:TRP:HD1	1:A:291:MET:HE2	1.82	0.45
1:A:336:HIS:CE1	1:A:338:LEU:HB2	2.51	0.45
2:B:13:GLY:O	2:B:14:LYS:C	2.60	0.45
1:A:323:THR:HG22	1:A:325:VAL:O	2.16	0.45
1:A:16:VAL:HG22	1:A:16:VAL:O	2.17	0.45
1:A:351:LYS:HG3	1:A:360:ARG:NH2	2.32	0.45
1:A:55:VAL:HG21	6:A:462:EDO:H12	1.99	0.44
1:A:107:THR:HG23	1:A:109:ASN:N	2.33	0.44
1:A:134:GLY:H	6:A:458:EDO:H22	1.82	0.44
1:A:114:GLU:CD	1:A:114:GLU:N	2.75	0.44
1:A:154:ASP:O	1:A:171:LYS:HD2	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:LEU:HD12	1:A:371:VAL:HG22	1.98	0.44
1:A:380:ARG:HG2	1:A:380:ARG:NH1	2.31	0.44
1:A:86:THR:O	1:A:90:GLU:HG3	2.18	0.44
1:A:89[A]:ARG:HG2	1:A:89[A]:ARG:HH11	1.82	0.44
1:A:346:ALA:O	1:A:350:GLU:HG3	2.18	0.44
1:A:383:ARG:HG2	1:A:383:ARG:NH1	2.32	0.44
1:A:116:LEU:HD12	1:A:213:LEU:HD11	2.00	0.43
1:A:379:PHE:HE1	1:A:440:VAL:HG21	1.83	0.43
1:A:215:TRP:CE2	1:A:371:VAL:HG21	2.53	0.43
1:A:267:TYR:O	1:A:319:HIS:HA	2.17	0.43
1:A:268:LEU:HD23	1:A:319:HIS:HB3	2.00	0.43
1:A:380:ARG:NH1	1:A:436:GLU:OE2	2.51	0.43
1:A:63:LYS:O	1:A:64:LYS:CB	2.66	0.43
1:A:208:LYS:HE2	1:A:212:LYS:CE	2.45	0.42
1:A:191:ILE:HA	1:A:236:LEU:HD13	2.00	0.42
1:A:134:GLY:H	6:A:458:EDO:C2	2.32	0.42
1:A:291:MET:CE	1:A:296:GLN:NE2	2.80	0.42
1:A:49:HIS:ND1	1:A:59:PRO:O	2.51	0.42
1:A:103:ILE:HG22	1:A:124:PRO:HB2	2.00	0.42
1:A:421:PRO:HB2	1:A:424:VAL:HG23	2.00	0.42
1:A:425:ARG:CG	1:A:428:GLN:HB3	2.49	0.42
2:B:11:THR:C	2:B:13:GLY:H	2.27	0.42
1:A:78:LYS:HG2	1:A:79:PRO:CD	2.32	0.42
1:A:107:THR:HG22	1:A:110:GLN:CG	2.49	0.41
1:A:108:GLY:HA3	1:A:131:ASP:O	2.20	0.41
1:A:137:LEU:HD23	1:A:204:VAL:HG13	2.02	0.41
1:A:150:TYR:CZ	1:A:203:LEU:HD11	2.54	0.41
1:A:107:THR:HG22	1:A:110:GLN:H	1.85	0.41
1:A:119:ASN:HD22	1:A:119:ASN:H	1.69	0.41
1:A:287:ASN:O	1:A:288:GLN:C	2.64	0.41
1:A:119:ASN:ND2	1:A:119:ASN:C	2.75	0.41
1:A:105:LYS:HE2	1:A:126:LEU:HD23	2.03	0.40
1:A:291:MET:CE	1:A:296:GLN:HE22	2.35	0.40
1:A:136:THR:HB	1:A:205:GLU:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/447 (96%)	414 (96%)	12 (3%)	4 (1%)	14	14
2	B	12/24 (50%)	10 (83%)	2 (17%)	0	100	100
All	All	442/471 (94%)	424 (96%)	14 (3%)	4 (1%)	14	14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	VAL
1	A	288	GLN
1	A	223	GLU
1	A	224	CYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/407 (94%)	379 (99%)	4 (1%)	68	81
2	B	10/15 (67%)	10 (100%)	0	100	100
All	All	393/422 (93%)	389 (99%)	4 (1%)	68	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	THR
1	A	119	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	232	GLN
1	A	236	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	ASN
1	A	288	GLN
1	A	289	ASN
1	A	296	GLN
1	A	307	GLN
1	A	343	GLN
1	A	374	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	M3L	B	4	2	10,11,12	0.68	0	9,14,16	0.46	0
2	MLY	B	9	2	9,10,11	0.51	0	6,11,13	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M3L	B	4	2	-	1/9/10/12	-
2	MLY	B	9	2	-	0/8/9/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4	M3L	C-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	4	M3L	1	0
2	B	9	MLY	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	A	463	-	3,3,3	0.57	0	2,2,2	0.37	0
6	EDO	A	455	-	3,3,3	0.47	0	2,2,2	0.32	0
6	EDO	A	457	-	3,3,3	0.34	0	2,2,2	0.45	0
6	EDO	A	456	-	3,3,3	0.46	0	2,2,2	0.32	0
6	EDO	A	452	-	3,3,3	0.35	0	2,2,2	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	A	453	-	3,3,3	0.42	0	2,2,2	0.34	0
6	EDO	A	459	-	3,3,3	0.43	0	2,2,2	0.36	0
6	EDO	A	458	-	3,3,3	0.43	0	2,2,2	0.29	0
6	EDO	A	464	-	3,3,3	0.47	0	2,2,2	0.42	0
6	EDO	A	462	-	3,3,3	0.49	0	2,2,2	0.44	0
7	OGA	A	5798	4	9,9,9	1.50	1 (11%)	8,11,11	1.72	3 (37%)
6	EDO	A	454	-	3,3,3	0.44	0	2,2,2	0.28	0
6	EDO	A	461	-	3,3,3	0.49	0	2,2,2	0.36	0
6	EDO	A	460	-	3,3,3	0.44	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	463	-	-	0/1/1/1	-
6	EDO	A	455	-	-	0/1/1/1	-
6	EDO	A	457	-	-	0/1/1/1	-
6	EDO	A	456	-	-	1/1/1/1	-
6	EDO	A	452	-	-	1/1/1/1	-
6	EDO	A	453	-	-	0/1/1/1	-
6	EDO	A	459	-	-	1/1/1/1	-
6	EDO	A	458	-	-	0/1/1/1	-
6	EDO	A	464	-	-	0/1/1/1	-
6	EDO	A	462	-	-	1/1/1/1	-
7	OGA	A	5798	4	-	0/8/9/9	-
6	EDO	A	454	-	-	0/1/1/1	-
6	EDO	A	461	-	-	0/1/1/1	-
6	EDO	A	460	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	5798	OGA	C2-C1	-3.65	1.50	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	5798	OGA	O2'-C2-C1	-3.23	117.12	121.36
7	A	5798	OGA	C4-N1-C2	2.92	125.80	121.29
7	A	5798	OGA	O2-C1-O1	-2.05	119.02	123.90

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	462	EDO	O1-C1-C2-O2
6	A	460	EDO	O1-C1-C2-O2
6	A	459	EDO	O1-C1-C2-O2
6	A	452	EDO	O1-C1-C2-O2
6	A	456	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	458	EDO	3	0
6	A	462	EDO	1	0
6	A	461	EDO	2	0
6	A	460	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	432/447 (96%)	0.55	49 (11%) 10 8	8, 23, 51, 64	2 (0%)
2	B	12/24 (50%)	0.85	2 (16%) 4 3	9, 22, 43, 47	2 (16%)
All	All	444/471 (94%)	0.56	51 (11%) 9 7	8, 23, 51, 64	4 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	447	ASN	5.9
1	A	62	MET	5.3
1	A	56	LEU	5.3
1	A	57	HIS	4.3
1	A	64	LYS	4.2
1	A	55	VAL	4.2
1	A	444	PHE	4.0
1	A	446	GLN	3.8
1	A	58	GLY	3.8
1	A	78	LYS	3.7
1	A	445	GLN	3.7
1	A	137	LEU	3.6
1	A	61	ILE	3.6
1	A	145	ARG	3.6
2	B	14	LYS	3.6
1	A	26	CYS	3.5
1	A	222	GLU	3.4
1	A	226	PHE	3.3
1	A	224	CYS	3.3
2	B	13	GLY	3.2
1	A	25	MET	3.1
1	A	51	PRO	3.1
1	A	149	HIS	3.1
1	A	388	HIS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	VAL	3.0
1	A	49	HIS	3.0
1	A	3	SER	2.8
1	A	43	ALA	2.8
1	A	89[A]	ARG	2.7
1	A	384	GLU	2.7
1	A	60	SER	2.7
1	A	47	LEU	2.6
1	A	182	GLY	2.6
1	A	59	PRO	2.6
1	A	184	ARG	2.6
1	A	53	CYS	2.6
1	A	52	ASN	2.5
1	A	387	ARG	2.5
1	A	50	CYS	2.4
1	A	63	LYS	2.4
1	A	409	ARG	2.3
1	A	204	VAL	2.2
1	A	289	ASN	2.2
1	A	152	GLY	2.2
1	A	183	LYS	2.2
1	A	438	ARG	2.1
1	A	46	ASP	2.1
1	A	223	GLU	2.1
1	A	79	PRO	2.1
1	A	380	ARG	2.1
1	A	144	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	M3L	B	4	12/13	0.92	0.10	13,17,18,18	0
2	MLY	B	9	11/12	0.95	0.07	10,14,18,19	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	A	464	4/4	0.65	0.20	39,40,40,41	0
6	EDO	A	458	4/4	0.69	0.22	40,41,44,45	0
6	EDO	A	461	4/4	0.73	0.22	35,36,36,37	0
6	EDO	A	463	4/4	0.76	0.16	39,39,41,41	0
6	EDO	A	459	4/4	0.79	0.18	46,47,48,50	0
6	EDO	A	456	4/4	0.80	0.18	37,37,37,37	0
6	EDO	A	462	4/4	0.83	0.12	39,41,41,43	0
6	EDO	A	454	4/4	0.85	0.13	35,36,36,38	0
6	EDO	A	457	4/4	0.85	0.17	41,41,41,43	0
6	EDO	A	453	4/4	0.88	0.13	30,32,34,34	0
6	EDO	A	460	4/4	0.90	0.13	27,29,30,30	0
7	OGA	A	5798	10/10	0.91	0.11	26,34,38,40	0
6	EDO	A	455	4/4	0.93	0.10	28,31,32,33	0
6	EDO	A	452	4/4	0.96	0.09	13,14,14,19	0
3	ZN	A	449	1/1	0.96	0.06	37,37,37,37	0
5	NI	A	451	1/1	0.99	0.13	33,33,33,33	0
4	FE2	A	450	1/1	1.00	0.02	16,16,16,16	0
3	ZN	A	448	1/1	1.00	0.01	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.