



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 12:41 AM UTC

PDB ID : 3KVR / pdb_00003kvr
Title : Trapping of an oxocarbenium ion intermediate in UP crystals
Authors : Paul, D.; O'Leary, S.; Rajashankar, K.; Bu, W.; Toms, A.; Settembre, E.;
Sanders, J.; Begley, T.P.; Ealick, S.E.
Deposited on : 2009-11-30
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	 4% 62% 27% 5% • 5%
1	B	309	 3% 68% 23% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	URF	A	2001	-	X	-	-
2	URF	B	2011	-	X	-	-
4	R2G	A	311	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2238	1416	386	414	22			
1	B	293	Total	C	N	O	S	0	0	0
			2251	1422	390	417	22			

- Molecule 2 is 5-FLUOROURACIL (CCD ID: URF) (formula: $C_4H_3FN_2O_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			9	4	1	2	2		
2	B	1	Total	C	F	N	O	0	0
			9	4	1	2	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 2,5-anhydro-4-deoxy-D-erythro-pent-4-enitol (CCD ID: R2G) (formula: C₅H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	5	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	5	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	63	Total	O	0	0
			63	63		
5	B	57	Total	O	0	0
			57	57		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uridine phosphorylase



• Molecule 1: Uridine phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.42Å 82.42Å 258.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.60 78.52 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.7 ((Not available)-2.60) 99.4 (78.52-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.87 (at 2.61Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.167 , 0.205 0.182 , 0.222	Depositor DCC
R_{free} test set	1426 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	36.2	Xtriage
Anisotropy	0.696	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4653	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: R2G, SO4, URF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.49	11/2282 (0.5%)	1.41	11/3086 (0.4%)
1	B	1.46	5/2294 (0.2%)	1.44	19/3099 (0.6%)
All	All	1.48	16/4576 (0.3%)	1.42	30/6185 (0.5%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	111	VAL	CA-CB	9.50	1.58	1.54
1	A	101	PRO	CA-C	6.89	1.59	1.52
1	A	157	ALA	CA-C	-6.34	1.44	1.52
1	B	89	GLU	CA-CB	6.19	1.64	1.53
1	B	22	ASN	N-CA	-6.01	1.41	1.46
1	B	29	LYS	CA-CB	5.69	1.62	1.53
1	A	20	LEU	CA-C	-5.62	1.46	1.52
1	A	63	ARG	CG-CD	5.52	1.69	1.52
1	A	301	PHE	N-CA	-5.43	1.39	1.46
1	A	21	CYS	CA-C	-5.41	1.47	1.53
1	A	129	HIS	CA-C	5.35	1.60	1.53
1	A	111	VAL	N-CA	-5.27	1.40	1.46
1	B	272	LEU	CA-C	-5.18	1.46	1.52
1	B	265	ALA	CA-CB	-5.11	1.44	1.54
1	A	42	THR	CA-CB	5.10	1.61	1.53
1	A	91	THR	N-CA	-5.01	1.39	1.45

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	252	VAL	CB-CA-C	-9.99	98.43	112.22
1	B	252	VAL	CB-CA-C	-9.39	97.04	112.26
1	B	220	ASP	N-CA-CB	-8.52	97.74	110.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123	LYS	N-CA-C	-7.89	102.62	111.14
1	A	74	GLU	N-CA-C	-6.99	105.68	114.56
1	B	292	GLN	N-CA-C	-6.75	104.52	112.89
1	A	157	ALA	N-CA-C	-6.74	100.51	110.48
1	B	60	SER	CA-C-N	-6.22	112.43	119.28
1	B	60	SER	C-N-CA	-6.22	112.43	119.28
1	B	157	ALA	N-CA-C	-6.17	99.59	109.96
1	B	219	LEU	CA-C-N	6.06	131.89	122.49
1	B	219	LEU	C-N-CA	6.06	131.89	122.49
1	B	251	SER	N-CA-C	6.00	118.34	111.02
1	A	193	SER	N-CA-C	-5.95	104.87	111.36
1	A	153	ILE	CA-CB-CG2	5.90	120.53	110.50
1	A	240	ALA	CA-C-N	-5.86	113.29	122.73
1	A	240	ALA	C-N-CA	-5.86	113.29	122.73
1	B	153	ILE	CB-CA-C	-5.63	103.27	110.98
1	B	15	ASN	N-CA-C	5.57	117.82	108.96
1	A	252	VAL	N-CA-C	-5.55	105.11	110.72
1	B	193	SER	N-CA-C	-5.55	105.31	111.36
1	B	164	PRO	CB-CA-C	-5.50	105.68	112.89
1	A	102	VAL	N-CA-C	5.43	115.68	107.80
1	A	77	PHE	N-CA-C	-5.40	104.97	113.02
1	B	22	ASN	CA-C-N	5.32	124.93	119.56
1	B	22	ASN	C-N-CA	5.32	124.93	119.56
1	B	54	PHE	CA-C-N	-5.31	116.22	123.12
1	B	54	PHE	C-N-CA	-5.31	116.22	123.12
1	B	93	ARG	CB-CG-CD	5.17	123.19	111.30
1	A	158	VAL	N-CA-CB	-5.05	103.02	112.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2238	0	2220	68	0
1	B	2251	0	2230	58	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	9	0	3	2	0
2	B	9	0	3	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	8	0	0	4	0
4	B	8	0	0	1	0
5	A	63	0	0	0	0
5	B	57	0	0	2	0
All	All	4653	0	4456	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2001:URF:HN1	4:A:311:R2G:C1	1.69	1.03
1:A:222:ALA:H	1:B:119:HIS:HD2	1.06	0.98
1:A:153:ILE:HD11	1:A:186:VAL:HB	1.45	0.95
2:B:2011:URF:HN1	4:B:311:R2G:C1	1.81	0.92
1:B:74:GLU:OE2	1:B:303:LYS:NZ	2.04	0.90
1:A:119:HIS:HD2	1:B:222:ALA:H	1.28	0.80
1:A:222:ALA:H	1:B:119:HIS:CD2	1.95	0.79
1:A:222:ALA:N	1:B:119:HIS:HD2	1.83	0.76
1:A:73:MET:CB	1:A:78:ALA:HB2	2.16	0.76
1:A:279:GLN:HE22	1:B:32:ILE:H	1.35	0.74
1:B:67:PHE:O	1:B:71:VAL:HG23	1.89	0.73
1:A:79:HIS:HB3	1:A:80:PRO:HD3	1.70	0.72
1:B:30:GLU:HG3	1:B:41:SER:HB3	1.69	0.72
1:B:138:ILE:O	1:B:138:ILE:HG13	1.91	0.71
1:A:67:PHE:HD2	1:A:295:GLN:HG2	1.54	0.71
1:B:182:ASP:OD2	1:B:263:ARG:HD2	1.89	0.71
1:A:154:THR:HG1	1:A:206:THR:HG1	1.03	0.69
1:A:79:HIS:HB3	1:A:80:PRO:CD	2.25	0.67
1:B:63:ARG:HD2	1:B:291:GLN:HG2	1.77	0.66
1:B:35:HIS:HD2	1:B:120:GLU:OE2	1.78	0.66
1:A:67:PHE:CD2	1:A:295:GLN:HG2	2.30	0.66
1:B:166:PHE:CZ	1:B:252:VAL:HG13	2.31	0.66
1:A:73:MET:HB3	1:A:78:ALA:HB2	1.78	0.65
1:B:31:ASP:OD2	1:B:127:HIS:HE1	1.80	0.65
1:B:297:LEU:C	1:B:297:LEU:HD12	2.21	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:VAL:HG22	1:B:252:VAL:HG22	1.80	0.63
1:B:150:SER:O	1:B:270:THR:HB	1.98	0.63
1:A:107:HIS:HD2	1:A:108:GLY:O	1.82	0.62
1:A:166:PHE:CZ	1:A:252:VAL:HG13	2.35	0.62
1:B:195:GLU:OE1	1:B:304:LYS:HD2	2.01	0.61
1:B:15:ASN:HB2	1:B:132:GLY:N	2.16	0.61
1:A:55:VAL:HG11	1:A:121:LEU:HD21	1.83	0.61
1:B:124:LEU:C	1:B:124:LEU:HD23	2.27	0.60
1:B:291:GLN:HE21	1:B:291:GLN:CA	2.14	0.60
1:B:161:CYS:HB2	1:B:163:LYS:HE3	1.84	0.59
1:B:63:ARG:HD2	1:B:291:GLN:CG	2.32	0.59
1:B:269:VAL:HG22	1:B:270:THR:N	2.19	0.57
1:A:153:ILE:O	1:A:153:ILE:HG13	2.04	0.57
4:A:311:R2G:O5	1:B:35:HIS:HE1	1.88	0.57
1:B:269:VAL:HG22	1:B:270:THR:H	1.69	0.56
1:B:297:LEU:HD12	1:B:297:LEU:O	2.05	0.56
1:A:167:GLU:CD	1:A:174:ARG:HE	2.14	0.56
1:B:144:ILE:HD11	1:B:246:ILE:HD11	1.89	0.55
1:A:119:HIS:CD2	1:B:222:ALA:H	2.18	0.55
1:B:135:LEU:O	1:B:264:ALA:HA	2.07	0.54
1:A:107:HIS:HB3	1:A:117:MET:HE2	1.88	0.54
1:B:107:HIS:HD2	1:B:108:GLY:O	1.90	0.54
1:A:68:ILE:HG13	1:A:99:VAL:HG12	1.89	0.53
1:A:135:LEU:O	1:A:264:ALA:HA	2.08	0.53
1:B:15:ASN:HB2	1:B:132:GLY:CA	2.39	0.53
1:A:32:ILE:H	1:B:279:GLN:HE22	1.57	0.53
1:A:36:PHE:HB3	1:A:38:LEU:HD22	1.91	0.52
1:B:185:LEU:HD13	1:B:265:ALA:HB2	1.91	0.52
1:A:45:PHE:N	1:A:46:PRO:CD	2.72	0.51
1:A:158:VAL:HG22	1:A:162:PHE:C	2.35	0.51
1:A:308:GLN:O	1:A:309:ALA:HB2	2.11	0.51
1:A:97:PHE:O	1:A:103:LEU:HD12	2.11	0.51
1:A:185:LEU:HD13	1:A:265:ALA:HB2	1.94	0.50
1:A:111:VAL:HG13	1:A:252:VAL:CG2	2.42	0.50
1:A:111:VAL:HG13	1:A:252:VAL:HG22	1.92	0.50
1:B:199:PHE:N	5:B:331:HOH:O	2.33	0.50
1:A:178:ASN:ND2	1:A:180:ASP:H	2.10	0.49
1:A:159:ASP:HB2	1:A:160:PRO:CD	2.42	0.49
1:A:73:MET:HB2	1:A:78:ALA:HB2	1.94	0.49
1:B:211:ASP:OD2	1:B:214:GLU:N	2.45	0.49
1:A:36:PHE:O	1:A:37:SER:HB3	2.11	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84:TYR:N	1:B:84:TYR:CD2	2.80	0.49
1:A:107:HIS:HE1	1:A:137:ARG:HH11	1.61	0.49
1:B:304:LYS:O	1:B:308:GLN:HG3	2.12	0.49
1:A:31:ASP:OD2	1:A:127:HIS:HE1	1.96	0.48
1:A:45:PHE:HB2	1:A:46:PRO:HD3	1.94	0.48
1:A:223:LEU:HD12	1:B:21:CYS:HB2	1.96	0.48
1:A:107:HIS:CE1	1:A:137:ARG:HH11	2.32	0.47
1:A:176:VAL:O	1:A:176:VAL:HG23	2.14	0.47
1:A:20:LEU:CD2	1:A:122:ILE:HG22	2.45	0.47
1:B:138:ILE:O	1:B:138:ILE:CG1	2.63	0.47
4:A:311:R2G:C5	1:B:35:HIS:HE1	2.28	0.46
1:B:138:ILE:HA	1:B:267:VAL:O	2.16	0.46
1:A:124:LEU:C	1:A:124:LEU:HD23	2.40	0.46
1:B:211:ASP:HB3	1:B:215:GLY:HA3	1.97	0.45
1:A:20:LEU:HD23	1:A:122:ILE:HG22	1.98	0.45
1:A:119:HIS:HD2	1:B:222:ALA:N	2.07	0.45
1:B:15:ASN:CB	1:B:132:GLY:HA3	2.47	0.45
1:A:154:THR:OG1	1:A:206:THR:OG1	1.91	0.45
1:B:111:VAL:HG22	1:B:252:VAL:CG2	2.47	0.45
1:B:159:ASP:OD2	1:B:159:ASP:C	2.59	0.45
1:A:187:GLN:O	1:A:187:GLN:HG3	2.17	0.45
1:A:58:GLY:O	1:A:106:SER:HA	2.17	0.44
1:A:96:MET:HG3	1:A:105:VAL:HG22	1.98	0.44
1:A:55:VAL:HB	1:A:135:LEU:HD13	1.99	0.44
1:B:107:HIS:CD2	1:B:108:GLY:O	2.71	0.44
1:A:48:MET:CE	1:A:87:ILE:HD13	2.48	0.44
1:A:154:THR:HG21	1:A:206:THR:HB	1.99	0.44
1:A:162:PHE:N	1:A:162:PHE:CD2	2.86	0.43
1:A:308:GLN:O	1:A:309:ALA:CB	2.66	0.43
1:A:289:GLU:O	1:A:290:TYR:C	2.60	0.43
1:B:291:GLN:HE21	1:B:291:GLN:N	2.16	0.43
1:B:22:ASN:ND2	1:B:25:ILE:HD11	2.33	0.43
1:B:155:ARG:HD3	5:B:313:HOH:O	2.17	0.43
1:B:15:ASN:HB2	1:B:132:GLY:HA3	2.00	0.43
1:A:57:VAL:HA	1:A:105:VAL:O	2.19	0.43
1:A:166:PHE:CE1	1:A:252:VAL:HG13	2.54	0.43
1:B:22:ASN:HA	1:B:23:PRO:HD2	1.90	0.43
1:A:86:ASN:HB3	1:A:89:GLU:HG2	2.01	0.42
1:A:283:PRO:O	1:A:284:HIS:C	2.60	0.42
1:B:288:ALA:O	1:B:291:GLN:HB2	2.20	0.42
1:A:68:ILE:HD13	1:A:104:SER:HB2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:PHE:O	1:B:78:ALA:O	2.38	0.42
1:A:38:LEU:HD21	1:A:87:ILE:HD11	2.02	0.42
2:A:2001:URF:N1	4:A:311:R2G:C1	2.54	0.42
1:A:20:LEU:HD11	1:A:126:TYR:CD1	2.55	0.42
1:A:300:GLN:HE21	1:A:303:LYS:NZ	2.17	0.42
1:A:110:GLY:HA3	1:B:112:PRO:HB2	2.02	0.41
1:A:84:TYR:HA	1:A:85:PRO:HD2	1.91	0.41
1:A:196:LEU:HG	1:A:300:GLN:HG3	2.03	0.41
1:B:31:ASP:OD1	1:B:123:LYS:NZ	2.54	0.41
1:A:59:GLY:HA2	1:A:107:HIS:CD2	2.55	0.41
1:A:107:HIS:HB3	1:A:117:MET:CE	2.51	0.40
1:B:75:LEU:HB2	1:B:77:PHE:HD1	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/309 (94%)	276 (95%)	11 (4%)	4 (1%)	9	19
1	B	289/309 (94%)	279 (96%)	9 (3%)	1 (0%)	36	58
All	All	580/618 (94%)	555 (96%)	20 (3%)	5 (1%)	14	30

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	78	ALA
1	A	82	ALA
1	A	80	PRO
1	A	76	GLY
1	A	79	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/255 (94%)	215 (90%)	24 (10%)	7	16
1	B	242/255 (95%)	226 (93%)	16 (7%)	15	34
All	All	481/510 (94%)	441 (92%)	40 (8%)	10	23

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LEU
1	A	21	CYS
1	A	38	LEU
1	A	40	THR
1	A	63	ARG
1	A	71	VAL
1	A	83	ASP
1	A	87	ILE
1	A	153	ILE
1	A	158	VAL
1	A	162	PHE
1	A	174	ARG
1	A	185	LEU
1	A	186	VAL
1	A	189	LEU
1	A	202	VAL
1	A	203	VAL
1	A	223	LEU
1	A	247	GLU
1	A	252	VAL
1	A	279	GLN
1	A	281	SER
1	A	303	LYS
1	A	304	LYS
1	B	38	LEU
1	B	73	MET
1	B	79	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	93	ARG
1	B	158	VAL
1	B	185	LEU
1	B	186	VAL
1	B	189	LEU
1	B	203	VAL
1	B	247	GLU
1	B	251	SER
1	B	252	VAL
1	B	270	THR
1	B	280	ILE
1	B	291	GLN
1	B	297	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	HIS
1	A	119	HIS
1	A	127	HIS
1	A	129	HIS
1	A	178	ASN
1	A	258	ASN
1	A	279	GLN
1	A	291	GLN
1	A	300	GLN
1	B	35	HIS
1	B	107	HIS
1	B	119	HIS
1	B	127	HIS
1	B	168	GLN
1	B	187	GLN
1	B	279	GLN
1	B	291	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	310	-	4,4,4	0.46	0	6,6,6	0.27	0
4	R2G	A	311	-	7,8,8	1.41	1 (14%)	5,10,10	5.10	4 (80%)
2	URF	A	2001	-	9,9,9	2.66	3 (33%)	12,12,12	3.88	8 (66%)
2	URF	B	2011	-	9,9,9	2.43	4 (44%)	12,12,12	4.18	9 (75%)
3	SO4	A	310	-	4,4,4	0.89	0	6,6,6	0.83	0
4	R2G	B	311	-	7,8,8	1.46	1 (14%)	5,10,10	4.24	4 (80%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	URF	A	2001	-	-	-	0/1/1/1
2	URF	B	2011	-	-	-	0/1/1/1
4	R2G	A	311	-	-	2/2/12/12	0/1/1/1
4	R2G	B	311	-	-	2/2/12/12	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	URF	C6-C5	6.63	1.39	1.33
2	B	2011	URF	C4-C5	4.38	1.49	1.44
2	B	2011	URF	C6-C5	3.16	1.36	1.33
2	B	2011	URF	C4-N3	-2.87	1.33	1.38
4	A	311	R2G	O3-C3	-2.80	1.38	1.43
2	A	2001	URF	C4-N3	-2.79	1.33	1.38
2	B	2011	URF	C2-N1	-2.77	1.33	1.36
4	B	311	R2G	C1-C2	2.56	1.36	1.31
2	A	2001	URF	C2-N1	-2.26	1.33	1.36

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	311	R2G	C5-C4-C3	7.96	133.88	115.10
2	A	2001	URF	O4-C4-C5	-7.87	118.99	125.69
4	B	311	R2G	C5-C4-C3	7.11	131.89	115.10
4	A	311	R2G	O4-C1-C2	-7.09	106.88	114.79
2	B	2011	URF	N1-C2-N3	6.66	122.19	115.17
2	A	2001	URF	N1-C2-N3	6.37	121.89	115.17
2	B	2011	URF	F5-C5-C4	6.36	122.05	116.47
2	B	2011	URF	C5-C4-N3	5.83	117.97	112.64
2	B	2011	URF	O2-C2-N1	-4.97	117.66	122.79
2	B	2011	URF	C4-N3-C2	-4.61	120.02	126.37
2	A	2001	URF	F5-C5-C6	4.10	124.59	121.58
4	B	311	R2G	O4-C1-C2	-4.06	110.25	114.79
4	B	311	R2G	O4-C4-C5	4.04	113.85	108.37
2	B	2011	URF	F5-C5-C6	-3.74	118.83	121.58
2	B	2011	URF	O4-C4-C5	-3.68	122.56	125.69
2	A	2001	URF	C5-C4-N3	3.62	115.95	112.64
2	A	2001	URF	O2-C2-N1	-3.56	119.11	122.79
2	B	2011	URF	C6-C5-C4	-3.49	119.11	122.11
2	A	2001	URF	C4-N3-C2	-3.34	121.77	126.37
4	A	311	R2G	C3-C2-C1	-3.13	100.15	106.88
2	A	2001	URF	F5-C5-C4	-2.73	114.08	116.47
2	A	2001	URF	O4-C4-N3	2.63	125.05	120.11
4	B	311	R2G	C3-C2-C1	-2.56	101.39	106.88
4	A	311	R2G	O4-C4-C5	-2.40	105.12	108.37
2	B	2011	URF	C5-C6-N1	-2.12	117.71	119.49

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	311	R2G	C3-C4-C5-O5
4	B	311	R2G	C3-C4-C5-O5
4	B	311	R2G	O4-C4-C5-O5
4	A	311	R2G	O4-C4-C5-O5

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	311	R2G	4	0
2	A	2001	URF	2	0
2	B	2011	URF	1	0
4	B	311	R2G	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/309 (94%)	-0.38	13 (4%) 39 33	16, 30, 57, 93	0
1	B	293/309 (94%)	-0.46	8 (2%) 56 50	18, 31, 54, 89	0
All	All	586/618 (94%)	-0.42	21 (3%) 46 40	16, 31, 57, 93	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	233	ASP	5.2
1	A	80	PRO	4.4
1	B	79	HIS	4.2
1	B	309	ALA	4.1
1	A	17	LEU	4.1
1	A	309	ALA	4.1
1	B	14	HIS	3.7
1	B	285	ASP	3.5
1	A	79	HIS	3.1
1	A	233	ASP	2.9
1	A	154	THR	2.7
1	B	78	ALA	2.7
1	B	307	MET	2.6
1	B	220	ASP	2.4
1	A	81	GLY	2.4
1	A	82	ALA	2.4
1	A	73	MET	2.3
1	A	307	MET	2.3
1	A	77	PHE	2.2
1	A	78	ALA	2.1
1	A	285	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	R2G	A	311	8/8	0.88	0.17	40,45,47,49	0
4	R2G	B	311	8/8	0.92	0.14	39,41,44,45	0
2	URF	B	2011	9/9	0.96	0.09	20,23,27,27	0
2	URF	A	2001	9/9	0.97	0.12	24,24,27,29	0
3	SO4	A	310	5/5	0.98	0.10	27,30,36,37	0
3	SO4	B	310	5/5	0.98	0.09	31,34,36,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.