



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 11:22 PM EDT

PDB ID : 1KWX / pdb_00001kwx
Title : Rat mannose protein A complexed with b-Me-Fuc.
Authors : Ng, K.K.; Kolatkar, A.R.; Park-Snyder, S.; Feinberg, H.; Clark, D.A.; Drick-amer, K.; Weis, W.I.
Deposited on : 2002-01-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	71% 26% .
1	B	149	77% 23% .
1	C	149	3% 80% 20%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MANNOSE-BINDING PROTEIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	0	3	0
			1157	722	198	228	9			
1	B	149	Total	C	N	O	S	0	4	0
			1158	722	198	229	9			
1	C	149	Total	C	N	O	S	0	2	0
			1159	724	200	227	8			

- Molecule 2 is methyl beta-L-fucopyranoside (CCD ID: MFB) (formula: $C_7H_{14}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		
2	C	1	Total	C	O	0	0
			11	6	5		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total 4	Ca 4	0	0
3	B	3	Total 3	Ca 3	0	0
3	C	3	Total 3	Ca 3	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Cl 1	0	0
4	B	1	Total 1	Cl 1	0	0
4	C	1	Total 1	Cl 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	156	Total 156	O 156	0	0
5	B	129	Total 129	O 129	0	0
5	C	128	Total 128	O 128	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: MANNOSE-BINDING PROTEIN A



• Molecule 1: MANNOSE-BINDING PROTEIN A



• Molecule 1: MANNOSE-BINDING PROTEIN A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	78.48Å 85.43Å 97.80Å 90.00° 107.37° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.00) 96.5 (30.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.03 (at 2.00Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.200 , 0.229 0.188 , 0.214	Depositor DCC
R_{free} test set	4169 reflections (10.19%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3933	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MFB, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	1/1193 (0.1%)	1.02	5/1605 (0.3%)
1	B	0.48	0/1199	0.92	2/1613 (0.1%)
1	C	0.47	0/1190	1.04	10/1602 (0.6%)
All	All	0.51	1/3582 (0.0%)	0.99	17/4820 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	167	GLN	CD-OE1	-6.38	1.11	1.23

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	PHE	CA-C-N	6.73	126.75	119.89
1	A	219	PHE	C-N-CA	6.73	126.75	119.89
1	C	219	PHE	CA-C-N	6.07	126.03	119.78
1	C	219	PHE	C-N-CA	6.07	126.03	119.78
1	C	198	ILE	N-CA-C	-6.03	99.89	108.58
1	C	182	LYS	N-CA-C	-5.98	102.63	110.53
1	A	198	ILE	N-CA-C	-5.72	100.34	108.58
1	B	189	HIS	N-CA-C	5.62	117.54	110.24
1	C	115	ASN	N-CA-C	-5.62	106.01	112.92
1	A	115	ASN	N-CA-C	-5.56	105.71	112.88
1	A	182	LYS	N-CA-C	-5.47	103.30	110.53
1	C	202	GLY	N-CA-C	-5.46	108.11	115.21
1	C	189	HIS	N-CA-C	5.31	117.93	110.23
1	C	201	ASN	N-CA-C	-5.29	106.76	114.12
1	B	198	ILE	N-CA-C	-5.13	100.39	108.23
1	C	176	LEU	N-CA-C	5.09	117.69	110.10
1	C	174	GLY	N-CA-C	5.02	117.92	112.29

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1157	0	1136	37	0
1	B	1158	0	1137	26	0
1	C	1159	0	1137	21	0
2	A	11	0	9	0	0
2	B	11	0	9	0	0
2	C	11	0	10	0	0
3	A	4	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	156	0	0	2	0
5	B	129	0	0	3	0
5	C	128	0	0	1	0
All	All	3933	0	3438	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:169[B]:MET:HE1	1:B:175:ARG:HG2	1.61	0.79
1:C:105:LYS:HD3	1:C:111:PHE:CD2	2.18	0.78
1:A:145:LYS:O	1:A:149:GLU:HG2	1.87	0.75
1:B:138:PRO:HG3	1:B:147:ILE:HD12	1.70	0.71
1:A:81:MET:SD	1:B:81:MET:HE2	2.36	0.66
1:B:88:LEU:HD21	1:C:89:LYS:HG2	1.77	0.65
1:A:110:LYS:HB2	1:A:219:PHE:O	1.97	0.65
1:A:78:LEU:HD21	1:C:78:LEU:HD13	1.79	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:LEU:HA	1:C:94:LEU:HD11	1.81	0.61
1:C:99[A]:HIS:CE1	1:C:103:MET:SD	2.94	0.61
1:A:78:LEU:HD21	1:C:78:LEU:CD1	2.31	0.61
1:A:138:PRO:HG3	1:A:147:ILE:HD12	1.80	0.60
1:A:132:ARG:HD3	1:C:94:LEU:CD2	2.31	0.60
1:B:169[B]:MET:SD	1:B:175:ARG:HA	2.41	0.60
1:B:99:HIS:HA	1:C:99[A]:HIS:CE1	2.38	0.58
1:B:151:ALA:O	1:B:152:LYS:HB2	2.06	0.56
5:B:613:HOH:O	1:C:89:LYS:HE2	2.05	0.56
1:A:134:THR:HG22	1:A:135:VAL:N	2.20	0.56
1:A:149:GLU:O	1:A:152:LYS:HE3	2.07	0.55
1:C:179:SER:HA	1:C:204:TRP:CH2	2.42	0.55
1:A:161:ASP:OD1	1:A:194:ASP:HA	2.07	0.55
1:A:132:ARG:HD3	1:C:94:LEU:HD21	1.89	0.55
1:A:151:ALA:O	1:A:152:LYS:HB2	2.09	0.53
1:C:161:ASP:OD1	1:C:194:ASP:HA	2.09	0.52
1:C:100:ALA:HA	1:C:103:MET:HE3	1.91	0.51
1:B:145:LYS:O	1:B:149:GLU:HG2	2.11	0.51
1:B:169[B]:MET:HE3	5:B:656:HOH:O	2.11	0.50
1:A:127:LEU:C	1:A:127:LEU:HD23	2.37	0.50
1:A:110:LYS:HE2	1:A:143:GLU:CD	2.38	0.49
1:A:77:LYS:HB2	1:B:78:LEU:HD21	1.95	0.49
1:B:161:ASP:OD1	1:B:194:ASP:HA	2.13	0.49
1:A:88:LEU:HD21	1:B:89:LYS:HG2	1.96	0.48
1:B:120:PRO:HD2	1:B:123:LYS:HD2	1.95	0.48
1:B:145:LYS:HB2	1:B:145:LYS:NZ	2.28	0.48
1:C:111:PHE:HB2	5:C:757:HOH:O	2.14	0.48
1:A:207:ILE:HG13	1:A:208:SER:H	1.79	0.47
1:B:138:PRO:HG3	1:B:147:ILE:CD1	2.43	0.47
1:B:176:LEU:HB3	5:B:708:HOH:O	2.13	0.47
1:A:74:ILE:CG2	1:A:75:GLU:N	2.78	0.47
1:A:78:LEU:HD22	1:C:81:MET:CE	2.45	0.46
1:B:78:LEU:O	1:B:82:GLU:HG3	2.15	0.46
1:A:120:PRO:HD2	1:A:123:LYS:HG3	1.99	0.45
1:A:169[B]:MET:HE3	5:A:738:HOH:O	2.15	0.45
1:C:151:ALA:O	1:C:152:LYS:HB2	2.15	0.44
1:A:81:MET:HG3	1:B:85:ILE:CD1	2.48	0.44
1:B:106:LYS:HD2	1:C:117:GLU:HB2	1.98	0.44
1:C:207:ILE:HG13	1:C:208:SER:H	1.82	0.44
1:B:207:ILE:HG13	1:B:208:SER:H	1.82	0.44
1:A:200:ASP:O	1:A:201:ASN:HB2	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:LEU:HD13	1:B:132:ARG:HG3	2.01	0.43
1:B:183:LYS:O	1:B:184:ASP:HB2	2.19	0.43
1:A:138:PRO:HG3	1:A:147:ILE:CD1	2.47	0.42
1:A:199:VAL:HG23	1:A:200:ASP:N	2.34	0.42
1:A:169[B]:MET:CE	5:A:738:HOH:O	2.67	0.42
1:A:187:ASN:O	1:A:188:ASP:C	2.62	0.42
1:B:181:TRP:CD2	1:B:186:PRO:HD3	2.55	0.42
1:A:74:ILE:HG23	1:A:75:GLU:N	2.35	0.42
1:A:78:LEU:HD22	1:C:81:MET:HE1	2.02	0.42
1:A:182:LYS:HE2	1:A:182:LYS:HB3	1.85	0.42
1:A:181:TRP:CD2	1:A:186:PRO:HD3	2.55	0.42
1:B:151:ALA:C	1:B:153:THR:H	2.28	0.41
1:B:169[B]:MET:CE	1:B:175:ARG:HG2	2.41	0.41
1:A:183:LYS:HG2	1:A:184:ASP:OD1	2.20	0.41
1:A:207:ILE:HG13	1:A:208:SER:N	2.35	0.41
1:A:78:LEU:O	1:A:82:GLU:HG3	2.20	0.41
1:C:132:ARG:HA	1:C:132:ARG:HD3	1.86	0.41
1:B:179:SER:HA	1:B:204:TRP:CH2	2.56	0.41
1:C:207:ILE:HG13	1:C:208:SER:N	2.35	0.40
1:A:168:PHE:C	1:A:169[B]:MET:HG2	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/149 (101%)	146 (97%)	3 (2%)	1 (1%)	18	14
1	B	151/149 (101%)	146 (97%)	4 (3%)	1 (1%)	18	14
1	C	149/149 (100%)	144 (97%)	3 (2%)	2 (1%)	9	5
All	All	450/447 (101%)	436 (97%)	10 (2%)	4 (1%)	14	9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	200	ASP
1	A	199	VAL
1	B	199	VAL
1	C	74	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/126 (102%)	128 (99%)	1 (1%)	73	80
1	B	130/126 (103%)	128 (98%)	2 (2%)	57	64
1	C	128/126 (102%)	128 (100%)	0	100	100
All	All	387/378 (102%)	384 (99%)	3 (1%)	73	80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	167	GLN
1	B	94	LEU
1	B	145	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	ASN
1	A	86	ASN
1	A	96	ASN
1	A	148	GLN
1	A	167	GLN
1	A	210	GLN
1	B	80	ASN
1	B	86	ASN
1	B	96	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	148	GLN
1	B	180	ASN
1	C	80	ASN
1	C	96	ASN
1	C	148	GLN
1	C	189	HIS
1	C	205	ASN
1	C	210	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 13 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MFB	A	501	3	11,11,12	0.40	0	16,16,17	0.27	0
2	MFB	B	601	3	11,11,12	0.45	0	16,16,17	0.42	0
2	MFB	C	701	3	11,11,12	0.44	0	16,16,17	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MFB	A	501	3	-	-	0/1/1/1
2	MFB	B	601	3	-	-	0/1/1/1
2	MFB	C	701	3	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	149/149 (100%)	-0.28	2 (1%) 75 74	12, 20, 35, 44	3 (2%)
1	B	149/149 (100%)	-0.09	2 (1%) 75 74	15, 24, 40, 50	4 (2%)
1	C	149/149 (100%)	-0.03	5 (3%) 48 47	11, 24, 40, 56	2 (1%)
All	All	447/447 (100%)	-0.13	9 (2%) 65 64	11, 23, 40, 56	9 (2%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	199	VAL	3.3
1	A	200	ASP	3.3
1	C	200	ASP	2.6
1	A	201	ASN	2.3
1	B	183	LYS	2.2
1	C	108	GLY	2.2
1	C	169	MET	2.1
1	C	201	ASN	2.1
1	B	182	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	A	502	1/1	0.43	0.21	16,16,16,16	0
3	CA	A	504	1/1	0.63	0.15	17,17,17,17	0
3	CA	C	702	1/1	0.86	0.08	18,18,18,18	0
2	MFB	B	601	11/12	0.87	0.10	32,36,37,37	0
3	CA	C	703	1/1	0.89	0.07	19,19,19,19	0
3	CA	B	603	1/1	0.92	0.06	29,29,29,29	0
3	CA	A	606	1/1	0.93	0.07	30,30,30,30	1
3	CA	A	503	1/1	0.94	0.05	20,20,20,20	0
2	MFB	C	701	11/12	0.94	0.07	20,23,25,27	0
2	MFB	A	501	11/12	0.94	0.08	22,27,29,32	0
3	CA	C	704	1/1	0.95	0.05	19,19,19,19	0
4	CL	B	605	1/1	0.96	0.08	37,37,37,37	0
3	CA	B	602	1/1	0.99	0.03	20,20,20,20	0
4	CL	A	505	1/1	0.99	0.09	20,20,20,20	0
3	CA	B	604	1/1	0.99	0.02	22,22,22,22	0
4	CL	C	705	1/1	0.99	0.06	23,23,23,23	0

6.5 Other polymers

There are no such residues in this entry.