



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:40 AM UTC

PDB ID : 4KWD / pdb_00004kwd
Title : Crystal structure of Aspergillus terreus aristolochene synthase complexed with (1R,8R,9aS)-1,9a-dimethyl-8-(prop-1-en-2-yl)decahydroquinolizin-5-ium
Authors : Chen, M.; Al-lami, N.; Janvier, M.; D'Antonio, E.L.; Faraldos, J.A.; Cane, D.E.; Allemann, R.K.; Christianson, D.W.
Deposited on : 2013-05-23
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	88% 8% ..
1	B	314	90% 6% .
1	C	314	2% 84% 12% ..
1	D	314	35% 80% 14% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11000 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aristolochene synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	1	0
			2459	1570	415	458	16			
1	B	304	Total	C	N	O	S	0	1	0
			2459	1570	415	458	16			
1	C	304	Total	C	N	O	S	0	1	0
			2459	1570	415	458	16			
1	D	304	Total	C	N	O	S	0	0	0
			2454	1566	415	458	15			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q9UR08
A	2	HIS	-	expression tag	UNP Q9UR08
A	3	HIS	-	expression tag	UNP Q9UR08
A	4	HIS	-	expression tag	UNP Q9UR08
A	5	HIS	-	expression tag	UNP Q9UR08
A	6	HIS	-	expression tag	UNP Q9UR08
A	7	HIS	-	expression tag	UNP Q9UR08
B	1	MET	-	expression tag	UNP Q9UR08
B	2	HIS	-	expression tag	UNP Q9UR08
B	3	HIS	-	expression tag	UNP Q9UR08
B	4	HIS	-	expression tag	UNP Q9UR08
B	5	HIS	-	expression tag	UNP Q9UR08
B	6	HIS	-	expression tag	UNP Q9UR08
B	7	HIS	-	expression tag	UNP Q9UR08
C	1	MET	-	expression tag	UNP Q9UR08
C	2	HIS	-	expression tag	UNP Q9UR08
C	3	HIS	-	expression tag	UNP Q9UR08
C	4	HIS	-	expression tag	UNP Q9UR08
C	5	HIS	-	expression tag	UNP Q9UR08
C	6	HIS	-	expression tag	UNP Q9UR08
C	7	HIS	-	expression tag	UNP Q9UR08

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	MET	-	expression tag	UNP Q9UR08
D	2	HIS	-	expression tag	UNP Q9UR08
D	3	HIS	-	expression tag	UNP Q9UR08
D	4	HIS	-	expression tag	UNP Q9UR08
D	5	HIS	-	expression tag	UNP Q9UR08
D	6	HIS	-	expression tag	UNP Q9UR08
D	7	HIS	-	expression tag	UNP Q9UR08

- Molecule 2 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 9 7 2	0	0
2	B	1	Total O P 9 7 2	0	0
2	C	1	Total O P 9 7 2	0	0
2	D	1	Total O P 9 7 2	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Mg 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Mg	0	0
			3	3		
3	C	3	Total	Mg	0	0
			3	3		
3	D	3	Total	Mg	0	0
			3	3		

- Molecule 4 is (1R,5R,8R,9aS)-1,9a-dimethyl-8-(prop-1-en-2-yl)octahydro-2H-quinolizinium (CCD ID: JF2) (formula: C₁₄H₂₆N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			15	14	1		
4	B	1	Total	C	N	0	0
			15	14	1		
4	C	1	Total	C	N	0	0
			15	14	1		
4	D	1	Total	C	N	0	0
			15	14	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	336	Total	O	0	0
			336	336		
6	B	353	Total	O	0	0
			353	353		
6	C	229	Total	O	0	0
			229	229		
6	D	131	Total	O	0	0
			131	131		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aristolochene synthase




- Molecule 1: Aristolochene synthase




- Molecule 1: Aristolochene synthase



- Molecule 1: Aristolochene synthase





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	123.86Å 123.86Å 202.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.37 – 1.86 47.37 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.37-1.86) 99.8 (47.37-1.86)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 1.86Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1370)	Depositor
R, R_{free}	0.193 , 0.226 0.195 , 0.226	Depositor DCC
R_{free} test set	7601 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.252	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.012 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11000	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, JF2, GOL, POP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	0/2517	0.94	2/3406 (0.1%)
1	B	0.93	0/2517	0.95	2/3406 (0.1%)
1	C	0.82	0/2517	0.91	3/3406 (0.1%)
1	D	0.69	0/2509	0.94	2/3396 (0.1%)
All	All	0.84	0/10060	0.93	9/13614 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	57	VAL	CB-CA-C	-8.06	99.20	112.26
1	B	10	PRO	CA-C-N	-6.54	113.64	120.38
1	B	10	PRO	C-N-CA	-6.54	113.64	120.38
1	A	9	GLU	CA-C-N	-6.06	114.14	120.38
1	A	9	GLU	C-N-CA	-6.06	114.14	120.38
1	C	40	PHE	CA-C-N	5.62	125.23	119.56
1	C	40	PHE	C-N-CA	5.62	125.23	119.56
1	C	107	ARG	NE-CZ-NH1	-5.55	115.95	121.50
1	D	11	PRO	O-C-N	5.13	123.56	121.15

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2459	0	2434	11	0
1	B	2459	0	2434	7	0
1	C	2459	0	2434	15	0
1	D	2454	0	2425	36	0
2	A	9	0	0	0	0
2	B	9	0	0	0	0
2	C	9	0	0	1	0
2	D	9	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
4	A	15	0	26	0	0
4	B	15	0	26	1	0
4	C	15	0	26	1	0
4	D	15	0	26	0	0
5	B	6	0	8	0	0
5	C	6	0	8	1	0
6	A	336	0	0	3	0
6	B	353	0	0	5	0
6	C	229	0	0	2	0
6	D	131	0	0	5	0
All	All	11000	0	9847	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:135:GLU:HG2	1:D:189:LYS:HE2	1.72	0.72
1:D:165:TYR:CZ	1:D:169:ARG:HG3	2.31	0.66
1:D:36:GLN:HG2	1:D:37:HIS:CD2	2.31	0.65
4:B:705:JF2:H6	6:B:1153:HOH:O	1.96	0.64
1:D:201:ILE:O	6:D:570:HOH:O	2.14	0.64
1:A:197:ARG:HD3	1:A:280:LEU:HB3	1.82	0.62
1:A:35:LEU:O	1:A:46:ARG:NH2	2.32	0.61
1:D:15:PHE:HE2	1:D:296:MET:HE2	1.66	0.59
1:B:154:ARG:HD2	6:B:949:HOH:O	2.01	0.59
1:C:18:LEU:HD22	1:C:63:PRO:HB2	1.85	0.57
1:C:160:MET:HE1	1:C:165:TYR:HD2	1.72	0.55
1:B:236:CYS:O	6:B:1126:HOH:O	2.17	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:64:LYS:O	1:D:186:MET:HG2	2.06	0.55
1:C:35:LEU:O	1:C:46:ARG:NH2	2.39	0.54
1:A:222:LEU:HD22	1:A:226:LYS:HE3	1.90	0.54
1:D:57:VAL:HG22	1:D:302:TRP:CE2	2.43	0.54
1:D:69:ARG:NH2	1:D:133:ASP:OD1	2.33	0.53
1:D:112:PRO:HD3	1:D:123:TYR:CD1	2.44	0.53
1:B:183:ARG:NH2	6:B:1116:HOH:O	2.41	0.52
1:C:196:GLN:HA	1:C:199:ARG:HD3	1.91	0.52
1:B:200:GLU:OE1	1:B:267:ARG:NH2	2.43	0.51
1:C:183:ARG:NH1	6:C:568:HOH:O	2.44	0.51
1:D:15:PHE:CE2	1:D:296:MET:HE2	2.45	0.51
1:D:57:VAL:HG13	1:D:302:TRP:CG	2.46	0.51
1:A:267:ARG:NH1	6:A:670:HOH:O	2.42	0.50
1:A:25:GLU:HB2	6:A:614:HOH:O	2.11	0.50
1:A:219:GLU:HG2	6:A:653:HOH:O	2.12	0.50
1:C:260[A]:MET:HE2	1:D:256:VAL:HG11	1.93	0.50
1:D:160:MET:HE1	1:D:165:TYR:HD2	1.76	0.49
1:D:307:LEU:O	1:D:311:VAL:HG22	2.12	0.49
1:D:75:ARG:HB2	1:D:125:LEU:HD21	1.95	0.48
1:A:160:MET:HE3	1:A:160:MET:HB2	1.68	0.47
1:D:15:PHE:HE2	1:D:296:MET:CE	2.27	0.47
1:A:57:VAL:HB	1:A:302:TRP:CE2	2.50	0.47
1:B:15:PHE:HB3	1:B:297:SER:HB3	1.97	0.47
1:D:46:ARG:HD2	6:D:541:HOH:O	2.15	0.46
1:D:66:LEU:HB2	1:D:69:ARG:HD3	1.98	0.46
1:C:213:ASN:O	1:C:217:SER:OG	2.34	0.46
1:D:38:TRP:HB3	1:D:40:PHE:CE1	2.51	0.45
1:D:93:GLU:HG3	1:D:154:ARG:HH22	1.81	0.45
1:D:43:GLU:HG3	1:D:47:LYS:HE2	1.99	0.45
1:D:175:LYS:HG3	6:D:551:HOH:O	2.17	0.45
1:D:62:PHE:CE1	1:D:181:LEU:HD13	2.52	0.45
1:C:15:PHE:HB3	1:C:297:SER:HB3	1.98	0.45
1:D:222:LEU:HD22	1:D:226:LYS:HE3	2.00	0.44
1:D:266:LEU:HD13	6:D:510:HOH:O	2.17	0.44
1:D:160:MET:HE3	1:D:165:TYR:HB2	2.00	0.43
1:A:57:VAL:HB	1:A:302:TRP:CD2	2.53	0.43
1:C:149:ARG:NH1	5:C:406:GOL:H11	2.33	0.43
1:A:153:ASP:CG	1:A:155:THR:HG22	2.43	0.43
1:C:28:LYS:HA	1:C:28:LYS:HD3	1.81	0.43
1:C:46:ARG:HD2	6:C:619:HOH:O	2.18	0.43
1:D:165:TYR:CE2	1:D:169:ARG:HG3	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:169:ARG:HG2	1:C:210:SER:HB3	2.00	0.43
2:C:401:POP:O	4:C:405:JF2:H17	2.18	0.42
1:B:236:CYS:HA	6:B:1023:HOH:O	2.18	0.42
1:C:40:PHE:HA	1:C:41:PRO:HD3	1.76	0.42
1:D:208:HIS:CG	1:D:296:MET:HE3	2.54	0.42
1:D:106:SER:HA	1:D:126:TRP:CD1	2.55	0.42
1:B:57:VAL:HB	1:B:302:TRP:CD2	2.54	0.42
1:D:184:PHE:CD2	1:D:184:PHE:C	2.98	0.41
1:D:194:GLU:OE1	1:D:282:THR:OG1	2.27	0.41
1:D:145:PHE:O	1:D:149:ARG:HD2	2.20	0.41
1:A:177:LEU:HD12	1:A:177:LEU:C	2.46	0.41
1:C:261:CYS:O	1:C:265:GLU:HG3	2.21	0.41
1:C:99:ASN:HB3	1:C:148:MET:HE2	2.03	0.41
1:D:57:VAL:HG22	1:D:302:TRP:NE1	2.36	0.41
1:D:303:SER:OG	6:D:555:HOH:O	2.22	0.40
1:D:165:TYR:OH	1:D:169:ARG:HD2	2.21	0.40
1:D:160:MET:CE	1:D:165:TYR:HD2	2.33	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/314 (96%)	300 (99%)	3 (1%)	0	100	100
1	B	303/314 (96%)	300 (99%)	3 (1%)	0	100	100
1	C	303/314 (96%)	299 (99%)	4 (1%)	0	100	100
1	D	302/314 (96%)	293 (97%)	8 (3%)	1 (0%)	36	25
All	All	1211/1256 (96%)	1192 (98%)	18 (2%)	1 (0%)	48	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	133	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/276 (97%)	253 (95%)	14 (5%)	21	7
1	B	267/276 (97%)	257 (96%)	10 (4%)	30	15
1	C	267/276 (97%)	252 (94%)	15 (6%)	19	6
1	D	266/276 (96%)	250 (94%)	16 (6%)	17	5
All	All	1067/1104 (97%)	1012 (95%)	55 (5%)	21	7

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	LEU
1	A	25	GLU
1	A	125	LEU
1	A	134	ARG
1	A	167	GLU
1	A	173	VAL
1	A	177	LEU
1	A	181	LEU
1	A	222	LEU
1	A	237	THR
1	A	266	LEU
1	A	285	LEU
1	A	301	LEU
1	A	307	LEU
1	B	111	LEU
1	B	181	LEU
1	B	222	LEU
1	B	237	THR
1	B	240	GLN
1	B	258	PHE
1	B	266	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	285	LEU
1	B	301	LEU
1	B	311	VAL
1	C	28	LYS
1	C	111	LEU
1	C	125	LEU
1	C	181	LEU
1	C	222	LEU
1	C	237	THR
1	C	240	GLN
1	C	247	ASP
1	C	251	GLU
1	C	266	LEU
1	C	267	ARG
1	C	285	LEU
1	C	301	LEU
1	C	307	LEU
1	C	311	VAL
1	D	36	GLN
1	D	57	VAL
1	D	93	GLU
1	D	125	LEU
1	D	162	LEU
1	D	167	GLU
1	D	169	ARG
1	D	177	LEU
1	D	181	LEU
1	D	185	SER
1	D	222	LEU
1	D	258	PHE
1	D	266	LEU
1	D	275	LEU
1	D	301	LEU
1	D	311	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	196	GLN
1	B	204	ASN
1	B	240	GLN
1	B	299	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	39	ASN
1	C	204	ASN
1	D	132	HIS
1	D	240	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 12 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	POP	B	704	3	6,8,8	1.01	0	12,13,13	0.85	0
2	POP	A	401	3	6,8,8	0.87	0	12,13,13	0.76	0
4	JF2	C	405	-	14,16,16	0.78	1 (7%)	19,24,24	1.36	3 (15%)
2	POP	C	401	3	6,8,8	0.93	0	12,13,13	1.21	1 (8%)
4	JF2	B	705	-	14,16,16	0.62	0	19,24,24	1.88	5 (26%)
4	JF2	D	405	-	14,16,16	0.74	1 (7%)	19,24,24	1.19	1 (5%)
5	GOL	B	706	-	5,5,5	0.26	0	5,5,5	0.80	0
2	POP	D	401	3	6,8,8	0.50	0	12,13,13	1.59	2 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	JF2	A	405	-	14,16,16	0.86	1 (7%)	19,24,24	1.98	4 (21%)
5	GOL	C	406	-	5,5,5	0.21	0	5,5,5	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	POP	B	704	3	-	0/6/6/6	-
2	POP	A	401	3	-	0/6/6/6	-
4	JF2	C	405	-	-	0/4/32/32	0/2/2/2
2	POP	C	401	3	-	0/6/6/6	-
4	JF2	B	705	-	-	0/4/32/32	0/2/2/2
4	JF2	D	405	-	-	0/4/32/32	0/2/2/2
5	GOL	B	706	-	-	0/4/4/4	-
2	POP	D	401	3	-	0/6/6/6	-
4	JF2	A	405	-	-	0/4/32/32	0/2/2/2
5	GOL	C	406	-	-	2/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	405	JF2	CAO-CAM	-2.27	1.52	1.55
4	A	405	JF2	CAH-NAN	2.23	1.50	1.47
4	C	405	JF2	CAO-CAM	-2.16	1.52	1.55

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	405	JF2	CAF-CAE-CAH	4.43	116.57	110.75
4	A	405	JF2	CAE-CAF-CAM	-4.35	106.11	112.68
4	B	705	JF2	CAI-NAN-CAH	-4.19	104.67	111.02
4	B	705	JF2	CAD-CAO-CAJ	-3.60	105.15	110.76
2	D	401	POP	O6-P2-O5	3.21	119.85	107.80
4	C	405	JF2	CAD-CAO-CAM	-3.16	107.34	112.16
4	A	405	JF2	CAJ-CAL-CAK	-2.87	108.06	112.61
4	B	705	JF2	CAJ-CAL-CAG	-2.70	106.57	109.72
4	C	405	JF2	CAE-CAF-CAM	-2.56	108.81	112.68
4	B	705	JF2	CAG-CAL-CAK	-2.55	106.99	113.27
2	D	401	POP	O3-P1-O2	2.30	116.42	107.80
4	D	405	JF2	CAI-NAN-CAH	-2.27	107.58	111.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	405	JF2	CAI-NAN-CAH	-2.13	107.79	111.02
4	C	405	JF2	CAI-NAN-CAH	-2.11	107.82	111.02
2	C	401	POP	O6-P2-O4	2.08	118.93	110.83
4	B	705	JF2	CAE-CAF-CAM	-2.07	109.55	112.68

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	406	GOL	C1-C2-C3-O3
5	C	406	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	405	JF2	1	0
2	C	401	POP	1	0
4	B	705	JF2	1	0
5	C	406	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/314 (96%)	-0.43	2 (0%) 84 87	8, 17, 32, 50	1 (0%)
1	B	304/314 (96%)	-0.55	0 100 100	8, 15, 27, 41	1 (0%)
1	C	304/314 (96%)	0.27	6 (1%) 65 68	13, 25, 44, 58	1 (0%)
1	D	304/314 (96%)	1.65	110 (36%) 1 0	21, 38, 66, 96	0
All	All	1216/1256 (96%)	0.24	118 (9%) 13 13	8, 23, 49, 96	3 (0%)

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	280	LEU	5.9
1	D	285	LEU	5.8
1	C	311	VAL	5.5
1	D	195	LEU	5.0
1	D	61	TYR	4.9
1	D	18	LEU	4.8
1	D	272	VAL	4.8
1	D	287	ALA	4.8
1	D	126	TRP	4.7
1	D	311	VAL	4.7
1	D	275	LEU	4.7
1	C	8	LEU	4.7
1	D	188	LEU	4.3
1	D	279	GLY	4.3
1	D	277	ALA	4.2
1	D	8	LEU	4.1
1	D	284	GLY	4.1
1	D	270	LEU	4.0
1	D	23	VAL	3.9
1	D	200	GLU	3.9
1	D	73	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	283	PRO	3.8
1	D	294	TYR	3.7
1	D	198	VAL	3.7
1	D	64	LYS	3.7
1	D	19	CYS	3.6
1	D	268	HIS	3.5
1	D	65	ALA	3.5
1	D	273	ALA	3.4
1	D	57	VAL	3.4
1	D	72	PHE	3.4
1	D	288	TYR	3.3
1	D	111	LEU	3.3
1	D	199	ARG	3.3
1	D	136	MET	3.3
1	D	131	ALA	3.2
1	D	286	ALA	3.2
1	D	70	ILE	3.2
1	D	181	LEU	3.2
1	D	186	MET	3.2
1	D	56	ARG	3.1
1	D	179	ALA	3.1
1	D	205	CYS	3.1
1	D	137	ALA	3.1
1	D	154	ARG	3.0
1	D	22	LEU	3.0
1	D	74	CYS	3.0
1	D	59	CYS	3.0
1	D	144	VAL	2.9
1	D	123	TYR	2.9
1	D	282	THR	2.9
1	D	15	PHE	2.8
1	D	62	PHE	2.8
1	D	66	LEU	2.8
1	D	184	PHE	2.8
1	D	276	SER	2.8
1	D	182	MET	2.8
1	D	143	PRO	2.7
1	D	197	ARG	2.7
1	D	108	GLY	2.7
1	D	292	LEU	2.6
1	D	26	VAL	2.6
1	D	187	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	191	SER	2.6
1	D	261	CYS	2.6
1	D	190	LEU	2.6
1	D	139	GLU	2.6
1	D	20	HIS	2.5
1	D	129	MET	2.5
1	D	271	LEU	2.5
1	A	311	VAL	2.5
1	D	14	THR	2.5
1	D	194	GLU	2.5
1	D	266	LEU	2.5
1	D	291	GLY	2.5
1	D	140	ILE	2.4
1	D	63	PRO	2.4
1	A	154	ARG	2.4
1	D	60	LEU	2.4
1	D	146	LEU	2.4
1	C	246	ALA	2.4
1	D	264	TRP	2.4
1	D	183	ARG	2.4
1	D	135	GLU	2.4
1	D	281	GLU	2.4
1	D	185	SER	2.4
1	D	203	ALA	2.3
1	D	174	GLY	2.3
1	D	27	SER	2.3
1	D	16	GLN	2.3
1	D	105	ILE	2.3
1	D	76	LEU	2.3
1	D	36	GLN	2.3
1	D	289	VAL	2.3
1	D	125	LEU	2.3
1	D	89	TYR	2.2
1	D	11	PRO	2.2
1	D	180	ALA	2.2
1	D	302	TRP	2.2
1	D	141	LEU	2.2
1	D	128	SER	2.2
1	D	163	GLY	2.2
1	D	71	HIS	2.2
1	D	301	LEU	2.2
1	D	132	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	278	GLU	2.1
1	C	28	LYS	2.1
1	D	274	ARG	2.1
1	D	54	PHE	2.1
1	D	258	PHE	2.1
1	D	193	SER	2.1
1	C	247	ASP	2.1
1	D	189	LYS	2.0
1	D	21	PRO	2.0
1	D	77	LEU	2.0
1	D	296	MET	2.0
1	C	89	TYR	2.0
1	D	299	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	JF2	D	405	15/15	0.85	0.21	36,45,53,54	0
5	GOL	C	406	6/6	0.85	0.17	24,40,46,49	0
4	JF2	A	405	15/15	0.90	0.11	18,21,26,28	0
4	JF2	B	705	15/15	0.90	0.10	18,22,25,27	0
3	MG	D	404	1/1	0.92	0.05	24,24,24,24	0
4	JF2	C	405	15/15	0.92	0.12	25,31,35,37	0
3	MG	D	403	1/1	0.93	0.05	24,24,24,24	0
5	GOL	B	706	6/6	0.95	0.07	25,26,29,33	0
2	POP	D	401	9/9	0.95	0.08	23,28,28,29	0
3	MG	C	404	1/1	0.96	0.04	22,22,22,22	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	POP	C	401	9/9	0.97	0.06	18,19,22,24	0
3	MG	C	402	1/1	0.97	0.04	19,19,19,19	0
3	MG	C	403	1/1	0.97	0.03	22,22,22,22	0
3	MG	D	402	1/1	0.98	0.03	23,23,23,23	0
3	MG	A	403	1/1	0.98	0.02	13,13,13,13	0
3	MG	A	402	1/1	0.98	0.03	14,14,14,14	0
2	POP	A	401	9/9	0.99	0.03	10,11,12,13	0
2	POP	B	704	9/9	0.99	0.02	8,10,10,13	0
3	MG	A	404	1/1	0.99	0.03	11,11,11,11	0
3	MG	B	702	1/1	0.99	0.02	10,10,10,10	0
3	MG	B	701	1/1	1.00	0.02	12,12,12,12	0
3	MG	B	703	1/1	1.00	0.02	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.