



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:00 AM UTC

PDB ID : 4KXA / pdb_00004kxa
Title : Crystal structure of human aminopeptidase A complexed with aspartate and calcium
Authors : Yang, Y.; Liu, C.; Lin, Y.Y.; Li, F.
Deposited on : 2013-05-24
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	888	 84% 13% ..
2	B	2	 100%
2	C	2	 100%
2	F	2	 100%
2	G	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	3	 33% 67%
3	E	3	 67% 33%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamyl aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	875	Total	C	N	O	S	0	2	0
			7159	4584	1192	1358	25			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	213	ARG	GLN	variant	UNP Q07075
A	218	ALA	VAL	variant	UNP Q07075
A	958	HIS	-	expression tag	UNP Q07075
A	959	HIS	-	expression tag	UNP Q07075
A	960	HIS	-	expression tag	UNP Q07075
A	961	HIS	-	expression tag	UNP Q07075
A	962	HIS	-	expression tag	UNP Q07075
A	963	HIS	-	expression tag	UNP Q07075

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			42	24	3	15			
3	E	3	Total	C	N	O	0	0	0
			42	24	3	15			


- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is ASPARTIC ACID (CCD ID: ASP) (formula: $C_4H_7NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			9	4	1	4		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	507	Total	O	0	0
			507	507		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamyl aminopeptidase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  33% 67%

MAG1
MAG2
MAG3

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  67% 33%

MAG1
MAG2
MAG3

4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	142.09Å 142.09Å 237.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.57 – 2.40 48.57 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.2 (48.57-2.40) 97.2 (48.57-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.152 , 0.224 0.156 , 0.224	Depositor DCC
R_{free} test set	2838 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	45.4	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 78.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7901	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/7344	0.78	4/9985 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174	GLU	N-CA-C	-5.52	101.47	109.59
1	A	614	ASN	N-CA-C	5.49	118.50	109.72
1	A	616	PHE	N-CA-C	5.28	117.20	108.13
1	A	355	GLY	N-CA-C	5.16	120.28	113.37

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7159	0	6949	69	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
2	F	28	0	25	0	0
2	G	28	0	25	0	0
3	D	42	0	37	2	0
3	E	42	0	37	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	28	0	26	0	0
7	A	9	0	3	1	0
8	A	507	0	0	10	0
All	All	7901	0	7152	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:3:NAG:H83	3:E:3:NAG:H3	1.54	0.88
1:A:175:GLU:O	1:A:176:GLU:HB3	1.77	0.84
1:A:229:LYS:HE3	8:A:1568:HOH:O	1.82	0.79
1:A:423:GLU:O	1:A:427:VAL:HG12	1.84	0.78
1:A:118:THR:HG22	1:A:191:GLU:HG2	1.64	0.78
1:A:501:GLY:HA3	1:A:523:LEU:HD23	1.65	0.77
3:E:3:NAG:H3	3:E:3:NAG:C8	2.16	0.75
1:A:736:ALA:O	1:A:745:ARG:NH1	2.18	0.73
1:A:248:THR:OG1	1:A:273:ARG:HG3	1.91	0.70
1:A:761:LEU:HD22	1:A:793:ASN:HD21	1.54	0.70
1:A:101:HIS:CD2	1:A:242:THR:HG23	2.30	0.66
1:A:361:ASN:HB2	1:A:364:LEU:O	1.96	0.65
1:A:735:ASP:OD1	1:A:781:ASN:HB2	1.96	0.65
1:A:195:TRP:HZ3	8:A:1569:HOH:O	1.80	0.64
1:A:244:THR:HG22	1:A:278:LYS:HA	1.81	0.62
1:A:674:LYS:HD3	3:D:1:NAG:H62	1.81	0.60
1:A:717:GLU:HG2	1:A:755:MET:O	2.03	0.59
1:A:923:LYS:HG3	8:A:1478:HOH:O	2.02	0.58
1:A:175:GLU:O	1:A:176:GLU:CB	2.51	0.57
1:A:158:ARG:HB2	1:A:174:GLU:HG2	1.87	0.57
1:A:119:VAL:O	1:A:189:THR:HA	2.06	0.56
1:A:273:ARG:HD3	8:A:1499:HOH:O	2.09	0.52
1:A:455:MET:HE3	1:A:566:ALA:HB1	1.92	0.52
1:A:703:ILE:HD12	1:A:713:TYR:CE2	2.46	0.51
1:A:440:MET:HE2	8:A:1480:HOH:O	2.11	0.50
1:A:488:MET:HE3	1:A:535:MET:HE1	1.92	0.50
1:A:215:LYS:HE2	8:A:1580:HOH:O	2.12	0.49
1:A:616:PHE:HE1	1:A:618:LYS:HG3	1.76	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:ARG:HE	1:A:177:LEU:HD21	1.78	0.49
1:A:390:VAL:O	1:A:394:GLU:HG2	2.13	0.48
1:A:596:PHE:C	1:A:596:PHE:CD1	2.93	0.47
1:A:937:GLU:HG2	8:A:1449:HOH:O	2.14	0.47
1:A:130:ARG:NH2	1:A:172:GLU:OE1	2.48	0.47
1:A:77:ILE:HD11	1:A:144:LEU:N	2.30	0.47
1:A:427:VAL:HG11	1:A:436:MET:HE2	1.97	0.46
1:A:713:TYR:HB3	1:A:714:PRO:HD3	1.97	0.46
1:A:131:TYR:CD2	1:A:172:GLU:HB3	2.50	0.46
1:A:105:HIS:HE1	1:A:248:THR:OG1	1.98	0.46
1:A:101:HIS:HD2	1:A:242:THR:HG23	1.75	0.46
1:A:493:ILE:O	1:A:494:LYS:HB2	2.16	0.46
1:A:914:LYS:HD3	1:A:915:TYR:CE2	2.51	0.46
1:A:689:PHE:HA	1:A:744:LEU:HD13	1.98	0.46
1:A:424:PHE:CG	1:A:440:MET:HG2	2.52	0.45
1:A:125:LEU:HD21	1:A:127:ALA:O	2.17	0.45
1:A:793:ASN:HB3	8:A:1492:HOH:O	2.16	0.45
1:A:402:ASN:N	1:A:402:ASN:HD22	2.15	0.44
1:A:356:THR:HG21	7:A:1019:ASP:CG	2.42	0.44
1:A:943:ARG:HH12	1:A:947:ARG:HH11	1.66	0.44
1:A:359:MET:O	1:A:365:ILE:HG23	2.18	0.44
1:A:397:HIS:HA	1:A:400:PHE:O	2.18	0.44
1:A:585:TRP:CZ2	1:A:592:SER:HB2	2.53	0.43
1:A:161:PHE:CZ	1:A:170:VAL:HB	2.53	0.43
1:A:908:MET:SD	1:A:932:VAL:HG21	2.59	0.43
1:A:616:PHE:CE1	1:A:618:LYS:HG3	2.53	0.43
1:A:103:ASP:HB2	1:A:120:SER:HB2	2.01	0.43
1:A:565:ARG:CZ	1:A:937:GLU:HG3	2.49	0.43
1:A:133:TRP:CB	1:A:170:VAL:HG22	2.49	0.42
1:A:202:GLY:O	1:A:220:THR:HA	2.18	0.42
1:A:100:VAL:O	1:A:242:THR:HG22	2.19	0.42
1:A:596:PHE:C	1:A:596:PHE:HD1	2.27	0.42
1:A:215:LYS:CE	8:A:1580:HOH:O	2.68	0.41
1:A:175:GLU:HG3	1:A:176:GLU:O	2.21	0.41
1:A:585:TRP:CD1	1:A:617:LEU:CD2	3.03	0.41
1:A:952:ASN:HB3	3:D:2:NAG:O4	2.20	0.41
1:A:863:GLY:HA2	1:A:866:MET:HB2	2.01	0.41
1:A:724:VAL:O	1:A:725:LYS:C	2.63	0.41
1:A:923:LYS:HB2	1:A:924:PRO:HD3	2.02	0.41
1:A:139:THR:OG1	1:A:229:LYS:O	2.39	0.41
1:A:784:LEU:O	1:A:788:ARG:HG3	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:923:LYS:HD2	1:A:923:LYS:HA	1.91	0.40
1:A:834:ARG:HD2	8:A:1595:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	873/888 (98%)	839 (96%)	29 (3%)	5 (1%)	21 32

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	175	GLU
1	A	552	VAL
1	A	176	GLU
1	A	494	LYS
1	A	224	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	787/797 (99%)	744 (94%)	43 (6%)	19 34

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ILE
1	A	84	GLU
1	A	132	LEU
1	A	136	LEU
1	A	177	LEU
1	A	190	MET
1	A	200	LEU
1	A	252	GLU
1	A	273	ARG
1	A	278	LYS
1	A	298	LYS
1	A	299	ARG
1	A	303	SER
1	A	325	ILE
1	A	356	THR
1	A	406	MET
1	A	427	VAL
1	A	462	VAL
1	A	493	ILE
1	A	506	LEU
1	A	508	LYS
1	A	524	GLU
1	A	527	SER
1	A	539	THR
1	A	575	LEU
1	A	590	ILE
1	A	598	ARG
1	A	604	ILE
1	A	605	THR
1	A	614	ASN
1	A	658	LEU
1	A	665	LEU
1	A	677	LEU
1	A	703	ILE
1	A	712	LEU
1	A	724	VAL
1	A	738	ASP
1	A	748	VAL
1	A	778	LEU
1	A	794	SER
1	A	844	LEU
1	A	903	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	937	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	HIS
1	A	154	GLN
1	A	312	GLN
1	A	385	GLN
1	A	402	ASN
1	A	415	ASN
1	A	429	HIS
1	A	439	GLN
1	A	497	ASN
1	A	510	GLN
1	A	570	GLN
1	A	646	ASN
1	A	793	ASN
1	A	808	GLN
1	A	843	ASN
1	A	874	ASN
1	A	907	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.53	0	17,19,21	0.97	0
2	NAG	B	2	2	14,14,15	0.48	0	17,19,21	0.79	0
2	NAG	C	1	1,2	14,14,15	0.54	0	17,19,21	1.56	3 (17%)
2	NAG	C	2	2	14,14,15	0.65	0	17,19,21	2.06	2 (11%)
3	NAG	D	1	3,1	14,14,15	0.57	0	17,19,21	1.46	4 (23%)
3	NAG	D	2	3	14,14,15	0.83	1 (7%)	17,19,21	1.92	4 (23%)
3	NAG	D	3	3	14,14,15	0.72	0	17,19,21	0.91	0
3	NAG	E	1	3,1	14,14,15	0.48	0	17,19,21	0.96	1 (5%)
3	NAG	E	2	3	14,14,15	0.60	0	17,19,21	1.85	5 (29%)
3	NAG	E	3	3	14,14,15	1.08	1 (7%)	17,19,21	2.44	6 (35%)
2	NAG	F	1	1,2	14,14,15	0.67	1 (7%)	17,19,21	2.11	4 (23%)
2	NAG	F	2	2	14,14,15	0.50	0	17,19,21	1.02	1 (5%)
2	NAG	G	1	1,2	14,14,15	0.53	0	17,19,21	0.75	0
2	NAG	G	2	2	14,14,15	0.46	0	17,19,21	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	C	2	2	-	3/6/23/26	0/1/1/1
3	NAG	D	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	NAG	D	3	3	-	2/6/23/26	0/1/1/1
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	4/6/23/26	0/1/1/1
3	NAG	E	3	3	-	5/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	3	NAG	O6-C6	2.90	1.54	1.42
3	D	2	NAG	C1-C2	2.20	1.55	1.52
2	F	1	NAG	C1-C2	2.04	1.55	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	3	NAG	C2-N2-C7	6.25	131.27	122.90
2	C	2	NAG	C1-O5-C5	5.68	119.80	112.19
2	F	1	NAG	C1-O5-C5	5.60	119.70	112.19
2	C	2	NAG	C2-N2-C7	4.92	129.50	122.90
3	E	3	NAG	C4-C3-C2	4.38	117.44	111.02
3	D	2	NAG	C3-C4-C5	4.28	117.99	110.23
3	E	2	NAG	O5-C1-C2	-3.96	105.16	111.29
3	D	2	NAG	C4-C3-C2	3.94	116.79	111.02
2	F	1	NAG	O4-C4-C3	3.90	119.57	110.38
3	D	2	NAG	C1-O5-C5	3.77	117.24	112.19
3	E	3	NAG	C8-C7-N2	3.52	121.96	116.12
2	C	1	NAG	O4-C4-C3	3.43	118.45	110.38
3	E	2	NAG	C2-N2-C7	3.41	127.47	122.90
2	C	1	NAG	C1-O5-C5	3.26	116.55	112.19
2	F	1	NAG	C4-C3-C2	-3.24	106.27	111.02
3	E	3	NAG	C1-O5-C5	3.17	116.44	112.19
3	E	2	NAG	C1-O5-C5	3.08	116.32	112.19
3	E	3	NAG	C3-C4-C5	2.77	115.25	110.23
2	F	1	NAG	O4-C4-C5	2.75	116.10	109.32
2	F	2	NAG	C1-O5-C5	2.68	115.77	112.19
3	E	3	NAG	O7-C7-C8	-2.66	117.33	122.05
3	D	1	NAG	O7-C7-C8	-2.57	117.49	122.05
3	E	2	NAG	C1-C2-N2	2.54	114.44	110.43
3	D	1	NAG	C8-C7-N2	2.50	120.27	116.12
3	D	1	NAG	C2-N2-C7	2.43	126.15	122.90
2	C	1	NAG	C8-C7-N2	2.37	120.05	116.12
3	D	2	NAG	C6-C5-C4	-2.22	107.57	113.02
3	E	1	NAG	C1-O5-C5	2.15	115.06	112.19
3	D	1	NAG	C1-C2-N2	-2.10	107.12	110.43
3	E	2	NAG	O5-C5-C4	2.01	115.72	110.83

There are no chirality outliers.

All (27) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	3	NAG	C3-C2-N2-C7
2	C	2	NAG	O5-C5-C6-O6
3	D	3	NAG	O5-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
3	D	3	NAG	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	F	2	NAG	C8-C7-N2-C2
2	F	2	NAG	O7-C7-N2-C2
3	D	1	NAG	C8-C7-N2-C2
3	D	1	NAG	O7-C7-N2-C2
3	E	3	NAG	C8-C7-N2-C2
3	E	3	NAG	O7-C7-N2-C2
3	E	2	NAG	O5-C5-C6-O6
3	E	3	NAG	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
3	E	3	NAG	O5-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
3	E	2	NAG	C3-C2-N2-C7
3	E	2	NAG	C1-C2-N2-C7
2	C	2	NAG	C3-C2-N2-C7
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	2	NAG	1	0
3	E	3	NAG	2	0
3	D	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	ASP	A	1019	-	7,8,8	1.14	1 (14%)	6,10,10	1.55	1 (16%)
6	NAG	A	1008	1	14,14,15	0.54	0	17,19,21	1.52	2 (11%)
6	NAG	A	1007	1	14,14,15	0.61	0	17,19,21	1.00	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ASP	A	1019	-	-	3/8/8/8	-
6	NAG	A	1008	1	-	4/6/23/26	0/1/1/1
6	NAG	A	1007	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1019	ASP	OXT-C	-2.28	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1008	NAG	C2-N2-C7	4.25	128.59	122.90
7	A	1019	ASP	OXT-C-O	-3.11	117.03	124.08
6	A	1007	NAG	C4-C3-C2	2.33	114.43	111.02
6	A	1008	NAG	C1-C2-N2	2.18	113.87	110.43

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1008	NAG	C4-C5-C6-O6
6	A	1008	NAG	O5-C5-C6-O6
7	A	1019	ASP	C-CA-CB-CG
7	A	1019	ASP	N-CA-CB-CG
6	A	1008	NAG	C1-C2-N2-C7
6	A	1008	NAG	C3-C2-N2-C7
7	A	1019	ASP	OXT-C-CA-N

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1019	ASP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	875/888 (98%)	-0.61	1 (0%) 92 90	18, 58, 107, 155	2 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	615	ALA	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.70	0.12	102,161,196,205	0
3	NAG	D	3	14/15	0.79	0.13	123,160,173,182	0
3	NAG	E	3	14/15	0.79	0.13	126,160,177,189	0
2	NAG	B	2	14/15	0.82	0.10	136,154,174,175	0
2	NAG	G	2	14/15	0.83	0.12	110,137,156,163	0
2	NAG	F	2	14/15	0.89	0.10	120,134,157,166	0
3	NAG	D	2	14/15	0.89	0.17	112,143,157,159	0
3	NAG	D	1	14/15	0.93	0.10	79,106,121,122	0
2	NAG	C	1	14/15	0.93	0.10	81,111,127,131	0
2	NAG	B	1	14/15	0.94	0.08	74,103,128,144	0
2	NAG	F	1	14/15	0.94	0.09	66,90,112,115	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	E	2	14/15	0.95	0.10	102,120,140,144	0
3	NAG	E	1	14/15	0.96	0.08	70,91,105,108	0
2	NAG	G	1	14/15	0.97	0.07	58,83,106,115	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain G:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	A	1007	14/15	0.61	0.14	132,159,172,176	0
6	NAG	A	1008	14/15	0.84	0.12	131,156,172,183	0
7	ASP	A	1019	9/9	0.86	0.16	75,90,110,122	0
4	CA	A	1001	1/1	0.97	0.04	50,50,50,50	1
5	ZN	A	1002	1/1	0.98	0.10	103,103,103,103	0

6.5 Other polymers

There are no such residues in this entry.