



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:24 PM UTC

PDB ID : 6KXF / pdb_00006kxf
Title : The ishigamide ketosynthase/chain length factor
Authors : Du, D.; Katsuyama, Y.; Horiuchi, M.; Fushinobu, S.; Chen, A.; Davis, T.;
Burkart, M.; Ohnishi, Y.
Deposited on : 2019-09-10
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	409	2% 93% 6%
2	B	378	17% 89% 6% 5%
3	C	83	55% 49% 25% 10% 12%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6432 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ketosynthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	403	Total	C	N	O	S	0	0	0
			2953	1821	537	582	13			


- Molecule 2 is a protein called Ketosynthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	360	Total	C	N	O	S	0	0	0
			2572	1602	465	496	9			

- Molecule 3 is a protein called ACP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	73	Total	C	N	O	S	0	0	0
			553	347	88	115	3			

- Molecule 4 is [(3 {R})-2,2-dimethyl-4-[[3-[2-[({E})-oct-2-enoyl]amino]ethylamino]-3-oxidanylidene-propyl]amino]-3-oxidanyl-4-oxidanylidene-butyl] dihydrogen phosphate (CCD ID: DYF) (formula: C₁₉H₃₆N₃O₈P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	P	0	0
			31	19	3	8	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	216	Total	O	0	0
			216	216		
5	B	101	Total	O	0	0
			101	101		
5	C	6	Total	O	0	0
			6	6		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.87Å 104.41Å 114.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.56 – 1.98 47.56 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.56-1.98) 99.7 (47.56-1.98)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0230, PHENIX	Depositor
R, R_{free}	0.179 , 0.229 (Not available) , 0.214	Depositor DCC
R_{free} test set	3452 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6432	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DYF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	0/2999	1.06	3/4075 (0.1%)
2	B	0.92	0/2616	1.07	0/3570
3	C	0.84	1/557 (0.2%)	1.21	3/757 (0.4%)
All	All	0.96	1/6172 (0.0%)	1.08	6/8402 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	5
3	C	0	3
All	All	0	10

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	22	VAL	N-CA	5.32	1.53	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	PRO	N-CA-CB	-6.38	96.55	103.25
1	A	406	THR	CB-CA-C	6.34	119.99	109.53
3	C	34	ILE	CA-C-N	5.72	132.00	121.70
3	C	34	ILE	C-N-CA	5.72	132.00	121.70
1	A	359	PRO	N-CA-CB	5.46	106.06	103.22
1	A	108	THR	CB-CA-C	5.05	118.22	110.14

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	141	ARG	Sidechain
1	A	8	ARG	Sidechain
2	B	178	ARG	Sidechain
2	B	190	ARG	Sidechain
2	B	246	ARG	Sidechain
2	B	301	ARG	Sidechain
2	B	79	ARG	Sidechain
3	C	11	ARG	Sidechain
3	C	13	ARG	Sidechain
3	C	34	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2953	0	2898	13	0
2	B	2572	0	2566	10	0
3	C	553	0	538	25	0
4	C	31	0	0	0	0
5	A	216	0	0	3	0
5	B	101	0	0	0	0
5	C	6	0	0	1	0
All	All	6432	0	6002	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:34:ILE:HG12	3:C:68:THR:HA	1.53	0.88
3:C:34:ILE:HB	3:C:35:ASP:HB2	1.70	0.72
3:C:34:ILE:CB	3:C:35:ASP:HB2	2.23	0.69
3:C:26:GLY:C	3:C:69:ASN:OD1	2.37	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:360:VAL:HG12	2:B:366:VAL:HG22	1.77	0.66
2:B:242:GLU:CD	2:B:242:GLU:H	2.01	0.65
1:A:408:GLY:HA3	5:A:697:HOH:O	1.98	0.63
3:C:34:ILE:HD12	3:C:35:ASP:HB2	1.80	0.63
3:C:34:ILE:CG1	3:C:68:THR:HA	2.26	0.62
1:A:408:GLY:CA	5:A:697:HOH:O	2.48	0.61
2:B:146:ALA:HA	2:B:149:ILE:HD12	1.80	0.61
3:C:69:ASN:C	3:C:71:GLU:H	2.08	0.58
3:C:6:PRO:HG2	5:C:206:HOH:O	2.04	0.57
3:C:48:LEU:HD23	3:C:48:LEU:C	2.30	0.56
1:A:9:ARG:HD2	1:A:256:GLU:OE2	2.06	0.56
3:C:22:VAL:HG12	3:C:23:ASP:H	1.71	0.55
3:C:34:ILE:HD12	3:C:35:ASP:CB	2.36	0.55
3:C:69:ASN:O	3:C:73:THR:HG23	2.08	0.53
3:C:34:ILE:CD1	3:C:35:ASP:HB2	2.39	0.52
2:B:322:MET:HA	2:B:322:MET:HE2	1.91	0.52
1:A:268:HIS:ND1	1:A:269:PRO:HD2	2.25	0.52
1:A:108:THR:HA	1:A:164:THR:O	2.10	0.52
1:A:147:ARG:O	1:A:151:VAL:HG22	2.11	0.51
3:C:35:ASP:HA	3:C:39:ALA:CB	2.41	0.51
3:C:33:PHE:CB	3:C:69:ASN:ND2	2.73	0.51
2:B:239:GLU:OE1	2:B:246:ARG:NH2	2.44	0.50
1:A:13:THR:O	1:A:93:ALA:HA	2.12	0.50
3:C:35:ASP:N	3:C:39:ALA:HB2	2.26	0.49
3:C:40:ASP:OD2	3:C:42:LEU:HB2	2.12	0.48
3:C:34:ILE:CA	3:C:35:ASP:HB2	2.43	0.48
1:A:24:GLY:N	1:A:88:MET:HE1	2.30	0.47
2:B:275:ARG:NH2	2:B:279:ARG:HD3	2.29	0.47
3:C:33:PHE:CB	3:C:69:ASN:HD21	2.28	0.46
3:C:8:ARG:HG2	3:C:83:TRP:CH2	2.51	0.46
2:B:356:LEU:C	2:B:356:LEU:HD23	2.41	0.45
1:A:24:GLY:HA2	1:A:88:MET:CE	2.47	0.45
3:C:9:LEU:HD22	3:C:74:TYR:CD1	2.52	0.44
3:C:34:ILE:HG22	3:C:44:LEU:HD11	2.00	0.44
1:A:327:THR:O	1:A:375:VAL:HA	2.18	0.43
1:A:99:ASP:O	1:A:103:ARG:HD2	2.19	0.43
3:C:69:ASN:C	3:C:71:GLU:N	2.74	0.42
3:C:70:VAL:HG12	3:C:70:VAL:O	2.20	0.42
2:B:109:ALA:O	2:B:199:LYS:HE3	2.20	0.42
1:A:268:HIS:CE1	1:A:269:PRO:HD2	2.55	0.42
2:B:247:PRO:HA	2:B:248:PRO:HD2	1.81	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:5:ALA:HB3	3:C:6:PRO:HD3	2.03	0.41
2:B:117:VAL:HA	2:B:172:THR:O	2.21	0.41
1:A:408:GLY:HA2	5:A:697:HOH:O	2.18	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/409 (98%)	385 (96%)	16 (4%)	0	100	100
2	B	358/378 (95%)	343 (96%)	15 (4%)	0	100	100
3	C	69/83 (83%)	51 (74%)	12 (17%)	6 (9%)	0	0
All	All	828/870 (95%)	779 (94%)	43 (5%)	6 (1%)	18	9

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	6	PRO
3	C	35	ASP
3	C	37	LEU
3	C	40	ASP
3	C	22	VAL
3	C	25	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/304 (98%)	295 (99%)	4 (1%)	61	57
2	B	250/264 (95%)	243 (97%)	7 (3%)	38	29
3	C	58/70 (83%)	44 (76%)	14 (24%)	1	0
All	All	607/638 (95%)	582 (96%)	25 (4%)	27	16

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	ASP
1	A	108	THR
1	A	274	ARG
1	A	365	ARG
2	B	17	PRO
2	B	47	LYS
2	B	79	ARG
2	B	178	ARG
2	B	200	TYR
2	B	242	GLU
2	B	282	VAL
3	C	8	ARG
3	C	9	LEU
3	C	11	ARG
3	C	20	ILE
3	C	22	VAL
3	C	23	ASP
3	C	24	LEU
3	C	25	ASP
3	C	34	ILE
3	C	46	ASP
3	C	47	VAL
3	C	63	GLU
3	C	73	THR
3	C	75	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	73	GLN
2	B	75	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	336	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DYF	C	101	1,3	28,30,30	1.86	4 (14%)	36,39,39	4.41	12 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DYF	C	101	1,3	-	14/37/37/37	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	101	DYF	C21-N22	-4.74	1.35	1.46
4	C	101	DYF	C12-N14	4.53	1.44	1.33
4	C	101	DYF	C17-N19	4.28	1.43	1.33
4	C	101	DYF	C25-C23	3.31	1.55	1.48

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	101	DYF	O24-C23-N22	-16.90	95.78	122.29
4	C	101	DYF	C21-N22-C23	13.81	142.37	122.46
4	C	101	DYF	C25-C23-N22	10.46	138.00	115.07
4	C	101	DYF	C20-C21-N22	-5.83	93.40	111.54
4	C	101	DYF	C16-C17-N19	-4.15	108.78	116.34
4	C	101	DYF	O09-P06-O05	2.88	114.19	106.67
4	C	101	DYF	O11-C10-C02	-2.59	104.17	110.18
4	C	101	DYF	C20-N19-C17	2.53	127.53	122.82
4	C	101	DYF	C03-C02-C10	2.42	112.89	108.77
4	C	101	DYF	C26-C25-C23	-2.25	117.23	122.94
4	C	101	DYF	O18-C17-N19	2.14	127.23	123.03
4	C	101	DYF	O09-P06-O08	2.07	118.89	110.83

There are no chirality outliers.


All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	101	DYF	C25-C23-N22-C21
4	C	101	DYF	O24-C23-N22-C21
4	C	101	DYF	C04-O05-P06-O07
4	C	101	DYF	C04-O05-P06-O09
4	C	101	DYF	N22-C23-C25-C26
4	C	101	DYF	N19-C20-C21-N22
4	C	101	DYF	O24-C23-C25-C26
4	C	101	DYF	C16-C15-N14-C12
4	C	101	DYF	C04-O05-P06-O08
4	C	101	DYF	C27-C28-C29-C30
4	C	101	DYF	C15-C16-C17-O18
4	C	101	DYF	C15-C16-C17-N19
4	C	101	DYF	C25-C26-C27-C28
4	C	101	DYF	C02-C04-O05-P06

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	403/409 (98%)	0.32	7 (1%) 69 78	32, 41, 60, 79	0
2	B	360/378 (95%)	1.24	64 (17%) 4 4	37, 53, 83, 107	0
3	C	73/83 (87%)	2.69	46 (63%) 0 0	56, 89, 114, 120	0
All	All	836/870 (96%)	0.92	117 (13%) 6 9	32, 48, 89, 120	0

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	70	VAL	6.5
3	C	22	VAL	6.0
3	C	5	ALA	5.8
3	C	24	LEU	5.6
3	C	34	ILE	5.4
3	C	26	GLY	5.1
3	C	37	LEU	5.0
3	C	15	ILE	4.8
3	C	17	ALA	4.5
3	C	74	TYR	4.3
3	C	9	LEU	4.3
3	C	20	ILE	4.3
2	B	296	PRO	4.1
1	A	6	ALA	4.1
3	C	16	ILE	4.0
3	C	33	PHE	3.9
3	C	39	ALA	3.7
3	C	12	ILE	3.7
2	B	324	LEU	3.6
3	C	72	ALA	3.6
3	C	6	PRO	3.5
3	C	76	VAL	3.5
3	C	68	THR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	56	LEU	3.4
2	B	322	MET	3.4
3	C	78	ALA	3.4
3	C	83	TRP	3.4
2	B	102	ALA	3.3
2	B	18	GLY	3.3
2	B	348	ALA	3.2
3	C	69	ASN	3.2
2	B	297	THR	3.2
2	B	354	ILE	3.1
3	C	38	GLY	3.1
2	B	15	LEU	3.1
2	B	33	ILE	3.0
3	C	19	ASN	3.0
3	C	73	THR	3.0
2	B	374	ILE	3.0
3	C	10	SER	2.9
2	B	362	ARG	2.9
2	B	62	SER	2.9
2	B	16	PRO	2.9
2	B	49	ALA	2.9
2	B	346	ALA	2.8
2	B	17	PRO	2.8
1	A	371	CYS	2.7
2	B	31	TYR	2.7
2	B	343	ALA	2.7
3	C	81	ALA	2.7
1	A	408	GLY	2.7
3	C	75	GLN	2.6
2	B	36	ALA	2.6
2	B	319	VAL	2.6
2	B	48	THR	2.6
3	C	65	PRO	2.5
3	C	67	MET	2.5
2	B	57	GLY	2.5
2	B	344	ALA	2.5
2	B	316	LEU	2.5
2	B	352	GLY	2.5
3	C	13	ARG	2.5
2	B	64	VAL	2.5
3	C	23	ASP	2.5
2	B	50	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	44	ALA	2.4
2	B	109	ALA	2.4
2	B	294	ALA	2.4
2	B	32	GLY	2.4
3	C	49	SER	2.4
2	B	94	ALA	2.4
2	B	351	ARG	2.4
2	B	243	ALA	2.4
3	C	80	ALA	2.4
2	B	237	LEU	2.3
3	C	51	LEU	2.3
3	C	55	TYR	2.3
2	B	54	ALA	2.3
2	B	45	GLY	2.3
3	C	21	ASP	2.3
2	B	25	TRP	2.3
2	B	288	TRP	2.3
2	B	309	ALA	2.3
2	B	42	VAL	2.3
2	B	310	LEU	2.3
1	A	323	ALA	2.3
2	B	46	ALA	2.3
2	B	308	ALA	2.3
2	B	285	GLY	2.3
2	B	287	VAL	2.3
2	B	55	GLY	2.2
3	C	79	ALA	2.2
3	C	82	GLY	2.2
2	B	240	GLN	2.2
3	C	25	ASP	2.2
2	B	69	GLY	2.2
2	B	52	ALA	2.2
2	B	315	ALA	2.2
3	C	7	GLU	2.2
1	A	129	ALA	2.2
2	B	107	ALA	2.2
2	B	226	ALA	2.2
2	B	342	ALA	2.2
3	C	71	GLU	2.1
3	C	77	THR	2.1
1	A	130	GLY	2.1
2	B	373	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	53	MET	2.1
2	B	295	ALA	2.1
2	B	337	ILE	2.1
1	A	306	LYS	2.1
2	B	80	GLY	2.1
2	B	97	GLY	2.1
2	B	360	VAL	2.1
2	B	59	LEU	2.1
2	B	325	LEU	2.0
3	C	40	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	DYF	C	101	31/31	0.94	0.12	41,52,71,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.