



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 05:26 AM UTC

PDB ID : 1KYF / pdb_00001kyf
Title : AP-2 CLATHRIN ADAPTOR ALPHA-APPENDAGE IN COMPLEX WITH EPS15 DPF PEPTIDE
Authors : Brett, T.J.; Traub, L.M.; Fremont, D.H.
Deposited on : 2002-02-04
Resolution : 1.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2061 (1.24-1.20)
Ramachandran outliers	187476	2009 (1.24-1.20)
Sidechain outliers	187428	2008 (1.24-1.20)
RSRZ outliers	180081	2000 (1.24-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	4% 83% 13% ..
2	P	6	50% 33% 33% 17% 17%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4239 atoms, of which 1897 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-ADAPTIN C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	247	Total	C	H	N	O	S	0	0	0
			3854	1242	1897	336	370	9			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	692	GLY	-	cloning artifact	UNP P17427
A	693	SER	-	cloning artifact	UNP P17427
A	694	PRO	-	cloning artifact	UNP P17427
A	695	GLY	-	cloning artifact	UNP P17427
A	696	ILE	-	cloning artifact	UNP P17427
A	697	ARG	-	cloning artifact	UNP P17427
A	698	LEU	-	cloning artifact	UNP P17427
A	699	GLY	-	cloning artifact	UNP P17427
A	700	SER	-	cloning artifact	UNP P17427

- Molecule 2 is a protein called Epidermal growth factor receptor substrate 15.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	P	5	Total	C	N	O	0	0	0
			42	27	6	9			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	338	Total	O	0	0
			338	338		
3	P	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALPHA-ADAPTIN C



- Molecule 2: Epidermal growth factor receptor substrate 15



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.62Å 72.82Å 41.86Å 90.00° 99.95° 90.00°	Depositor
Resolution (Å)	20.00 – 1.22 20.00 – 1.22	Depositor EDS
% Data completeness (in resolution range)	94.4 (20.00-1.22) 87.1 (20.00-1.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.22Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.153 , 0.211 0.150 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.736	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.50 , 76.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.035 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4239	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	1/1994 (0.1%)	1.66	24/2697 (0.9%)
2	P	0.65	0/43	1.93	1/55 (1.8%)
All	All	0.81	1/2037 (0.0%)	1.66	25/2752 (0.9%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	707	ARG	NE-CZ	-5.18	1.27	1.33

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	707	ARG	CD-NE-CZ	36.95	176.12	124.40
1	A	806	ARG	CD-NE-CZ	16.32	147.25	124.40
1	A	761	ASP	CA-CB-CG	8.71	121.31	112.60
1	A	849	GLU	CB-CG-CD	8.09	126.36	112.60
1	A	751	ASN	CA-CB-CG	7.33	119.93	112.60
1	A	844	SER	O-C-N	7.32	131.10	122.09
1	A	773	VAL	O-C-N	-6.85	114.88	122.69
1	A	760	ASP	CA-C-O	-6.72	113.68	121.07
1	A	858	HIS	O-C-N	6.49	125.87	121.71
1	A	813	ASN	OD1-CG-ND2	6.33	128.94	122.60
1	A	825	PHE	CA-CB-CG	-6.26	107.54	113.80
1	A	845	ASN	CA-CB-CG	6.12	118.72	112.60
1	A	837	PHE	CA-CB-CG	-5.90	107.90	113.80
1	A	924	ASP	CA-CB-CG	5.89	118.49	112.60
1	A	705	PHE	O-C-N	5.78	128.11	122.09
1	A	704	ASN	CA-CB-CG	-5.75	106.85	112.60
1	A	773	VAL	CB-CA-C	5.68	119.88	111.31
1	A	913	GLN	OE1-CD-NE2	5.68	128.28	122.60
1	A	849	GLU	CB-CA-C	-5.54	101.10	110.19
1	A	843	LEU	CA-C-N	-5.34	112.85	122.46
1	A	843	LEU	C-N-CA	-5.34	112.85	122.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	872	PHE	CA-CB-CG	5.26	119.06	113.80
2	P	631	PHE	O-C-N	5.09	128.89	122.37
1	A	740	PHE	CA-CB-CG	-5.07	108.73	113.80
1	A	833	SER	O-C-N	-5.04	116.78	122.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1957	1897	1953	16	0
2	P	42	0	37	2	0
3	A	338	0	0	9	0
3	P	5	0	0	0	0
All	All	2342	1897	1990	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:936:GLU:HG3	3:A:252:HOH:O	1.90	0.71
1:A:744:LYS:HE2	3:A:275:HOH:O	1.95	0.67
1:A:736:ARG:HD2	3:A:172:HOH:O	1.97	0.63
2:P:631:PHE:O	2:P:632:LYS:HB3	2.06	0.56
1:A:774:ASP:HB3	3:A:81:HOH:O	2.10	0.51
1:A:712:ASN:C	1:A:713:ASN:HD22	2.18	0.51
1:A:783:VAL:HG23	3:A:72:HOH:O	2.13	0.49
1:A:848:GLN:HB3	1:A:921:THR:O	2.16	0.46
2:P:632:LYS:HB3	2:P:632:LYS:HE2	1.37	0.46
1:A:834:GLN:NE2	1:A:838:GLN:OE1	2.49	0.45
1:A:806:ARG:NE	3:A:287:HOH:O	2.50	0.45
1:A:834:GLN:O	1:A:838:GLN:HG3	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:813:ASN:ND2	3:A:194:HOH:O	2.50	0.45
1:A:761:ASP:OD1	1:A:765:ASN:ND2	2.52	0.43
1:A:832:ALA:HA	3:A:276:HOH:O	2.20	0.41
1:A:693:SER:O	1:A:693:SER:OG	2.30	0.41
1:A:697:ARG:HG3	3:A:347:HOH:O	2.21	0.41
1:A:713:ASN:ND2	1:A:727:LYS:HA	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/247 (99%)	241 (98%)	3 (1%)	1 (0%)	30	10
2	P	3/6 (50%)	3 (100%)	0	0	100	100
All	All	248/253 (98%)	244 (98%)	3 (1%)	1 (0%)	30	10

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	693	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/220 (100%)	208 (94%)	12 (6%)	19	1
2	P	5/5 (100%)	4 (80%)	1 (20%)	1	0
All	All	225/225 (100%)	212 (94%)	13 (6%)	18	1

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	693	SER
1	A	697	ARG
1	A	761	ASP
1	A	769	GLN
1	A	773	VAL
1	A	792	ILE
1	A	806	ARG
1	A	813	ASN
1	A	834	GLN
1	A	844	SER
1	A	864	ILE
1	A	878	GLU
2	P	628	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	713	ASN
1	A	767	ASN
1	A	802	ASN
1	A	804	GLN
1	A	813	ASN
1	A	834	GLN
1	A	845	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/247 (100%)	-0.02	11 (4%) 38 40	14, 23, 47, 86	0
2	P	5/6 (83%)	2.10	3 (60%) 0 0	29, 31, 53, 62	0
All	All	252/253 (99%)	0.03	14 (5%) 30 32	14, 23, 50, 86	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	692	GLY	4.9
1	A	693	SER	4.1
2	P	632	LYS	3.4
2	P	628	SER	3.1
1	A	761	ASP	2.7
1	A	844	SER	2.7
1	A	764	THR	2.6
1	A	760	ASP	2.6
1	A	772	PRO	2.6
1	A	845	ASN	2.6
2	P	631	PHE	2.4
1	A	834	GLN	2.3
1	A	694	PRO	2.2
1	A	846	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.